

Bilaga till rapport

1 (1)

Diagnos och behandling av epilepsi / Diagnosis
and treatment of epilepsy, rapport [281] (2018)

Bilaga 2 - Exkluderade studier samt flödesträd över sökning och urval av litteratur

Innehåll

Inledning.....	2
Diagnos	2
EEG inom 12 – 24 timmar (7)	2
Sömn-EEG (23).....	2
Video-EEG (60).....	3
MRT-DT (147)	6
Läkemedelsbehandling.....	12
Insättning (9)	12
Utsättning (15).....	12
Koncentrationsbestämning (27).....	13
Tilläggsbehandling – fokala epilepsi (105).....	14
Tilläggsbehandling – epilepsi generaliserade tonisk-kloniska anfall (69)	19
Anfallskuperande läkemedel (5).....	22
Anti-depressiva läkemedel (79).....	22
Anti-psykotiska läkemedel (19)	25
Centralstimulerande läkemedel (35).....	26
Kirurgis behandling.....	28
Resekтив kirurgi (33).....	28
Resekтив kirurgi tidsfaktor (62)	29
Annan behandling	32
Ketogen kost (121)	32
DBS (27)	37
VNS (121).....	38
Hälsoekonomi.....	43
Flödesträd över litteraturgenomgången.....	47

Inledning

I denna bilaga listas de studier som exkluderats efter att ha granskats i fulltext. Att en artikel exkluderats betyder att den inte uppfyller ett eller flera urvalskriterier definierade i respektive kapitel. En studie kan även ha exkluderats om den inte uppfyller basala krav på vetenskaplig tillförlitlighet. Ibland kan det vara så att fler eller samtliga urvalskriterier inte är uppfyllda. En och samma artikel kan förekomma på flera ställen eftersom det kan preliminärt ha bedömts vara relevant för flera åtgärder. Det kan även förekomma artiklar som exkluderats för vissa åtgärder i den första urvalsprocessen med stöd av information tillgänglig i de elektroniska referenserna. Urvalet här är med andra ord sensitivt.

Diagnos

EEG inom 12 – 24 timmar (7)

- Bouma, H. K., Labos, C., Gore, G. C., Wolfson, C., & Keezer, M. R. (2016). The diagnostic accuracy of routine electroencephalography after a first unprovoked seizure. *European Journal of Neurology*, 23(3), 455-463.
- Hamiwka LD, Singh N, Niosi J, Wirrell EC. Diagnostic inaccuracy in children referred with "first seizure": role for a first seizure clinic. *Epilepsia* 2007; 48:1062–6.
- Hamiwka, L., Singh, N., Kozlik, S., & Wirrell, E. (2008). Feasibility and clinical utility of early electroencephalogram (EEG) in children with first seizure. *Journal of Child Neurology*, 23(7), 762–765.
- Maillard L, Jonas J, Boyer R, Frismand S, Mathey G, et al. One-year outcome after a first clinically possible epileptic seizure: Predictive value of clinical classification and early EEG. *Neurophysiologie Clinique/Clinical Neurophysiology*, Elsevier Masson, 2012, 42 (6), pp.355–362.
- Sierra-Marcos, A., Toledo, M., Quintana, M., Edo, M. C., Centeno, M., Santamarina, E., Alvarez-Sabin, J. (2011). Diagnosis of epileptic syndrome after a new onset seizure and its correlation at long-term follow-up: longitudinal study of 131 patients from the emergency room. *Epilepsy Research*, 97(1–2), 30–36.
- Wikström S, Hansen Pupp I, Rosén I, Norman E, Fellman V, Ley D, Hellström-Westas L. Early single-channel aEEG/EEG predicts outcome in very preterm infants. *Acta Paediatr*. 2012 Jul; 101(7): 719–726.
- Yigit, O., Eray, O., Mihci, E., Yilmaz, D., Eray, B., & Ozkaynak, S. (2013). EEG as a part of the decision-making process in the emergency department. *European Journal of Emergency Medicine*, 20(6), 402–407.

Sömn-EEG (23)

- Borkowski WJ Jr, Ellington RJ, Sverdrup EK. Effect of sleep deprivation on the EEG of learning-impaired children with absence seizures. *Clin Electroencephalogr*. 1992 Apr;23(2):62–4.
- Burkholder DB, Britton JW, Rajasekaran V, Fabris RR, Cherian PJ, Kelly-Williams KM, So EL, Nickels KC, Wong-Kisiel LC, Lagerlund TD, Cascino GD, Worrell GA, Wirrell EC. *Neurology*. 2016 Apr 19;86(16):1524–30.
- Carpay JA1, de Weerd AW, Schimsheimer RJ, Stroink H, Brouwer OF, Peters AC, van Donselaar CA, Geerts AT, Arts WF. The diagnostic yield of a second EEG after partial sleep deprivation: a prospective study in children with newly diagnosed seizures. *Epilepsia*. 1997 May;38(5):595–9.
- de Carvalho Sousal NA, da Silva Sousal P, Garzonl E, Sakamotol AC, Bragall NIO, Yacubian EMT. EEG recording after sleep deprivation in a series of patients with juvenile myoclonic epilepsy. *Arq Neuropsiquiatr* 2005;63(2-B):383–388
- Deisenhammer E, Klingler D, Trägner H. Epileptic seizures in alcoholism and diagnostic value of EEG after sleep deprivation. *Epilepsia*. 1984 Aug;25(4):526–30.
- Gandelman-Marton R, Theitler J. When should a sleep-deprived EEG be performed following a presumed first seizure in adults? *Acta Neurol Scand*. 2011 Sep;124(3):202–5.
- Geut, I., Weenink, S., Knottnerus, I. L. H., & van Putten, M. (2017). Detecting interictal discharges in first seizure patients: ambulatory EEG or EEG after sleep deprivation? *Seizure*, 51, 52–54.
- Gilbert DL, DeRoos S, Bare MA. Does sleep or sleep deprivation increase epileptiform discharges in pediatric electroencephalograms? *Pediatrics*. 2004 Sep;114(3):658–62.
- Giorgi, F. S., Maestri, M., Guida, M., Carnicelli, L., Caciagli, L., Ferri, R., Bonanni, E. (2017). Cyclic alternating pattern and interictal epileptiform discharges during morning sleep after sleep deprivation in temporal lobe epilepsy. *Epilepsy & Behavior*, 73, 131–136.
- Giorgi FS, Perini D, Maestri M, Guida M, Pizzanelli C, Caserta A, Iudice A, Bonanni E. Usefulness of a simple sleep-deprived EEG protocol for epilepsy diagnosis in de novo subjects. *Clin Neurophysiol*. 2013

- Nov;124(11):2101–7.
- King MA, Newton MR, Jackson GD, Fitt GJ, Mitchell LA, Silvapulle MJ, Berkovic SF. Epileptology of the first-seizure presentation: a clinical, electroencephalographic, and magnetic resonance imaging study of 300 consecutive patients. *Lancet*. 1998 Sep 26;352(9133):1007–11.
- Lawley A, Evans S, Manfredonia F, Cavanna AE. The role of outpatient ambulatory electroencephalography in the diagnosis and management of adults with epilepsy or nonepileptic attack disorder: A systematic literature review. *Epilepsy Behav*. 2015 Dec;53:26–30.
- Liporace J, Tatum W 4th, Morris GL 3rd, French J. Clinical utility of sleep-deprived versus computer-assisted ambulatory 16-channel EEG in epilepsy patients: a multi-center study. *Epilepsy Res*. 1998 Nov;32(3):357–62.
- Rafiei SM. Usefulness of sleep-deprived EEG in the diagnosis of seizure disorders in children. *Medical Journal of the Islamic Republic of Iran*. 18(1), p. 21–7.
- Renzel R, Baumann CR, Poryazova R. EEG after sleep deprivation is a sensitive tool in the first diagnosis of idiopathic generalized but not focal epilepsy. *Clin Neurophysiol*. 2016 Jan;127(1):209–213.
- Roupakiotis SC, Gatzonis SD, Triantafyllou N, Mantouvalos V, Chioni A, Zournas C, Siafakas A. The usefulness of sleep and sleep deprivation as activating methods in electroencephalographic recording: contribution to a long-standing discussion. *Seizure*. 2000 Dec;9(8):580–4.
- Schwarz JR, Zangemeister WH. The diagnostic value of the short sleep EEG and other provocative methods following sleep deprivation. *Journal of Neurology*, September 1978, Volume 218, Issue 3, pp 179–186
- Shahar E, Genizi J, Ravid S, Schif A. The complementary value of sleep-deprived EEG in childhood onset epilepsy. *Eur J Paediatr Neurol*. 2010 Jul;14(4):308–12.
- Tartara A, Moglia A, Manni R, Corbellini C. EEG findings and sleep deprivation. *Eur Neurol*. 1980;19(5):330–4.
- Theitler, J., Dassa, D., & Gandelman-Marton, R. (2016). How valuable is inpatient electroencephalogram for medical decision-making? *European Journal of Clinical Investigation*, 46(12), 1012–16.
- van Donselaar CA1, Geerts AT, Schimsheimer RJ. Idiopathic first seizure in adult life: who should be treated? *BMJ*. 1991 Mar 16;302(6777):620–3.
- Vardi J, Lipschitz M, Rosenberg P, Zevin S, Flechter S. EEG recording following sleep deprivation. A study utilizing the ‘source derivation’ technique. *Electroencephalography and Clinical Neurophysiology*. 66(5), 1987, Page S127.
- Veldhuizen R, Binnie CD, Beintema DJ. The effect of sleep deprivation on the EEG in epilepsy. *Electroencephalogr Clin Neurophysiol*. 1983 May;55(5):505–12.

Video-EEG (60)

- Alsaadi, T. M., Thieman, C., Shatzel, A., & Farias, S. (2004). Video-EEG telemetry can be a crucial tool for neurologists experienced in epilepsy when diagnosing seizure disorders. *Seizure*, 13(1), 32–34.
- Anonymous. (1983). Diagnostic and therapeutic technology assessment. 24-hour ambulatory EEG monitoring. *JAMA*, 250(24), 3340.
- Apakama, O., & Appleton, R. (2006). Non-epileptic clinical diagnoses in children referred for an outpatient EEG using video monitoring. *Epileptic Disorders*, 8(2), 156–58.
- Baxter, P. (2006). Diagnosing epilepsy. *Developmental Medicine & Child Neurology*, 48(5), 323.
- Benbadis, S. R., Siegrist, K., Tatum, W. O., Heriaud, L., & Anthony, K. (2004). Short-term outpatient EEG video with induction in the diagnosis of psychogenic seizures. *Neurology*, 63(9), 1728–30.
- Bergey, G. K. (2001). Evaluation of the patient with epilepsy. *American Journal of Managed Care*, 7(7 Suppl), S203–08.
- Binnie, C. D. (1987). Ambulatory diagnostic monitoring of seizures in adults. *Advances in Neurology*, 46, 169–82.
- Binnie, C. D., Aarts, J. H., Van Bentum-De Boer, P. T., & Wisman, T. (1985). Monitoring at the Instituut voor Epilepsiebestrijding Meer en Bosch. *Electroencephalography & Clinical Neurophysiology - Supplement*, 37, 341–55.
- Binnie, C. D., Rowan, A. J., & Overweg, J. (1981). Telemetric EEG and video monitoring in epilepsy. *Neurology*, 31(3), 298–303.
- Bleasel, A., Kotagal, P., Kankirawatana, P., & Rybicki, L. (1997). Lateralizing value and semiology of ictal limb posturing and version in temporal lobe and extratemporal epilepsy. *Epilepsia*, 38(2), 168–74.
- Bodde, N. M., Brooks, J. L., Baker, G. A., Boon, P. A., Hendriksen, J. G., & Aldenkamp, A. P. (2009). Psychogenic non-epileptic seizures--diagnostic issues: a critical review. *Clinical Neurology & Neurosurgery*, 111(1), 1–9.
- Boon, P., Vandekerckhove, T., Achten, E., Thiery, E., Goossens, L., Vonck, K., De Reuck, J. (1999). Epilepsy

- surgery in Belgium, the experience in Gent. *Acta Neurologica Belgica*, 99(4), 256-265.
- Bowman, E. S., & Coons, P. M. (2000). The differential diagnosis of epilepsy, pseudoseizures, dissociative identity disorder, and dissociative disorder not otherwise specified. *Bulletin of the Menninger Clinic*, 64(2), 164-180.
- Bridgers, S. L., & Ebersole, J. S. (1985). The clinical utility of ambulatory cassette EEG. *Neurology*, 35(2), 166-173.
- Bye, A. M. E., Kok, D. J. M., Ferenschild, F. T. J., & Vles, J. S. H. (2000). Paroxysmal non-epileptic events in children: A retrospective study over a period of 10 years. *Journal of Paediatrics and Child Health*, 36(3), 244-248.
- Carlson, C., Arnedo, V., Cahill, M., & Devinsky, O. (2009). Detecting nocturnal convulsions: efficacy of the MP5 monitor. *Seizure*, 18(3), 225-227.
- Cavanna, A. E., & Seri, S. (2016). Neurophysiological investigations for the diagnosis of non-epileptic attack disorder in neuropsychiatry services: From safety standards to improved effectiveness. *Acta Neuropsychiatrica*, 28(4), 185-194.
- Chayasirisobhon, S., Griggs, L., Westmoreland, S., & Kim, C. S. (1993). The usefulness of one to two hour video EEG monitoring in patients with refractory seizures. *Clinical Electroencephalography*, 24(2), 78-84.
- Chen, L. S., Mitchell, W. G., Horton, E. J., & Snead, O. C., 3rd. (1995). Clinical utility of video-EEG monitoring. *Pediatric Neurology*, 12(3), 220-224.
- Cuthill, F. M., & Espie, C. A. (2005). Sensitivity and specificity of procedures for the differential diagnosis of epileptic and non-epileptic seizures: a systematic review. *Seizure*, 14(5), 293-303.
- Davis, B. J. (2004). Predicting nonepileptic seizures utilizing seizure frequency, EEG, and response to medication. *European Neurology*, 51(3), 153-156.
- Del Giudice, E., Crisanti, A. F., & Romano, A. (2002). Short duration outpatient video electroencephalographic monitoring: the experience of a southern-Italian general pediatric department. *Epileptic Disorders*, 4(3), 197-202.
- Ebersole, J. S. (1985). Ambulatory cassette EEG. *Journal of Clinical Neurophysiology*, 2(4), 397-418.
- Ebersole, J. S. (1987). Ambulatory cassette EEG in epilepsy diagnosis. *Yale Journal of Biology & Medicine*, 60(2), 85-91.
- Ebersole, J. S., & Bridgers, S. L. (1985). Direct comparison of 3- and 8-channel ambulatory cassette EEG with intensive inpatient monitoring. *Neurology*, 35(6), 846-854.
- Ebersole, J. S., & Leroy, R. F. (1983). Evaluation of ambulatory cassette EEG monitoring: III. Diagnostic accuracy compared to intensive inpatient EEG monitoring. *Neurology*, 33(7), 853-860.
- Ebersole, J. S., Mattson, R. H., Williamson, P. D., Spencer, S. S., & Glaser, G. H. (1985). Monitoring at the West Haven VA/Yale University School of Medicine Epilepsy Center. *Electroencephalography & Clinical Neurophysiology - Supplement*, 37, 357-369.
- Engelsen, B. A., Gramstad, A., Lillebø, A., & Karlsen, B. (2013). An update on non-epileptic seizures: Diagnosis and brief notes on treatment. *Translational Neuroscience*, 4(2), 260-268.
- Faulkner, H. J., Arima, H., & Mohamed, A. (2012). The utility of prolonged outpatient ambulatory EEG. *Seizure*, 21(7), 491-495.
- Fois, A., Farnetani, M. A., Balestri, P., Buoni, S., Di Cosmo, G., Vattimo, A., Salvadori, P. A. (1995). EEG, PET, SPET and MRI in intractable childhood epilepsies: Possible surgical correlations. *Child's Nervous System*, 11(12), 672-678.
- Foley, C. M., Legido, A., Miles, D. K., & Grover, W. D. (1995). Diagnostic value of pediatric outpatient video-EEG. *Pediatric Neurology*, 12(2), 120-124.
- Gandelman-Marton, R., Kipervasser, S., & Neufeld, M. Y. (2012). Long-term video-EEG in patients with frontal seizures. *Neurological Research*, 34(10), 957-959.
- Güldiken, B., Baykan, B., Süt, N., Bebek, N., Gürses, C., & Gökyiğit, A. (2012). The evaluation of the agreements of different epilepsy classifications in seizures recorded with video EEG monitoring. *Journal of Neurological Sciences*, 29(2), 201-211.
- Hayes, & Inc. (2007). Video electroencephalograph (VEEG) monitoring for diagnosis of epilepsy in children (Structured abstract). *Health Technology Assessment Database*, (4).
- Hayes, & Inc. (2013). Video Electroencephalogram (VEEG) for diagnosis and management of epilepsy in adults (Structured abstract). *Health Technology Assessment Database*, (4).
- Heldberg, B. E., Kautz, T., Leutheuser, H., Hopfengartner, R., Kasper, B. S., & Eskofier, B. M. (2015). Using wearable sensors for semiology-independent seizure detection - towards ambulatory monitoring of epilepsy. *Conference Proceedings: ... Annual International Conference of the IEEE Engineering in Medicine & Biology Society, 2015*, 5593-5596.
- Holmes, G. L., McKeever, M., & Russman, B. S. (1982). Prolonged EEG and videotape monitoring in children.

- American Journal of Diseases of Children*, 136(7), 609-611.
- Kader, A. A. A., El-Kattan, M., Mostafa, S., El-Gohary, A., Ramzy, G., & Nawito, A. (2009). Long term polygraphic study of paroxysmal events in infancy and their relationship to sleep. *Egyptian Journal of Neurology, Psychiatry and Neurosurgery*, 46(2), 409-420.
- Lawley, A., Evans, S., Manfredonia, F., & Cavanna, A. E. (2015). The role of outpatient ambulatory electroencephalography in the diagnosis and management of adults with epilepsy or nonepileptic attack disorder: A systematic literature review. *Epilepsy & Behavior*, 53, 26-30.
- Lawley, A., Manfredonia, F., & Cavanna, A. E. (2016). Video-ambulatory EEG in a secondary care center: A retrospective evaluation of utility in the diagnosis of epileptic and nonepileptic seizures. *Epilepsy & Behavior*, 57(Pt A), 137-140.
- Mizrahi, E. M. (1984). Electroencephalographic/polygraphic/video monitoring in childhood epilepsy. *Journal of Pediatrics*, 105(1), 1-9.
- Nuwer, M. R., Engel, J., Jr., Sutherling, W. W., & Babb, T. L. (1985). Monitoring at the University of California, Los Angeles. *Electroencephalography & Clinical Neurophysiology - Supplement*, 37, 385-401.
- Olejniczak, P. W., Carey, M. C., Fisch, B. J., Trouppin, A. S., Tardo, C., & Hauck, M. (1996). The Louisiana State University Comprehensive Epilepsy Program: procedures and outcomes. *Journal of the Louisiana State Medical Society*, 148(12), 525-532.
- Otsubo, H., Hwang, P. A., Jay, V., Becker, L. E., Hoffman, H. J., Gilday, D., & Blaser, S. (1993). Focal cortical dysplasia in children with localization-related epilepsy: EEG, MRI, and SPECT findings. *Pediatric Neurology*, 9(2), 101-107.
- Pataria, E., Lurker, S., Serles, W., Lindinger, G., Aull, S., Leutmezer, F., Baumgartner, C. (1998). Ictal scalp EEG in unilateral mesial temporal lobe epilepsy.[Erratum appears in Epilepsia 1998 Sep;39(9):1022-3]. *Epilepsia*, 39(6), 608-614.
- Perkins, A. M., & Buchhalter, J. R. (2006). Optimizing patient care in the pediatric epilepsy monitoring unit. *Journal of Neuroscience Nursing*, 38(6), 416-421, 434.
- Perry, T. R., Gumnit, R. J., Gates, J. R., & Leppik, I. E. (1983). Routine EEG vs. intensive monitoring in the evaluation of intractable epilepsy. *Public Health Reports*, 98(4), 384-389.
- Pfander, M., Arnold, S., Henkel, A., Weil, S., Werhahn, K. J., Eisensehr, I., Noachtar, S. (2002). Clinical features and EEG findings differentiating mesial from neocortical temporal lobe epilepsy. *Epileptic Disorders*, 4(3), 189-195.
- Pritchard 3rd, P. B. (1992). Intensive neurodiagnostic monitoring. *Journal of the South Carolina Medical Association* (1975), 88(5), 230-233.
- Robinson, A. A., Pitiyanuvath, N., Abou-Khalil, B. W., Wang, L., Shi, Y., & Azar, N. J. (2011). Predictors of a nondiagnostic epilepsy monitoring study and yield of repeat study. *Epilepsy & Behavior*, 21(1), 76-79.
- Seneviratne, U., Rahman, Z., Diamond, A., & Brusco, M. (2012). The yield and clinical utility of outpatient short-term video-electroencephalographic monitoring: a five-year retrospective study. *Epilepsy & Behavior*, 25(3), 303-306.
- Shukla, G., Bhatia, M., Gaekwad, S. B., Singh, V. P., Jain, S., & Maheshwari, M. C. (2002). The lateralizing significance of version of head and dystonic limb posturing in epileptic seizures. *Neurology India*, 50(1), 33-36.
- Si, Y., Liu, L., Fang, J. J., Mu, J., Hu, J., Zhao, L. L., Zhou, D. (2011). Evaluation of the efficiency of inpatient 24-hour VEEG combined with MRI in consecutive patients with newly diagnosed epilepsies. *Epilepsy & Behavior*, 20(4), 633-637.
- Stores, G. (1987). Ambulatory diagnostic monitoring of seizures in children. *Advances in Neurology*, 46, 157-167.
- Troester, M., Haine-Schlagel, R., Ng, Y. T., Chapman, K., Chung, S., Drees, C., Kerrigan, J. F. (2011). EEG and video-EEG seizure monitoring has limited utility in patients with hypothalamic hamartoma and epilepsy. *Epilepsia*, 52(6), 1137-1143.
- Tsuchida, T. N. (2015). Diagnostic Yield of Continuous Video EEG for Neonatal Seizures. *Pediatric Neurology Briefs*, 29(8), 63.
- Tuunainen, A., & Nousianinen, U. (1993). Ictal recordings of ambulatory cassette EEG with sphenoidal electrodes in temporal lobe epilepsy: comparison with intensive videomonitoring. *Acta Neurologica Scandinavica*, 88(1), 21-25.
- Van den Noort, S., Economy, J., Davis, E., Dyck, P., Greenberg, J., Jacobs, L., Stuart, W. (1989). Assessment: Intensive EEG/video monitoring for epilepsy. *Neurology*, 39(8), 1101-1102.
- Watemberg, N., Lerman-Sagie, T., & Kramer, U. (2008). Diagnostic yield of electroencephalograms in infants and young children with frequent paroxysmal eye movements. *Journal of Child Neurology*, 23(6), 620-623.

Watemberg, N., Tziperman, B., Dabby, R., Hasan, M., Zehavi, L., & Lerman-Sagie, T. (2005). Adding video recording increases the diagnostic yield of routine electroencephalograms in children with frequent paroxysmal events. *Epilepsia*, 46(5), 716-719.

MRT-DT (147)

- Adams, C., Hwang, P. A., Gilday, D. L., Armstrong, D. C., Becker, L. E., & Hoffman, H. J. (1992). Comparison of SPECT, EEG, CT, MRI, and pathology in partial epilepsy. *Pediatric Neurology*, 8(2), 97-103.
- Ahmedani, B. K., Osborne, J., Nerenz, D. R., Haque, S., Pietrantoni, L., Mahone, D., & Smith, B. J. (2013). Diagnosis, Costs, and Utilization for Psychogenic Non-Epileptic Seizures in a US Health Care Setting. *Psychosomatics*, 54(1), 28-34.
- Alam-Eldeen, M. H., & Hasan, N. M. A. (2015). Assessment of the diagnostic reliability of brain CT and MRI in pediatric epilepsy patients. *Egyptian Journal of Radiology and Nuclear Medicine*.
- Alshami, R., Bessisso, M., El Said, M. F., & Venktraman, B. (2011). Neuro-images of patients with infantile spasm. *Qatar Medical Journal*, 20(2), 32-33.
- Andersen, A. R., Gram, L., Kjaer, L., Fuglsang-Frederiksen, A., Herning, M., Lassen, N. A., & Dam, M. (1988). SPECT in partial epilepsy: identifying side of the focus. *Acta Neurologica Scandinavica. Supplementum*, 117, 90-95.
- Anslow, P. (1990). Neuroradiology including magnetic resonance. *Acta Neurochirurgica - Supplementum*, 50, 76-79.
- Appel, B., Engelen, H., Moens, E., Mortelmans, L., & Lowenthal, A. (1986). Clinical evaluation of the first 365 patients studied with a 0.15 tesla resistive MRI system. *Acta neurologica Belgica*, 86(1), 5-10.
- Aprahamian, N., Harper, M. B., Prabhu, S. P., Monuteaux, M. C., Sadiq, Z., Torres, A., & Kimia, A. A. (2014). Pediatric first time non-febrile seizure with focal manifestations: Is emergent imaging indicated? *Seizure*, 23(9), 740-745.
- Arroyo, H. A., De Rosa, S., Ruggieri, V., de Dávila, M. T. G., Fejerman, N., Aldao, M., Toca, M. (2002). Epilepsy, occipital calcifications, and oligosymptomatic celiac disease in childhood. *Journal of Child Neurology*, 17(11), 800-806.
- Aslan, K., Bozdemir, H., Yapar, Z., & Burgut, R. (2010). The effect of electrophysiological and neuroimaging findings on the prognosis of juvenile myoclonic epilepsy proband. *Neurological Research*, 32(6), 620-624.
- Baker, H. L., Jr., Berquist, T. H., Kispert, D. B., Reese, D. F., Houser, O. W., Earnest, F. t., May, G. R. (1985). Magnetic resonance imaging in a routine clinical setting. *Mayo Clinic Proceedings*, 60(2), 75-90.
- Bakhsh, A. (2013). Value of neuroimaging in epilepsy: An experience from Pakistan. *Journal of Neurosciences in Rural Practice*, 4(5 SUPPL), S35-S39.
- Barkovich, A. J., Good, W. V., Koch, T. K., & Berg, B. O. (1993). Mitochondrial disorders: analysis of their clinical and imaging characteristics. *Ajnr: American Journal of Neuroradiology*, 14(5), 1119-1137.
- Bergen, D., Bleck, T., Ramsey, R., Clasen, R., Ristanovic, R., Smith, M., & Whisler, W. W. (1989). Magnetic resonance imaging as a sensitive and specific predictor of neoplasms removed for intractable epilepsy. *Epilepsia*, 30(3), 318-321.
- Bonilha, L., Montenegro, M. A., Cendes, F., & Li, L. M. (2004). The role of neuroimaging in the investigation of patients with single seizures, febrile seizures, or refractory partial seizures. *Medical Science Monitor*, 10(3), RA40-46.
- Boon, P. A., Williamson, P. D., Fried, I., Spencer, D. D., Novelly, R. A., Spencer, S. S., & Mattson, R. H. (1991). Intracranial, intraaxial, space-occupying lesions in patients with intractable partial seizures: An anatomic-clinical, neuropsychological, and surgical correlation. *Epilepsia*, 32(4), 467-476.
- Bozdemir, H., Sarica, Y., Tiryaki, G., Demirkiran, M., Kibar, M., & Bicakci, K. (1999). EEG, MRI and SPECT findings in partial epilepsies. *Annals of Medical Sciences*, 8(1), 26-30.
- Bronen, R. A., Fulbright, R. K., King, D., Kim, J. H., Spencer, S. S., Spencer, D. D., & Lange, R. (1997). Qualitative MR imaging of refractory temporal lobe epilepsy requiring surgery: Correlation with pathology and seizure outcome after surgery. *American Journal of Roentgenology*, 169(3), 875-882.
- Bronen, R. A., Fulbright, R. K., Spencer, D. D., Spencer, S. S., Kim, J. H., Lange, R. C., & Sutilla, C. (1996). Refractory epilepsy: comparison of MR imaging, CT, and histopathologic findings in 117 patients. *Radiology*, 201(1), 97-105.
- Camacho, D. L. A., & Castillo, M. (2007). MR Imaging of Temporal Lobe Epilepsy. *Seminars in Ultrasound, CT and MRI*, 28(6), 424-436.
- Carrilho, P. G., Yacubian, E. M., Cukiert, A., Fiore, L. A., Buchpiguel, C. A., Jorge, C. L., Marino Junior, R. (1994). MRI and brain spect findings in patients with unilateral temporal lobe epilepsy and normal CT scan.

- Arquivos de Neuro-Psiquiatria*, 52(2), 149-152.
- Cendes, F., Leproux, F., Melanson, D., Ethier, R., Evans, A., Peters, T., & Andermann, F. (1993). MRI of amygdala and hippocampus in temporal lobe epilepsy. *Journal of Computer Assisted Tomography*, 17(2), 206-210.
- Dayan, P. S., Lillis, K., Bennett, J., Conners, G., Bailey, P., Callahan, J., Kuppermann, N. (2015). Prevalence of and Risk Factors for Intracranial Abnormalities in Unprovoked Seizures. *Pediatrics*, 136(2), e351-360.
- Debets, R. M., van Veelen, C. W., Maquet, P., van Huffelen, A. C., van Emde Boas, W., Sadzot, B., Franck, G. (1990). Quantitative analysis of 18/FDG-PET in the presurgical evaluation of patients suffering from refractory partial epilepsy. Comparison with CT, MRI, and combined subdural and depth EEG. *Acta Neurochirurgica - Supplementum*, 50, 88-94.
- Del Brutto, O. H. (1994). CT Findings in neurologically normal adults with a single generalized seizure. *Journal of Epilepsy*, 7(1), 38-40.
- Dodick, D. W., Cascino, G. D., & Meyer, F. B. (1994). Vascular malformations and intractable epilepsy: Outcome after surgical treatment. *Mayo Clinic Proceedings*, 69(8), 741-745.
- Doelken, M. T., Mennecke, A., Huppertz, H. J., Rampp, S., Lukacs, E., Kasper, B. S., Hammen, T. (2012). Multimodality approach in cryptogenic epilepsy with focus on morphometric 3T MRI. *Journal of Neuroradiology*, 39(2), 87-96.
- Doelken, M. T., Richter, G., Stefan, H., Doerfler, A., Noemayr, A., Kuwert, T., Hammen, T. (2007). Multimodal coregistration in patients with temporal lobe epilepsy - Results of different imaging modalities in lateralization of the affected hemisphere in MR imaging positive and negative subgroups. *American Journal of Neuroradiology*, 28(3), 449-454.
- Dos Santos, S. L. M., Li, L. M., Ghizoni, E., & Cendes, F. (2004). Analysis of structural lesions in patients with partial epilepsy: Comparison between magnetic resonance imaging and computed tomography. *Journal of Epilepsy and Clinical Neurophysiology*, 10(3), 139-142.
- Duncan, J. S. (2002). Seizure-induced neuronal injury: Human data. *Neurology*, 59(9 SUPPL.), S15-S20.
- Duncan, R., Patterson, J., Hadley, D. M., Macpherson, P., Brodie, M. J., Bone, I., Wyper, D. J. (1990). CT, MR and SPECT imaging in temporal lobe epilepsy. *Journal of Neurology, Neurosurgery & Psychiatry*, 53(1), 11-15.
- Duplyakov, D., Golovina, G., Garkina, S., & Lyukshina, N. (2010). Is it possible to accurately differentiate neurocardiogenic syncope from epilepsy? *Cardiology Journal*, 17(4), 420-427.
- Ergün, E. L., Saygi, S., Yalnizoglu, D., Oguz, K. K., & Erbas, B. (2016). SPECT-PET in epilepsy and clinical approach in evaluation. *Seminars in Nuclear Medicine*, 46(4), 294-307.
- Flamini, J. R., Konkol, R. J., Wells, R. G., & Sty, J. R. (1990). I-123 lofetamine SPECT scan in children with neurological disorders. *Wisconsin Medical Journal*, 89(10), 584-587.
- Froment, J. C., Mauguere, F., Fischer, C., Revol, M., Bierme, T., & Convers, P. (1989). Magnetic resonance imaging in refractory focal epilepsy with normal CT scans. *Journal of Neuroradiology. Journal de Neuroradiologie*, 16(4), 285-291.
- Furune, S., Negoro, T., Maehara, M., Nomura, K., Miura, K., Takahashi, I., & Watanabe, K. (1989). Magnetic resonance imaging in complex partial seizures. *Japanese Journal of Psychiatry & Neurology*, 43(3), 361-367.
- Gaggero, R., Devescovi, R., Zacccone, A., & Ravera, G. (2001). Epilepsy associated with infantile hemiparesis: Predictors of long-term evolution. *Brain and Development*, 23(1), 12-17.
- Gaillard, W. D., Chiron, C., Cross, J. H., Harvey, A. S., Kuzniecky, R., Hertz-Pannier, L., Ilae, C. f. N. S. f. P. (2009). Guidelines for imaging infants and children with recent-onset epilepsy. *Epilepsia*, 50(9), 2147-2153.
- Gaillard, W. D., Cross, J. H., Duncan, J. S., Stefan, H., & Theodore, W. H. (2011). Epilepsy imaging study guideline criteria: Commentary on diagnostic testing study guidelines and practice parameters. *Epilepsia*, 52(9), 1750-1756.
- Goldstein, J. L., Leonhardt, D., Kmytyuk, N., Kim, F., Wang, D., & Wainwright, M. S. (2011). Abnormal neuroimaging is associated with early in-hospital seizures in pediatric abusive head trauma. *Neurocritical Care*, 15(1), 63-69.
- Gowda, V. K. N., Vasanna, S. H., Kumar, P., Shivananda, Lakshman, R. R., Govindraj, & Ramaswamy, P. (2013). Study of etiological profile of infantile and childhood focal seizures at a tertiary care centre in South India. *Journal of Pediatric Neurology*, 11(4), 235-240.
- Griffiths, P. D., & Baxter, P. (2001). Imaging epilepsy in childhood. *Imaging*, 13(4), 239-251.
- Grouiller, F., Delattre, B. M. A., Pittau, F., Heinzer, S., Lazeyras, F., Spinelli, L., Vulliemoz, S. (2015). All-in-one interictal presurgical imaging in patients with epilepsy: single-session EEG/PET/(f)MRI. *European Journal of Nuclear Medicine and Molecular Imaging*.
- Gupta, A. K., Sharma, R., & Sarma, D. (2000). Imaging in epilepsy. *Indian Journal of Pediatrics*, 67(1 Suppl), S40-60.
- Heinz, E. R., Heinz, T. R., Radtke, R., Darwin, R., Drayer, B. P., Fram, E., & Djang, W. T. (1989). Efficacy of MR vs

- CT in epilepsy. *AJR. American Journal of Roentgenology*, 152(2), 347-352.
- Heiskala, H., Launes, J., Pihko, H., Nikkinen, P., & Santavuori, P. (1993). Brain perfusion SPECT in children with frequent fits. *Brain and Development*, 15(3), 214-218.
- Ho, S. S., & Kuzniecky, R. I. (1997). Algorithm on the Clinical Evaluation of Epilepsy. *Journal of Neuroimaging*, 7(4), 236-241.
- Hsieh, D. T., Chang, T., Tsuchida, T. N., Vezina, L. G., Vanderver, A., Siedel, J., Gaillard, W. D. (2010). New-onset afebrile seizures in infants: Role of neuroimaging. *Neurology*, 74(2), 150-156.
- Hsu, Y. Y., Chang, C. N., Chu, N. S., & Hsu, J. C. (1997). Neuroimaging in intractable complex partial seizures. *Journal of the Formosan Medical Association*, 96(1), 51-54.
- Hugg, J. W., Laxer, K. D., Matson, G. B., Maudsley, A. A., & Weiner, M. W. (1993). Neuron loss localizes human temporal lobe epilepsy by in vivo proton magnetic resonance spectroscopic imaging. *Annals of Neurology*, 34(6), 788-794.
- Hwang, S. I., Kim, J. H., Park, S. W., Han, M. H., Yu, I. K., Lee, S. H., Chang, K. H. (2001). Comparative analysis of MR imaging, positron emission tomography, and ictal single-photon emission CT in patients with neocortical epilepsy. *Ajnr: American Journal of Neuroradiology*, 22(5), 937-946.
- Hwang, Y. S., & Kim, K. J. (1996). Tuberous sclerosis: Clinical, neuroradiologic and echocardiographic features. *Developmental Brain Dysfunction*, 9(5-6), 311-319.
- Jabbari, B., Gunderson, C. H., & Wippold, F. (1986). Magnetic resonance imaging in partial complex epilepsy. *Archives of Neurology*, 43(9), 869-872.
- Jackson, A., Teo, L., & Seneviratne, U. (2016). Challenges in the first seizure clinic for adult patients with epilepsy. *Epileptic Disorders*, 18(3), 305-314.
- Jan, M., Neville, B. G., Cox, T. C., & Scott, R. C. (2002). Convulsive status epilepticus in children with intractable epilepsy is frequently focal in origin. *Canadian Journal of Neurological Sciences*, 29(1), 65-67.
- Jensen, F. E. (2011). Epilepsy as a spectrum disorder: Implications from novel clinical and basic neuroscience. *Epilepsia*, 52(SUPPL. 1), 1-6.
- Kalifa, G. L., Chiron, C., Sellier, N., Demange, P., Ponsot, G., Lalande, G., & Robain, O. (1987). Hemimegalencephaly: MR imaging in five children. *Radiology*, 165(1), 29-33.
- Kato, H., Matsuda, K., Baba, K., Shimosegawa, E., Isohashi, K., Imaizumi, M., & Hatazawa, J. (2012). MR imaging-based correction for partial volume effect improves detectability of intractable epileptogenic foci on iodine 123 iomazenil brain SPECT images: An extended study with a larger sample size. *American Journal of Neuroradiology*, 33(11), 2088-2094.
- Kato, H., Shimosegawa, E., Oku, N., Kitagawa, K., Kishima, H., Saitoh, Y., Hatazawa, J. (2008). MRI-based correction for partial-volume effect improves detectability of intractable epileptogenic foci on 123I-iomazenil brain SPECT images. *Journal of Nuclear Medicine*, 49(3), 383-389.
- Kawakabani, A., Rossetti, A. O., & Despland, P. A. (2004). Survey of management of first-ever seizures in a hospital based community. *Swiss Medical Weekly*, 134(39-40), 586-592.
- Kilpatrick, C. J., Tress, B. M., O'Donnell, C., Rossiter, S. C., & Hopper, J. L. (1991). Magnetic resonance imaging and late-onset epilepsy. *Epilepsia*, 32(3), 358-364.
- Koirala, K. (2011). Magnetic resonance neuroimaging in patient with complain of seizure. *Journal of Nepal Health Research Council*, 9(1), 56-60.
- Kramer, U., Nevo, Y., Reider-Groszasser, I., Sheuer, E., Meyer, J. J., Leitner, Y., Harel, S. (1998). Neuroimaging of children with partial seizures. *Seizure*, 7(2), 115-118.
- Krsek, P., Hajek, M., Dezortova, M., Jiru, F., Skoch, A., Marusic, P., Komarek, V. (2007). 1H MR spectroscopic imaging in patients with MRI-negative extratemporal epilepsy: Correlation with ictal onset zone and histopathology. *European Radiology*, 17(8), 2126-2135.
- Krumholz, A., Wiebe, S., Gronseth, G., Shinnar, S., Levisohn, P., Ting, T., French, J. (2007). Practice parameter: Evaluating an apparent unprovoked first seizure in adults (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology*, 69(21), 1996-2007.
- Kuzniecky, R. I. (1996). Neuroimaging in pediatric epilepsy. *Epilepsia*, 37 Suppl 1, S10-21.
- Kuzniecky, R. I., & Knowlton, R. C. (2002). Neuroimaging of epilepsy. *Seminars in Neurology*, 22(3), 279-288.
- Lansberg, M. G., O'Brien, M. W., Norbash, A. M., Moseley, M. E., Morrell, M., & Albers, G. W. (1999). MRI abnormalities associated with partial status epilepticus. *Neurology*, 52(5), 1021-1027.
- Laster, D. W., Penry, J. K., & Moody, D. M. (1985). Chronic seizure disorders: Contribution of MR imaging when CT is normal. *American Journal of Neuroradiology*, 6(2), 177-180.
- Laxer, K. D., & Garcia, P. A. (1993). Imaging criteria to identify the epileptic focus. Magnetic resonance imaging, magnetic resonance spectroscopy, positron emission tomography scanning, and single photon emission

- computed tomography. *Neurosurgery Clinics of North America*, 4(2), 199-209.
- Lee, D. S., Lee, S. K., Kim, Y. K., Lee, J. S., Cheon, G. J., Kang, K. W., Lee, M. C. (2002). Superiority of HMPAO ictal SPECT to ECD ictal SPECT in localizing the epileptogenic zone. *Epilepsia*, 43(3), 263-269.
- Lesser, R. P., Modic, M. T., & Weinstein, M. A. (1986). Magnetic resonance imaging (1.5 tesla) in patients with intractable focal seizures. *Archives of Neurology*, 43(4), 367-371.
- Li, L. M., Fish, D. R., Sisodiya, S. M., Shorvon, S. D., Alsanjari, N., & Stevens, J. M. (1995). High resolution magnetic resonance imaging in adults with partial or secondary generalised epilepsy attending a tertiary referral unit. *Journal of Neurology Neurosurgery and Psychiatry*, 59(4), 384-387.
- Londoño, A., Castillo, M., Lee, Y. Z., & Smith, J. K. (2003). Apparent diffusion coefficient measurements in the hippocampi in patients with temporal lobe seizures. *American Journal of Neuroradiology*, 24(8), 1582-1586.
- Lyons, T. W., Johnson, K. B., Michelson, K. A., Nigrovic, L. E., Loddenkemper, T., Prabhu, S. P., & Kimia, A. A. (2016). Yield of emergent neuroimaging in children with new-onset seizure and status epilepticus. *Seizure*, 35, 4-10.
- Malik, M. A., Tarar, M. A., Hamid, H., Ur Rehman, M., Qureshi, A., Ossaid, M., Malik, S. (2010). Diagnostic importance of interictal electroencephalogram and neuroimaging of brain in new-onset idiopathic generalized epilepsy of childhood (IGEC). *Pakistan Paediatric Journal*, 34(1), 15-22.
- Marliani, A. F., Tampieri, D., Melancon, D., Ethier, R., Berkovic, S. F., & Andermann, F. (1991). Magnetic resonance imaging of hypothalamic hamartomas causing gelastic epilepsy. *Canadian Association of Radiologists Journal*, 42(5), 335-339.
- Masuoka, L. K., Anderson, A. W., Gore, J. C., McCarthy, G., Spencer, D. D., & Novotny, E. J. (1999). Functional magnetic resonance imaging identifies abnormal visual cortical function in patients with occipital lobe epilepsy. *Epilepsia*, 40(9), 1248-1253.
- Maytal, J., Krauss, J. M., Novak, G., Nagelberg, J., & Patel, M. (2000). The role of brain computed tomography in evaluating children with new onset of seizures in the emergency department. *Epilepsia*, 41(8), 950-954.
- Mengistu, G., Ewunetu, B. D., Johnston, J. C., & Metaferia, G. Z. (2014). Neuroimaging of Ethiopian patients with epilepsy: a retrospective review. *Ethiopian Medical Journal*, 52(2), 57-66.
- Mennink, S., van Nieuwenhuizen, O., Jennekens-Schinkel, A., van der Schouw, Y. T., van der Meij, W., & van Huffelen, A. C. (2003). Early prediction of seizure remission in children with occipital lobe epilepsy. *European Journal of Paediatric Neurology*, 7(4), 161-165.
- Menzel, C., Hufnagel, A., Grunwald, F., Pavics, L., Reichmann, K., Elger, C. E., & Biersack, H. J. (1995). The relevance of interictal rCBF brain SPECT in temporal lobe epilepsy: Diagnostical value and effects of spatial resolution. *Annals of Nuclear Medicine*, 9(4), 215-223.
- Menzel, C., Steidele, S., Grünwald, F., Hufnagel, A., Pavics, L., Elger, C. E., & Biersack, H. J. (1996). Evaluation of technetium-99m-ECD in childhood epilepsy. *Journal of Nuclear Medicine*, 37(7), 1106-1112.
- Merriam, M. A., Bronen, R. A., Spencer, D. D., & McCarthy, G. (1993). MR findings after depth electrode implantation for medically refractory epilepsy. *American Journal of Neuroradiology*, 14(6), 1343-1346.
- Michelucci, R., Poza, J. J., Sofia, V., De Feo, M. R., Binelli, S., Bisulli, F., Nobile, C. (2003). Autosomal Dominant Lateral Temporal Epilepsy: Clinical Spectrum, New Epitempin Mutations, and Genetic Heterogeneity in Seven European Families. *Epilepsia*, 44(10), 1289-1297.
- Mitsuyoshi, I., Tamaka, K., Okuno, T., Mutoh, K., Iwasaki, Y., Konishi, J., & Mikawa, H. (1993). Regional cerebral blood flow in diagnosis of childhood onset partial epilepsy. *Brain and Development*, 15(2), 97-102.
- Molla Mohammadi, M., Tonekaboni, S. H., Khatami, A., Azargashb, E., Tavasoli, A., Javadzadeh, M., & Zamani, G. (2013). Neuroimaging findings in first unprovoked seizures: A multicentric study in Tehran. *Iranian Journal of Child Neurology*, 7(4), 24-31.
- Montes, J. L., Rosenblatt, B., Farmer, J. P., O'Gorman, A. M., Andermann, F., Watters, G. V., & Meagher-Villemure, K. (1995). Lesionectomy of MRI detected lesions in children with epilepsy. *Pediatric Neurosurgery*, 22(4), 167-173.
- Murthy, J. M. K. (2013). Epilepsy: Diagnostic evaluation. *Journal of Association of Physicians of India*, 61(SPEC ISSUE 5), 10-14.
- Nair, P. P., Kalita, J., & Misra, U. K. (2009). Role of cranial imaging in epileptic status. *European Journal of Radiology*, 70(3), 475-480.
- Nam, H., Lee, S. K., Chung, C. K., Hong, K. S., Chang, K. H., & Lee, D. S. (2001). Incidence and clinical profile of extra-medial-temporal epilepsy with hippocampal atrophy. *Journal of Korean medical science*, 16(1), 95-102.
- Namer, I. J., Waydelich, R., Arnsbach, J. P., Hirsch, E., Marescaux, C., & Grucker, D. (1998). Contribution of T2 relaxation time mapping in the evaluation of cryptogenic temporal lobe epilepsy. *NeuroImage*, 7(4 I), 304-

313.

- Ng, K. K., Tang, K. W., Cheung, Y. L., & Li, P. C. (2000). Brain MRI of hippocampal volumetry in patients with refractory temporal lobe epilepsy. *Chinese medical journal*, 113(3), 254-256.
- Ormson, M. J., Kispert, D. B., & Sharbrough, F. W. (1986). Cryptic structural lesions in refractory partial epilepsy: MR imaging and CT studies. *Radiology*, 160(1), 215-219.
- Otsubo, H., Hoffman, H. J., Humphreys, R. P., Hendrick, E. B., Drake, J. M., Hwang, P. A., Chuang, S. H. (1992). Detection and management of gangliogliomas in children. *Surgical Neurology*, 38(5), 371-378.
- Owolabi, L. F., & Ogunniyi, A. (2015). Etiology and electroclinical pattern of late onset epilepsy in ibadan, southwestern nigeria. *African Journal of Neurological Sciences*, 34(2).
- Paliwal, P., Wakerley, B. R., Yeo, L. L. L., Ali, K. M., Ibrahim, I., Wilder-Smith, E., Rathakrishnan, R. (2015). Early electroencephalography in patients with Emergency Room diagnoses of suspected new-onset seizures: Diagnostic yield and impact on clinical decision-making. *Seizure*, 31, 22-26.
- Palmeri, B., Talamanca, S., Ragonese, P., Aridon, P., Daniele, O., Terruso, V., & D'Amelio, M. (2011). MRI abnormalities following repeated and incoming seizures. *Seizure*, 20(1), 90-92.
- Palmini, A., Andermann, F., Olivier, A., Tampieri, D., Robitaille, Y., Melanson, D., & Ethier, R. (1991). Neuronal migration disorders: a contribution of modern neuroimaging to the etiologic diagnosis of epilepsy. *Canadian Journal of Neurological Sciences*, 18(4 Suppl), 580-587.
- Peretti, P., Raybaud, C., Dravet, C., Mancini, J., & Pinsard, N. (1989). Magnetic resonance imaging in partial epilepsy of childhood. Seventy-nine cases. *Journal of Neuroradiology. Journal de Neuroradiologie*, 16(4), 308-316.
- Perissinotti, A., Setoain, X., Aparicio, J., Rubí, S., Fuster, B. M., Donaire, A., Pons, F. (2014). Clinical role of subtraction ictal SPECT coregistered to MR imaging and 18F-FDG PET in pediatric epilepsy. *Journal of Nuclear Medicine*, 55(7), 1099-1105.
- Phuttharak, W., Sawanyawisuth, K., Kawiungsanon, A., & Tiamkao, S. (2011). The appropriate neuroimaging study in persons with epilepsy. *Neurological Sciences*, 32(5), 969-971.
- Pohlmann-Eden, B., & Newton, M. (2008). First seizure: EEG and neuroimaging following an epileptic seizure. *Epilepsia*, 49 Suppl 1, 19-25.
- Prado, J. A. L. (2007). 3-Tesla MRI and Temporal Lobe Epilepsy. *Seminars in Ultrasound, CT and MRI*, 28(6), 451-461.
- Radue, E. W., & Scollo-Lavizzari, G. (1994). Computed tomography and magnetic resonance imaging in epileptic seizures. *European Neurology*, 34 Suppl 1, 55-57.
- Rajshekhar, V., & Chandy, M. J. (1996). Comparative study of CT and MRI in patients with seizures and a solitary cerebral cysticercus granuloma. *Neuroradiology*, 38(6), 542-546.
- Ramli, N., Rahmat, K., Lim, K. S., & Tan, C. T. (2015). Neuroimaging in refractory epilepsy. Current practice and evolving trends. *European Journal of Radiology*, 84(9), 1791-1800.
- Rasool, A., Choh, S. A., Wani, N. A., Mushtaq Ahmad, S., & Iqbal, Q. (2012). Role of electroencephalogram and neuroimaging in first onset afebrile and complex febrile seizures in children from Kashmir. *Journal of Pediatric Neurosciences*, 7(1), 9-15.
- Renou, P., Tourdias, T., Fleury, O., Debruxelles, S., Rouanet, F., & Sibon, I. (2012). Atraumatic nonaneurysmal sulcal subarachnoid hemorrhages: A diagnostic workup based on a case series. *Cerebrovascular Diseases*, 34(2), 147-152.
- Requena, I., Arias, M., Lopez-Ibor, L., Pereiro, I., Barba, A., Alonso, A., & Monton, E. (1991). Cavernomas of the central nervous system: clinical and neuroimaging manifestations in 47 patients. *Journal of Neurology, Neurosurgery & Psychiatry*, 54(7), 590-594.
- Resta, M., Palma, M., Dicuonzo, F., Spagnolo, P., Specchio, L. M., Laneve, A., La Selva, L. (1994). Imaging studies in partial epilepsy in children and adolescents. *Epilepsia*, 35(6), 1187-1193.
- Roffman, J. L., & Stern E, T. A. (2006). A complex presentation of complex partial seizures. *Primary Care Companion to the Journal of Clinical Psychiatry*, 8(2), 98-100.
- Saad, M., Zakarya, S., & Salama, H. (2011). Epilepsy in the elderly: Dilemma of diagnosis. *Egyptian Journal of Neurology, Psychiatry and Neurosurgery*, 48(1), 25-30.
- Salmenpera, T. M., & Duncan, J. S. (2005). Imaging in epilepsy. *Journal of Neurology, Neurosurgery & Psychiatry*, 76 Suppl 3, iii2-iii10.
- Sanmaneechai, O., Danchaivijitr, N., & Likasitwattanakul, S. (2015). Predictors of abnormal neuroimaging of the brain in children with epilepsy aged 1 month to 2 years. *Journal of Child Neurology*, 30(11), 1532-1536.
- Saygi, S., Katz, A., Marks, D. A., & Spencer, S. S. (1992). Frontal lobe partial seizures and psychogenic seizures: Comparison of clinical and ictal characteristics. *Neurology*, 42(7), 1274-1277.
- Scanlon, C., Mueller, S. G., Tosun, D., Cheong, I., Garcia, P., Barakos, J., Laxer, K. D. (2011). Impact of

- methodologic choice for automatic detection of different aspects of brain atrophy by using temporal lobe epilepsy as a model. *American Journal of Neuroradiology*, 32(9), 1669-1676.
- Schorner, W., Meencke, H. J., & Felix, R. (1987). Temporal-lobe epilepsy: comparison of CT and MR imaging. *AJR. American Journal of Roentgenology*, 149(6), 1231-1239.
- Shahar, E., Desatnik, H., Brand, N., Straussberg, R., & Hwang, P. A. (1996). Epileptic blindness in children: a localizing sign of various epileptic disorders. *Clinical Neurology & Neurosurgery*, 98(3), 237-241.
- Sharma, A. (2013). Seizures and epilepsy in children. *Indian Journal of Pediatrics*, 80(11), 925-935.
- Sharma, S., Riviello, J. J., Harper, M. B., & Baskin, M. N. (2003). The role of emergent neuroimaging in children with new-onset afebrile seizures. *Pediatrics*, 111(1), 1-5.
- Singh, G., Sachdev, M. S., Tirath, A., Gupta, A. K., & Avasthi, G. (2000). Focal cortical-subcortical calcifications (FCSCs) and epilepsy in the Indian subcontinent. *Epilepsia*, 41(6), 718-726.
- Singh, R. K., Stephens, S., Berl, M. M., Chang, T., Brown, K., Vezina, L. G., & Gaillard, W. D. (2010). Prospective study of new-onset seizures presenting as status epilepticus in childhood. *Neurology*, 74(8), 636-642.
- Spanaki, M. V., Spencer, S. S., Corsi, M., MacMullan, J., Seibyl, J., & Zubal, I. G. (1999). Sensitivity and specificity of quantitative difference SPECT analysis in seizure localization. *Journal of Nuclear Medicine*, 40(5), 730-736.
- Spencer, S. S., McCarthy, G., & Spencer, D. D. (1993). Diagnosis of medial temporal lobe seizure onset: Relative specificity and sensitivity of quantitative MRI. *Neurology*, 43(10), 2117-2124.
- Sperling, M. R., Wilson, G., Engel, J., Jr., Babb, T. L., Phelps, M., & Bradley, W. (1986). Magnetic resonance imaging in intractable partial epilepsy: correlative studies. *Annals of Neurology*, 20(1), 57-62.
- Stanescu, L., Ishak, G. E., Khanna, P. C., Biyyam, D. R., Shaw, D. W., & Parisi, M. T. (2013). FDG PET of the brain in pediatric patients: Imaging spectrum with MR imaging correlation. In (Vol. 33, pp. 1279-1303).
- Stefan, H., Pawlik, G., Bocher-Schwarz, H. G., Biersack, H. J., Burr, W., Penin, H., & Heiss, W. D. (1987). Functional and morphological abnormalities in temporal lobe epilepsy: a comparison of interictal and ictal EEG, CT, MRI, SPECT and PET. *Journal of Neurology*, 234(6), 377-384.
- Steffenburg, U., Hedstrom, A., Lindroth, A., Wiklund, L. M., Hagberg, G., & Kyllerman, M. (1998). Intractable epilepsy in a population-based series of mentally retarded children. *Epilepsia*, 39(7), 767-775.
- Swartz, B. E., Halgren, E., Delgado-Escueta, A. V., Feldstein, P., Maldonado, H., & Walsh, G. O. (1990). Multidisciplinary analysis of patients with extratemporal complex partial seizures. II. Predictive value of semiology. *Epilepsy Research*, 5(2), 146-154.
- Swartz, B. E., Tomiyasu, U., Delgado-Escueta, A. V., Mandelkern, M., & Khonsari, A. (1992). Neuroimaging in temporal lobe epilepsy: test sensitivity and relationships to pathology and postoperative outcome. *Epilepsia*, 33(4), 624-634.
- Tavassoli, A., & Noormohamadi, S. (2011). Factors related to abnormal neuroimaging in children with first unprovoked seizure. *Iranian Journal of Child Neurology*, 5(1), 15-20.
- Teixeira, R. A., Li, L. M., Santos, S. L. M., Amorim, B. J., Etchebehere, E. C. S. C., Zanardi, V. A., Cendes, F. (2004). Lateralization of epileptiform discharges in patients with epilepsy and precocious destructive brain insults. *Arquivos de Neuro-Psiquiatria*, 62(1), 1-8.
- Thakker, A. S., & Shetye, K. A. (2016). Neurodiagnostic evaluation of a child with first complex febrile seizure. *Journal of Krishna Institute of Medical Sciences University*, 5(2), 87-94.
- Theodore, W. H., Dorwart, R., & Holmes, M. (1986). Neuroimaging in refractory partial seizures: Comparison of PET, CT, and MRI. *Neurology*, 36(6), 750-759.
- Theodore, W. H., Katz, D., Kufta, C., Sato, S., Patronas, N., Smothers, P., & Bromfield, E. (1990). Pathology of temporal lobe foci: Correlation with CT, MRI, and PET. *Neurology*, 40(5), 797-803.
- Ulloa, C. M., & Gilliam, F. G. (2011). Epilepsy: Improving care for patients with epilepsy. *Nature Reviews Neurology*, 7(4), 195-196.
- Vanhainen, S. L., Puranen, J., Autti, T., Raininko, R., Liewendahl, K., Nikkinen, P., Häkkinen, A. M. (2004). Neuroradiological Findings (MRS, MRI, SPECT) in Infantile Neuronal Ceroid-Lipofuscinosis (Infantile CLN1) at Different Stages of the Disease. *Neuropediatrics*, 35(1), 27-35.
- Weinand, M. E., El-Saadany, W. F., Labiner, D. M., Talwar, D., & Ahern, G. (1995). Subdural strip electrode monitoring and surgical decision making in refractory epilepsy: Validity and prognostic value of noninvasive localizing data. *Journal of Epilepsy*, 8(2), 131-138.
- Welcker, S., Grams, A., Gizewski, E., Neubauer, B., & Hahn, A. (2013). Hemiconvulsion-hemiplegia-epilepsy syndrome with initially normal magnetic resonance imaging. *Klinische Padiatrie*, 225(2), 86-87.
- Verma, S. R., Sardana, V., Gupta, P. K., Verma, S. C., Munshi, A., & Suryavanshi, A. (2013). Evaluation of non febrile seizure disorder on MRI with correlation with seizure type and EEG records in a tertiary care teaching hospital. *Internet Journal of Third World Medicine*, 11(1).

- Widjaja, E., Li, B., & Santiago Medina, L. (2013). Diagnostic evaluation in patients with intractable epilepsy and normal findings on MRI: A decision analysis and cost-effectiveness study. *American Journal of Neuroradiology*, 34(5), 1004-1009.
- Wiesmann, U. C. (2003). Clinical application of neuroimaging in epilepsy. *Journal of Neurology, Neurosurgery & Psychiatry*, 74(4), 466-470.
- Yoong, M., Madari, R., Martinos, M., Clark, C., Chong, K., Neville, B., Scott, R. (2012). The role of magnetic resonance imaging in the follow-up of children with convulsive status epilepticus. *Developmental Medicine & Child Neurology*, 54(4), 328-333.
- Yucel, O., Aka, S., Yazicioglu, L., & Ceran, O. (2004). Role of early EEG and neuroimaging in determination of prognosis in children with complex febrile seizure. *Pediatrics International*, 46(4), 463-467.
- Zhou, Y. Z. (2005). Dynamic electroencephalogram monitoring in evaluation of paroxysmal cerebral disorder in children. *Chinese Journal of Clinical Rehabilitation*, 9(29), 184-186.
- Zhu, Z., & Chien, C. (2001). A preliminary study on comparison and fusion of metabolic images of PET with anatomic images of CT and MRI. *Chinese Medical Sciences Journal*, 16(2), 67-70.

Läkemedelsbehandling

Insättning (9)

- Arroyo, S., Dodson, W. E., Privitera, M. D., Glauser, T. A., Naritoku, D. K., Dlugos, D. J., Twyman, R. E. (2005). Randomized dose-controlled study of topiramate as first-line therapy in epilepsy. *Acta Neurol Scand*, 112(4), 214-222.
- Camfield, P., Camfield, C., Smith, S., Dooley, J., & Smith, E. (2002). Long-term outcome is unchanged by antiepileptic drug treatment after a first seizure: a 15-year follow-up from a randomized trial in childhood. *Epilepsia*, 43(6), 662-663.
- Fattore, C., Boniver, C., Capovilla, G., Cerminara, C., Citterio, A., Coppola, G., Perucca, E. (2011). A multicenter, randomized, placebo-controlled trial of levetiracetam in children and adolescents with newly diagnosed absence epilepsy. *Epilepsia*, 52(4), 802-809.
- Gilad, R., Lampl, Y., Gabbay, U., Eshel, Y., & Sarova-Pinhas, I. (1996). Early treatment of a single generalized tonic-clonic seizure to prevent recurrence. *Arch Neurol*, 53(11), 1149-1152.
- Glauser, T., Ben-Menachem, E., Bourgeois, B., Cnaan, A., Chadwick, D., Guerreiro, C., Tomson, T. (2006). ILAE treatment guidelines: evidence-based analysis of antiepileptic drug efficacy and effectiveness as initial monotherapy for epileptic seizures and syndromes. *Epilepsia*, 47(7), 1094-1120.
- Leone, MA, Giussani, G, Nolan, SJ, Marson, AG, Beghi, E. (2016). Immediate antiepileptic drug treatment, versus placebo, deferred, or no treatment for first unprovoked seizure. Cochrane Database of Systematic Reviews, Issue 5. Art. No.: CD007144.
- Musicco, M., Beghi, E., Solari, A., & Viani, F. (1997). Treatment of first tonic-clonic seizure does not improve the prognosis of epilepsy. First Seizure Trial Group (FIRST Group). *Neurology*, 49(4), 991-998.
- Najafi, M. R., Mehrabi, A., & Najafi, F. (2008). Seizure recurrence after a first unprovoked seizure: With and without treatment. *Journal of Research in Medical Sciences*, 13(4), 161-165.
- Young, K. D., Okada, P. J., Sokolove, P. E., Palchak, M. J., Panacek, E. A., Baren, J. M., Lewis, R. J. (2004). A randomized, double-blinded, placebo-controlled trial of phenytoin for the prevention of early posttraumatic seizures in children with moderate to severe blunt head injury. *Ann Emerg Med*, 43(4), 435-446.

Utsättning (15)

- Braathen, G., & Melander, H. (1997). Early discontinuation of treatment in children with uncomplicated epilepsy: a prospective study with a model for prediction of outcome. *Epilepsia*, 38(5), 561-569.
- Callaghan, N., Garrett, A., & Goggin, T. (1988). Withdrawal of anticonvulsant drugs in patients free of seizures for two years. A prospective study. *N Engl J Med*, 318(15), 942-946.
- Chiron, C., Dulac, O., & Gram, L. (1996). Vigabatrin withdrawal randomized study in children. *Epilepsy Res*, 25(3), 209-215.
- Hessen, E., Lossius, M. I., & Gjerstad, L. (2011). Improvement in verbal memory after withdrawal of carbamazepine and valproate in patients with well-controlled epilepsy: a randomized, double-blind study. *Acta Neurol Scand*, 123(6), 385-389.
- Hessen, E., Lossius, M. I., Reinvang, I., & Gjerstad, L. (2006). Influence of major antiepileptic drugs on attention,

- reaction time, and speed of information processing: results from a randomized, double-blind, placebo-controlled withdrawal study of seizure-free epilepsy patients receiving monotherapy. *Epilepsia*, 47(12), 2038-2045.
- Hessen, E., Lossius, M. I., Reinvang, I., & Gjerstad, L. (2007). Influence of major antiepileptic drugs on neuropsychological function: results from a randomized, double-blind, placebo-controlled withdrawal study of seizure-free epilepsy patients on monotherapy. *J Int Neuropsychol Soc*, 13(3), 393-400.
- Hessen, E., Lossius, M. I., Reinvang, I., & Gjerstad, L. (2007). Slight improvement in mood and irritability after antiepileptic drug withdrawal: a controlled study in patients on monotherapy. *Epilepsy Behav*, 10(3), 449-455.
- Hessen, E., Lossius, M. I., Reinvang, I., & Gjerstad, L. (2008). Improvement in speeded cognitive processing after anti-epileptic drug withdrawal - A controlled study in mono-therapy patients. *Progress in Neurotherapeutics and Neuropsychopharmacology*, 3(1), 199-209.
- Lossius MI, Hessen E, Mowinckel P, Stavem K, Eriksson J, Gulbrandsen P, et al. Consequences of antiepileptic drug withdrawal: a randomized, double-blind study (Akershus Study). *Epilepsia* 2008;49(3):455-63.
- Lossius, M. I., Nakken, K. O., Mowinckel, P., Taubøll, E., & Gjerstad, L. (2015). Favorable change of lipid profile after carbamazepine withdrawal. *Acta Neurol Scand*, 0.
- Lossius, M. I., Taubøll, E., Mowinckel, P., & Gjerstad, L. (2009). Reversible effects of antiepileptic drugs on thyroid hormones in men and women with epilepsy: a prospective randomized double-blind withdrawal study. *Epilepsy Behav*, 16(1), 64-68.
- MRC, Medical Research Council Antiepileptic Drug Withdrawal Study Group. Randomised study of antiepileptic drug withdrawal in patients in remission. *Lancet* 1991;337 (8751):1175-80.
- Strozzi I, Nolan SJ, Sperling MR, Wingerchuk DM, Sirven J. Early versus late antiepileptic drug withdrawal for people with epilepsy in remission. Cochrane Database of Systematic Reviews 2015, Issue 2. Art. No.: CD001902.
- Verrotti A, D'Egidio C, Agostinelli S, Parisi P, Spalice A, Chiarelli F, Coppola G. Antiepileptic drug withdrawal in childhood epilepsy: What are the risk factors associated with seizure relapse? *Journal of the European Paediatric Neurology Society*, 16, 599-604.
- Verrotti A, Morresi S, Cutarella R, Morgese G, Chiarelli F. Predictive value of EEG monitoring during drug withdrawal in children with cryptogenic partial epilepsy. *Neurophysiol Clin* 2000; 30:240-5.

Koncentrationsbestämning (27)

- Beydoun, A., Sackellares, J. C., & Shu, V. (1997). Safety and efficacy of divalproex sodium monotherapy in partial epilepsy: a double-blind, concentration-response design clinical trial. *Depakote Monotherapy for Partial Seizures Study Group. Neurology*, 48(1), 182-188.
- Christensen, J., Andreasen, F., Poulsen, J. H., & Dam, M. (2003). Randomized, concentration-controlled trial of topiramate in refractory focal epilepsy. *Neurology*, 61(9), 1210-1218.
- Dahlin, M., Knutsson, E., Amark, P., & Nergardh, A. (2000). Reduction of epileptiform activity in response to low-dose in children with epilepsy: a randomized double-blind study. *Epilepsia*, 41(3), 308-315.
- Faught, E., Wilder, B. J., Ramsay, R. E., Reife, R. A., Kramer, L. D., Pledger, G. W., & Karim, R. M. (1996). Topiramate placebo-controlled dose-ranging trial in refractory partial epilepsy using 200-, 400-, and 600-mg daily dosages. *Topiramate YD Study Group. Neurology*, 46(6), 1684-1690.
- Garnett, W. R., Levy, B., McLean, A. M., Zhang, Y., Couch, R. A., Rudnic, E. M., Belendiuk, G. W. (1998). Pharmacokinetic evaluation of twice-daily extended-release carbamazepine (CBZ) and four-times-daily immediate-release CBZ in patients with epilepsy. *Epilepsia*, 39(3), 274-279.
- Gilliam, F. G., Veloso, F., Bomhof, M. A., Gazda, S. K., Biton, V., Ter Bruggen, J. P., Wu, S. C. (2003). A dose-comparison trial of topiramate as monotherapy in recently diagnosed partial epilepsy. *Neurology*, 60(2), 196-202.
- McKee, P. J., Blacklaw, J., Butler, E., Gillham, R. A., & Brodie, M. J. (1991). Monotherapy with conventional and controlled-release carbamazepine: a double-blind, double-dummy comparison in epileptic patients. *Br J Clin Pharmacol*, 32(1), 99-104.
- Pledger, G. W., Sackellares, J. C., Treiman, D. M., Pellock, J. M., Wright, F. S., Mikati, M., al, e. (1994). Flunarizine for treatment of partial seizures: results of a concentration-controlled trial. *Neurology*, 44(10), 1830-1836.
- Pledger, G. W., Sackellares, J. C., Treiman, D. M., Pellock, J. M., Wright, F. S., Mikati, M., al, e. (1994). Flunarizine for treatment of partial seizures: results of a concentration-controlled trial. *Neurology*, 44(10), 1830-1836.
- Privitera, M. D., Welty, T. E., Gidal, B. E., Diaz, F. J., Krebill, R., Szaflarski, J. P., Berg, M. (2016). Generic-to-generic lamotrigine switches in people with epilepsy: the randomised controlled EQUIGEN trial. *Lancet Neurol*, 15(4), 365-372.

- Reed, R. C., Dutta, S., Cavanaugh, J. H., Locke, C., & Granneman, G. R. (2006). Every-12-hour administration of extended-release divalproex in patients with epilepsy: impact on plasma valproic acid concentrations. *Epilepsy Behav*, 8(2), 391-396.
- Reunanen, M., Heinonen, E., Anttila, M., Jarvensivu, P., Lehto, H., & Hokkanen, E. (1990). Multiple-dose pharmacokinetic study with a slow-release carbamazepine preparation. *Epilepsy Res*, 6(2), 126-133.
- Rey, E., Bulteau, C., Motte, J., Tran, A., Sturm, Y., D'Souza, J., Dulac, O. (2004). Oxcarbazepine pharmacokinetics and tolerability in children with inadequately controlled epilepsy. *J Clin Pharmacol*, 44(11), 1290-1300.
- Rimmer, E. M., & Richens, A. (1984). Double-blind study of gamma-vinyl GABA in patients with refractory epilepsy. *Lancet*, 1(8370), 189-190.
- Yegnanarayan, R., Mahesh, S. D., & Sangle, S. (2006). Chronotherapeutic dose schedule of phenytoin and carbamazepine in epileptic patients. *Chronobiol Int*, 23(5), 1035-1046.

Tilläggsbehandling – fokala epilepsi (105)

Studier av läkemedel som inte godkänts i Sverige på aktuell indikation har exkluderats.

- Andrews, J., Chadwick, D., Bates, D., Cartlidge, N., Boddie, G., Boile, P., Patterson, V. (1990). Gabapentin in partial epilepsy. *Lancet*, 335(8698), 1114-1117.
- Androsova, G., Krause, R., Borghei, M., Wassenaar, M., Auce, P., Avbersek, A., Zimprich, F. (2017). Comparative effectiveness of antiepileptic drugs in patients with mesial temporal lobe epilepsy with hippocampal sclerosis. *Epilepsia*, 58(10), 1734-1741.
- Asgarshirazi, M., Shariat, M., & Sheikh, M. (2017). Comparison of efficacy of folic acid and silymarin in the management of antiepileptic drug induced liver injury: a randomized clinical trial. *Hepatobiliary and Pancreatic Diseases International*, 16(3), 296-302.
- Baulac, M., Byrnes, W., Williams, P., Borghs, S., Webster, E., De Backer, M., & Dedeken, P. (2017). Lacosamide and sodium channel-blocking antiepileptic drug cross-titration against levetiracetam background therapy. *Acta Neurologica Scandinavica*, 135(4), 434-441.
- Baulac, M., Coulbaut, S., Doty, P., McShea, C., De Backer, M., Bartolomei, F., & Vlaicu, M. (2017). Adjunctive lacosamide for focal epilepsy: an open-label trial evaluating the impact of flexible titration and dosing on safety and seizure outcomes. *Epileptic Disorders*, 19(2), 186-194.
- Ben-Menachem, E., & Falter, U. (2000). Efficacy and tolerability of levetiracetam 3000 mg/d in patients with refractory partial seizures: a multicenter, double-blind, responder-selected study evaluating monotherapy. European Levetiracetam Study Group. *Epilepsia*, 41(10), 1276-1283.
- Beran, R. G., Berkovic, S. F., Buchanan, N., Danta, G., Mackenzie, R., Schapel, G., Vajda, F. (1996). A double-blind, placebo-controlled crossover study of vigabatrin 2 g/day and 3 g/day in uncontrolled partial seizures. *Seizure*, 5(4), 259-265.
- Berdichevsky, Y., Saponjian, Y., Park, K.-I., Roach, B., Pouliot, W., Lu, K., Staley, K. (2016). Staged anticonvulsant screening for chronic epilepsy. *Annals of clinical and translational neurology*, 3(12), 908-923.
- Berg, M., Welty, T. E., Gidal, B. E., Diaz, F. J., Krebill, R., Szaflarski, J. P., Privitera, M. D. (2017). Bioequivalence Between Generic and Branded Lamotrigine in People With Epilepsy: The EQUIGEN Randomized Clinical Trial. *JAMA Neurol*, 74(8), 919-926.
- Bernardina, B. D., Fontana, E., Vigevano, F., Fusco, L., Torelli, D., Galeone, D., Iudice, A. (1995). Efficacy and tolerability of vigabatrin in children with refractory partial seizures: A single-blind dose-increasing study. *Epilepsia*, 36(7), 687-691.
- Binnie, C. D., Debets, R. M. C., Engelsman, M., Meijer, J. W. A., Meinardi, H., Overweg, J., Yuen, W. C. (1989). Double-blind crossover trial of lamotrigine (Lamictal) as add-on therapy in intractable epilepsy. *Epilepsy Research*, 4(3), 222-229.
- Biton, V., Krauss, G., Vasquez-Santana, B., Bibbiani, F., Mann, A., Perdomo, C., & Narurkar, M. (2011). A randomized, double-blind, placebo-controlled, parallel-group study of rufinamide as adjunctive therapy for refractory partial-onset seizures. *Epilepsia*, 52(2), 234-242.
- Bonnett, L. J., Powell, G. A., Tudur Smith, C., & Marson, A. (2017). Risk of a seizure recurrence after a breakthrough seizure and the implications for driving: Further analysis of the standard versus new antiepileptic drugs (SANAD) randomised controlled trial. *BMJ Open*, 7(7).
- Brandt, C., Borghs, S., Elmoufti, S., Mueller, K., Townsend, R., & de la Loge, C. (2017). Health-related quality of life in double-blind Phase III studies of brivaracetam as adjunctive therapy of focal seizures: A pooled, post-hoc analysis. *Epilepsy and Behavior*, 69, 80-85.
- Brodie, M. J., Duncan, R., Vespignani, H., Solyom, A., Bitenskyy, V., & Lucas, C. (2005). Dose-dependent safety and efficacy of zonisamide: a randomized, double-blind, placebo-controlled study in patients with

- refractory partial seizures *Epilepsia*, 46(1), 31-41.
- Brodie, M. J., Lerche, H., Gil-Nagel, A., Elger, C., Hall, S., Shin, P., Mansbach, H. (2010). Efficacy and safety of adjunctive ezogabine (retigabine) in refractory partial epilepsy *Neurology*, 75(20), 1817-1824.
- Brodie, M. J., Rosenfeld, W. E., Vazquez, B., Sachdeo, R., Perdomo, C., Mann, A., & Arroyo, S. (2009). Rufinamide for the adjunctive treatment of partial seizures in adults and adolescents: a randomized placebo-controlled trial *Epilepsia*, 50(8), 1899-1909.
- Cao, J., Li, D., Gao, S.-S., & Song, Y.-J. (2016). Evaluation of the efficacy of adjunctive lamotrigine in the treatment of epilepsy and depression with Meta-analysis. *Chinese journal of contemporary neurology and neurosurgery*, 16(1), 23-29.
- Chaves, J., Breia, P., Pimentel, J., Pelejão, R., Carvalho, M., Mateus, P., Gala, A. (2017). Eslicarbazepine acetate as adjunctive therapy in clinical practice: ESLADOBA study. *Acta Neurologica Scandinavica*, 136(5), 407-413.
- Chiron, C., Tonnelier, S., Rey, E., Brunet, M. L., Tran, A., d'Athis, P., Pons, G. (2006). Stiripentol in childhood partial epilepsy: randomized placebo-controlled trial with enrichment and withdrawal design *J Child Neurol*, 21(6), 496-502.
- Chung, S. S., Fakhoury, T. A., Hogan, R. E., Nagaraddi, V. N., Blatt, I., Lawson, B., Halvorsen, M. B. (2014). Once-daily USL255 as adjunctive treatment of partial-onset seizures: randomized phase III study *Epilepsia*, 55(7), 1077-1087.
- Devinsky, O., Cross, J. H., Laux, L., Marsh, E., Miller, I., Nababout, R., Wright, S. (2017). Trial of cannabidiol for drug-resistant seizures in the dravet syndrome. *New England Journal of Medicine*, 376(21), 2011-2020.
- Devinsky, O., Marsh, E., Friedman, D., Thiele, E., Laux, L., Sullivan, J., Cilio, M. R. (2016). Cannabidiol in patients with treatment-resistant epilepsy: an open-label interventional trial. *The Lancet Neurology*, 15(3), 270-278.
- Elger, C. E., Stefan, H., Mann, A., Narurkar, M., Sun, Y., & Perdomo, C. (2010). A 24-week multicenter, randomized, double-blind, parallel-group, dose-ranging study of rufinamide in adults and adolescents with inadequately controlled partial seizures *Epilepsy Res*, 88(2-3), 255-263.
- Elger, C. E., Hong, S. B., Brandt, C., Mancione, L., Han, J., & Strohmaier, C. (2017). BGG492 as an adjunctive treatment in patients with partial-onset seizures: A 12-week, randomized, double-blind, placebo-controlled, phase II dose-titration study with an open-label extension. *Epilepsia*, 58(7), 1217-1226.
- Foldvary-Schaefer, N., Neme-Mercante, S., Andrews, N., Bruton, M., Wang, L., Morrison, S., Grigg-Damberger, M. (2017). Wake up to sleep: The effects of lacosamide on daytime sleepiness in adults with epilepsy. *Epilepsy and Behavior*, 75, 176-182.
- French, J. A., Baroldi, P., Brittain, S. T., & Johnson, J. K. (2014). Efficacy and safety of extended-release oxcarbazepine (Oxtellar XR) as adjunctive therapy in patients with refractory partial-onset seizures: a randomized controlled trial *Acta Neurol Scand*, 129(3), 143-153.
- French, J. A., Kanner, A. M., Bautista, J., Abou-Khalil, B., Browne, T., Harden, C. L., Glauser, T. A. (2007). Appendix D: Efficacy and tolerability of the new antiepileptic drugs II: Treatment of refractory epilepsy: Report of the Therapeutics and Technology Assessment Subcommittee and Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society *CONTINUUM Lifelong Learning in Neurology*, 13(4 EPILEPSY), 212-224.
- Gillham, R. A., Blacklaw, J., McKee, P. J., & Brodie, M. J. (1993). Effect of vigabatrin on sedation and cognitive function in patients with refractory epilepsy *J Neurol Neurosurg Psychiatry*, 56(12), 1271-1275.
- Glauser, T. A., Holland, K., O'Brien, V. P., Keddache, M., Martin, L. J., Clark, P. O., Grabowski, G. (2017). Pharmacogenetics of antiepileptic drug efficacy in childhood absence epilepsy. *Ann Neurol*, 81(3), 444-453.
- Gofshteyn, J., Wilfong, A., Devinsky, O., Bluvstein, J., Charuta, J., Ciliberto, M., Marsh, E. (2017). Cannabidiol as a Potential Treatment for Febrile Infection-Related Epilepsy Syndrome (FIRES) in the Acute and Chronic Phases. *Journal of child neurology*, 32(1), 35-40.
- Goyal, M., Lal, V., Sandal, R., Modi, M., Sachdeva, J., & Kharbanda, P. (2016). A comparative blinded study of cognitive effects of carbamazepine, valproate, lamotrigine and levetiracetam in adult epileptic patients. *Annals of indian academy of neurology. Conference: 24th annual conference of the indian academy of neurology, IANCON 2016. India. Conference start: 20161110. Conference end: 20161113*, 19(6), S93.
- Gram, L., Klosterskov, P., & Dam, M. (1985). γ -vinyl GABA: A double-blind placebo-controlled trial in partial epilepsy *Annals of Neurology*, 17(3), 262-266.
- Gram, L., Lyon, B. B., & Dam, M. (1983). Gamma-vinyl-GABA: A single-blind trial in patients with epilepsy *Acta Neurologica Scandinavica*, 68(1), 34-39.
- Guerrini, R., Rosati, A., Bradshaw, K., & Giorgi, L. (2014). Adjunctive zonisamide therapy in the long-term treatment of children with partial epilepsy: results of an open-label extension study of a phase III, randomized, double-blind, placebo-controlled trial *Epilepsia*, 55(4), 568-578.

- Hanin, A., Tubiana, R., Bedouet, M., Gales, A., Zahr, N., Liou, A., & Dupont, S. (2017). Rates of serum level determinations of antiepileptic drugs in accord with guidelines: A clinical study at a tertiary center. *Revue Neurologique*.
- Italian Study Group on Vigabatrin. Single-blind, placebo-controlled multicenter trial of vigabatrin in the treatment of epilepsy. (1992). *Ital J Neurol Sci*, 13(9), 741-747.
- Jawad, S., Richens, A., Goodwin, G., & Yuen, W. C. (1989). Controlled trial of lamotrigine (Lamictala) for refractory partial seizures *Epilepsia*, 30(3), 356-363.
- Kalvainen, R., Brodie, M. J., Duncan, J., Chadwick, D., Edwards, D., & Lyby, K. (1998). A double-blind, placebo-controlled trial of tiagabine given three-times daily as add-on therapy for refractory partial seizures. Northern European Tiagabine Study Group *Epilepsy Res*, 30(1), 31-40.
- Kerr, M. P., Baker, G. A., & Brodie, M. J. (2005). A randomized, double-blind, placebo-controlled trial of topiramate in adults with epilepsy and intellectual disability: impact on seizures, severity, and quality of life *Epilepsy Behav*, 7(3), 472-480.
- Kessler, S. K., Shinnar, S., Cnaan, A., Drugos, D., Conry, J., Hirtz, D. G., Glauser, T. A. (2017). Pretreatment seizure semiology in childhood absence epilepsy. *Neurology*, 89(7), 673-679.
- Kim, J. H., Lee, S. K., Loesch, C., Namgoong, K., Lee, H. W., & Hong, S. B. (2017). Comparison of levetiracetam and oxcarbazepine monotherapy among Korean patients with newly diagnosed focal epilepsy: A long-term, randomized, open-label trial. *Epilepsia*, 58(4), e70-e74.
- Koeppen, D., Baruzzi, A., Capozza, M., Chauvel, P., Courjon, J., Favel, P., Wolf, P. (1987). Clobazam in therapy-resistant patients with partial epilepsy: A double-blind placebo-controlled crossover study *Epilepsia*, 28(5), 495-506.
- Kothare, S., Kluger, G., Sachdeo, R., Williams, B., Olhaye, O., Perdomo, C., & Bibbiani, F. (2017). Dosing considerations for rufinamide in patients with Lennox–Gastaut syndrome: Phase III trial results and real-world clinical data. *Seizure*, 47, 25-33.
- Krauss, G. L., Bar, M., Biton, V., Klapper, J. A., Rektor, I., Vaiciene-Magistris, N., Kumar, D. (2012). Tolerability and safety of perampanel: two randomized dose-escalation studies *Acta Neurol Scand*, 125(1), 8-15.
- Krauss, G. L., Serratosa, J. M., Villanueva, V., Endziniene, M., Hong, Z., French, J., Laurenza, A. (2012). Randomized phase III study 306: adjunctive perampanel for refractory partial-onset seizures *Neurology*, 78(18), 1408-1415.
- Kroep, S., Bouwmeester, W., Dimova, S., Zhang, Y., Borghs, S., & Charokopou, M. (2017). Relative efficacy and safety of lacosamide as monotherapy for adults with newly diagnosed focal seizures: a network meta-analysis of randomized clinical trials. *Value in health. Conference: ISPOR 22nd annual international meeting. United states*, 20(5), A341.
- Lee, B. I. (1998). Double-blind placebo-controlled randomized clinical trial of topiramate add-on therapy in medically intractable partial epilepsies *Journal of the Korean Neurological Association*, 16(6), 809-819.
- Lee, B. I., Yi, S., Hong, S. B., Kim, M. K., Lee, S. A., Lee, S. K., Leon, T. (2009). Pregabalin add-on therapy using a flexible, optimized dose schedule in refractory partial epilepsies: a double-blind, randomized, placebo-controlled, multicenter trial *Epilepsia*, 50(3), 464-474.
- Levisor, P. M., Mintz, M., Hunter, S. J., Yang, H., & Jones, J. (2009). Neurocognitive effects of adjunctive levetiracetam in children with partial-onset seizures: a randomized, double-blind, placebo-controlled, noninferiority trial *Epilepsia*, 50(11), 2377-2389.
- Liu, M. J., Su, X. J., Md, X. Y. S., Wu, G. F., Zhang, Y. Q., Gao, L., Zou, L. P. (2017). Clinical features of benign epilepsy of childhood with centrotemporal spikes in Chinese children. *Medicine*, 96(4), e5623.
- Loiseau, P., Hardenberg, J. P., Pestre, M., Guyot, M., Schechter, P. J., & Tell, G. P. (1986). Double-blind, placebo-controlled study of vigabatrin (gamma-vinyl GABA) in drug-resistant epilepsy *Epilepsia*, 27(2), 115-120.
- Loiseau, P., Yuen, A. W., Duche, B., Menager, T., & Arne-Bes, M. C. (1990). A randomised double-blind placebo-controlled crossover add-on trial of lamotrigine in patients with treatment-resistant partial seizures *Epilepsy Res*, 7(2), 136-145.
- Lowrie, M., Thomson, S., Bessant, C., Sparkes, A., Harvey, R. J., & Garosi, L. (2017). Levetiracetam in the management of feline audiogenic reflex seizures: a randomised, controlled, open-label study. *J Feline Med Surg*, 19(2), 200-206.
- Luna, D., Dulac, O., Pajot, N., & Beaumont, D. (1989). Vigabatrin in the treatment of childhood epilepsies: A single-blind placebo-controlled study *Epilepsia*, 30(4), 430-437.
- Mann, D., Liu, J., Chew, M. L., Bockbrader, H., Alvey, C. W., Zegarac, E., Pitman, V. W. (2014). Safety, tolerability, and pharmacokinetics of pregabalin in children with refractory partial seizures: a phase 1, randomized controlled study *Epilepsia*, 55(12), 1934-1943.
- McGinty, R., & Costello, D. (2017). Long-term lacosamide retention—Real-world experience at a tertiary epilepsy

- center in Ireland. *Epilepsy & behavior*, 68, 141-145.
- McKee, P. J., Blacklaw, J., Friel, E., Thompson, G. G., Gillham, R. A., & Brodie, M. J. (1993). Adjuvant vigabatrin in refractory epilepsy: a ceiling to effective dosage in individual patients? *Epilepsia*, 34(5), 937-943.
- McLean, M. J., Ramsay, R. E., Leppik, L. E., Rowan, A. J., Shellenberger, M. K., & Wallace, J. (1993). Gabapentin as add-on therapy in refractory partial epilepsy: A double-blind, placebo-controlled, parallel-group study *Neurology*, 43(11), 2292-2298.
- McShea, C., Isojarvi, J., Doty, P., Berkovic, S., Kiley, M., O'Brien, T., Smith, P. (2016). Long-term adjunctive lacosamide treatment in patients with partial-onset seizures. *Acta Neurologica Scandinavica*, 133(2), 136-144.
- Michelucci, R., Cavaciocchi, P., Riguzzi, P., Passarelli, D., Parmeggiani, L., Santangelo, M., Tassinari, C. A. (1992). Single-blind, placebo-controlled dose-modification study of vigabatrin in refractory epileptic patients *Journal of Epilepsy*, 5(4), 248-252.
- Miller, R., Frame, B., Corrigan, B., Burger, P., Bockbrader, H., Garofalo, E., & Lalonde, R. (2003). Exposure-response analysis of pregabalin add-on treatment of patients with refractory partial seizures *Clin Pharmacol Ther*, 73(6), 491-505.
- Misra, U. K., & Kalita, J. (2016). A comparison of four antiepileptic drugs in status epilepticus: experience from India. *Int J Neurosci*, 126(11), 1013-1019.
- Morrell, M. J., Leppik, I., French, J., Ferrendelli, J., Han, J., & Magnus, L. (2003). The KEEPER trial: levetiracetam adjunctive treatment of partial-onset seizures in an open-label community-based study *Epilepsy Res*, 54(2-3), 153-161.
- Mukuria, C., Young, T., Keetharuth, A., Borghs, S., & Brazier, J. (2017). Sensitivity and responsiveness of the EQ-5D-3L in patients with uncontrolled focal seizures: an analysis of Phase III trials of adjunctive brivaracetam. *Qual Life Res*, 26(3), 749-759.
- Nickels, K. (2017). Cannabidiol in patients with intractable epilepsy due to TSC: A possible medication but not a miracle. *Epilepsy Currents*, 17(2), 91-92.
- Novotny, E., Renfroe, B., Yardi, N., Nordli, D., Ness, S., Wang, S., Ford, L. (2010). Randomized trial of adjunctive topiramate therapy in infants with refractory partial seizures *Neurology*, 74(9), 714-720.
- Peigné, S., Chhun, S., Tod, M., Rey, E., Rodrigues, C., Chiron, C., Jullien, V. (2017). Population Pharmacokinetics of Stiripentol in Paediatric Patients with Dravet Syndrome Treated with Stiripentol, Valproate and Clobazam Combination Therapy. *Clinical Pharmacokinetics*, 1-10.
- Perez, J., Chiron, C., Musial, C., Rey, E., Blehaut, H., D'Athis, P., Dulac, O. (1999). Stiripentol: Efficacy and tolerability in children with epilepsy *Epilepsia*, 40(11), 1618-1626.
- Pina-Garza, J. E., Elterman, R. D., Ayala, R., Corral, M., Mikati, M. A., Pina-Garza, M. J., Messenheimer, J. A. (2008). Long-term tolerability and efficacy of lamotrigine in infants 1 to 24 months old *J Child Neurol*, 23(8), 853-861.
- Pina-Garza, J. E., Levisohn, P., Gucuyener, K., Mikati, M. A., Warnock, C. R., Conklin, H. S., & Messenheimer, J. (2008). Adjunctive lamotrigine for partial seizures in patients aged 1 to 24 months *Neurology*, 70(22 Pt 2), 2099-2108.
- Pina-Garza, J. E., Nordli, D. R., Jr., Rating, D., Yang, H., Schiemann-Delgado, J., & Duncan, B. (2009). Adjunctive levetiracetam in infants and young children with refractory partial-onset seizures *Epilepsia*, 50(5), 1141-1149.
- Porter, R. J., Partiot, A., Sachdeo, R., Nohria, V., & Alves, W. M. (2007). Randomized, multicenter, dose-ranging trial of retigabine for partial-onset seizures *Neurology*, 68(15), 1197-1204.
- Premoli, I., Biondi, A., Carlesso, S., Rivolta, D., & Richardson, M. P. (2017). Lamotrigine and levetiracetam exert a similar modulation of TMS-evoked EEG potentials. *Epilepsia*, 58(1), 42-50.
- Qu, J., Lu, S. H., Lu, Z. L., Xu, P., Xiang, D. X., & Qu, Q. (2017). Pharmacogenetic and case-control study on potassium channel related gene variants and genetic generalized epilepsy. *Medicine (United States)*, 96(26).
- Rentmeester, T., Janssen, A., Hulsman, J., Scholtes, F., van der Kleij, B., Overweg, J., de Beukelaar, F. (1991). A double-blind, placebo-controlled evaluation of the efficacy and safety of lorcetazole as add-on therapy in patients with uncontrolled partial seizures *Epilepsy Res*, 9(1), 59-64.
- Reynolds, E. H., Ring, H. A., Farr, I. N., Heller, A. J., & Elwes, R. D. (1991). Open, double-blind and long-term study of vigabatrin in chronic epilepsy *Epilepsia*, 32(4), 530-538.
- Rosenfeld, W., Conry, J., Lagae, L., Rozentals, G., Yang, H., Fain, R., Laurenza, A. (2015). Efficacy and safety of perampanel in adolescent patients with drug-resistant partial seizures in three double-blind, placebo-controlled, phase III randomized clinical studies and a combined extension study *Eur J Paediatr Neurol*, 19(4), 435-445.

- Rudzinski, L., Vlez-Ruiz, N., Gedelman, E., Mauricio, E., Shih, J., & Karakis, I. (2016). New antiepileptic drugs: focus on ezogabine, clobazam, and perampanel. *Journal of investigative medicine*, 64(6), 1087-1101.
- Sachdeo, R. C., Leroy, R. F., Krauss, G. L., Drake, M. E., Jr., Green, P. M., Leppik, I. E., Sommerville, K. W. (1997). Tiagabine therapy for complex partial seizures. A dose-frequency study. The Tiagabine Study Group *Arch Neurol*, 54(5), 595-601.
- Sander, J. W., Patsalos, P. N., Oxley, J. R., Hamilton, M. J., & Yuen, W. C. (1990). A randomised double-blind placebo-controlled add-on trial of lamotrigine in patients with severe epilepsy *Epilepsy Res*, 6(3), 221-226.
- Schachter, S. C., Leppik, I. E., Matsuo, F., Messenheimer, J. A., Faught, E., Moore, E. L., & Risner, M. E. (1995). Lamotrigine: A six-month, placebo-controlled, safety and tolerance study *Journal of Epilepsy*, 8(3), 201-209.
- Schmidt, D., Beyenburg, S., D'Souza, J., & Stavem, K. (2013). Clinical features associated with placebo response in refractory focal epilepsy *Epilepsy Behav*, 27(2), 393-398.
- Schmidt, D. (2016). Drug treatment strategies for epilepsy revisited: starting early or late? One drug or several drugs? *Epileptic Disorders*, 18(4), 356-366.
- Schoemaker, R., Wade, J., & Stockis, A. (2017). Brivaracetam population pharmacokinetics in children with epilepsy aged 1 month to 16 years. *European journal of clinical pharmacology*, 1-7.
- Schoemaker, R., Wade, J. R., & Stockis, A. (2017). Extrapolation of a Brivaracetam Exposure-Response Model from Adults to Children with Focal Seizures. *Clinical Pharmacokinetics*, 1-12.
- Schoenberg, M. R., Rum, R. S., Osborn, K. E., & Werz, M. A. (2017). A randomized, double-blind, placebo-controlled crossover study of the effects of levetiracetam on cognition, mood, and balance in healthy older adults. *Epilepsia*, 58(9), 1566-1574.
- Sedighi, M., Asadi, F., Moradian, N., Vakiliyani, M., & Moradian, M. (2016). Efficacy and safety of levetiracetam in the management of seizures in neonates. *Neurosciences (Riyadh)*, 21(3), 232-235.
- Shih, J., Whitlock, J., Chimato, N., Vargas, E., Karceski, S., & Frank, R. (2016). Epilepsy treatment in adults and adolescents: expert opinion, 2016. *Epilepsy and behavior. (no pagination)*, 2016, Date of Publication: September 02.
- Shinnar, R. C., Shinnar, S., Cnaan, A., Clark, P., Dlugos, D., Hirtz, D. G., Glauser, T. A. (2017). Pretreatment behavior and subsequent medication effects in childhood absence epilepsy. *Neurology*, 89(16), 1698-1706.
- Sidhu, H. S., Srinivas, R., & Sadhotra, A. (2017). Evaluate the effects of long-term valproic acid treatment on metabolic profiles in newly diagnosed or untreated female epileptic patients: A prospective study. *Seizure*, 48, 15-21.
- Sperling, M. R., Greenspan, A., Cramer, J. A., Kwan, P., Kalvainen, R., Halford, J. J., Novak, G. (2010). Carisbamate as adjunctive treatment of partial onset seizures in adults in two randomized, placebo-controlled trials *Epilepsia*, 51(3), 333-343.
- Sperling, M. R., Klein, P., & Tsai, J. (2017). Randomized, double-blind, placebo-controlled phase 2 study of ganaxolone as add-on therapy in adults with uncontrolled partial-onset seizures. *Epilepsia*, 58(4), 558-564.
- Steinig, I., von Podewils, F., Möddel, G., Bauer, S., Klein, K. M., Paule, E., Strzelczyk, A. (2017). Postmarketing experience with brivaracetam in the treatment of epilepsies: A multicenter cohort study from Germany. *Epilepsia*, 58(7), 1208-1216.
- Striano, P., & Belcastro, V. (2017). Update on pharmacotherapy of myoclonic seizures. *Expert opinion on pharmacotherapy*, 18(2), 187-193.
- Theitler, J., Brik, A., Shaniv, D., Berkovitch, M., & Gandelman-Marton, R. (2017). Antiepileptic Drug Treatment in Community-Dwelling Older Patients with Epilepsy: A Retrospective Observational Study of Old- Versus New-Generation Antiepileptic Drugs. *Drugs and Aging*, 1-9.
- Uldall, P., Bulteau, C., Pedersen, S. A., Dulac, O., & Lyby, K. (2000). Tiagabine adjunctive therapy in children with refractory epilepsy: A single-blind dose escalating study *Epilepsy Research*, 42(2-3), 159-168.
- Uthman, B. M., Almas, M., Emir, B., Giordano, S., & Leon, T. (2011). Pregabalin or placebo used adjunctively with levetiracetam in refractory partial-onset epilepsy: A post hoc efficacy and safety analysis in combined clinical trials *Current Medical Research and Opinion*, 27(7), 1285-1293.
- Uthman, B. M., Rowan, A. J., Ahmann, P. A., Leppik, I. E., Schachter, S. C., Sommerville, K. W., & Shu, V. (1998). Tiagabine for complex partial seizures: a randomized, add-on, dose-response trial *Arch Neurol*, 55(1), 56-62.
- Wechsler, R. T., Yates, S. L., Messenheimer, J., Leroy, R., Beller, C., & Doty, P. (2017). Lacosamide for uncontrolled primary generalized tonic-clonic seizures: An open-label pilot study with 59-week extension. *Epilepsy Research*, 130, 13-20.
- Xu, X. X., Liu, X. R., Fan, C. Y., Lai, J. X., Shi, Y. W., Yang, W., Liao, W. P. (2017). Functional Investigation of a GRIN2A Variant Associated with Rolandic Epilepsy. *Neuroscience Bulletin*, 1-10.
- Yamamoto, T., Shimizu, M., Sato, K., Kurata, A., & Numachi, Y. (2017). Long-term efficacy and safety of

- lamotrigine: Results from the open-label extension of the lamotrigine monotherapy study in adult Japanese patients with epilepsy. *Epilepsy and Seizure*, 9(1), 11-17.
- Yamauchi, T., Kaneko, S., Yagi, K., & Sase, S. (2006). Treatment of partial seizures with gabapentin: double-blind, placebo-controlled, parallel-group study. *Psychiatry Clin Neurosci*, 60(4), 507-515.
- Yu, P., Zhou, D., Liao, W., Wang, X., Wang, Y., Wang, T., Hong, Z. (2017). An investigation of the characteristics of outpatients with epilepsy and antiepileptic drug utilization in a multicenter cross-sectional study in China. *Epilepsy & behavior*, 69, 126-132.
- Zhang, L., Zou, X., Li, S., & Li, H. (2016). Eslicarbazepine vs brivaracetam for adults with refractory partial seizures: a meta-analysis. *Epilepsia. Conference: 12th European Congress on Epileptology. Czech Republic. Conference start: 20160911. Conference end: 20160915*, 57, 181-182.

Tilläggsbehandling – epilepsi generaliserade tonisk-kloniska anfall (69)

Studier av läkemedel som inte godkänts i Sverige på aktuell indikation har exkluderats.

- Androsova, G., Krause, R., Borghei, M., Wassenaar, M., Auce, P., Avbersek, A., Zimprich, F. (2017). Comparative effectiveness of antiepileptic drugs in patients with mesial temporal lobe epilepsy with hippocampal sclerosis. *Epilepsia*, 58(10), 1734-1741.
- Asgarshirazi, M., Shariat, M., & Sheikh, M. (2017). Comparison of efficacy of folic acid and silymarin in the management of antiepileptic drug induced liver injury: a randomized clinical trial. *Hepatobiliary and Pancreatic Diseases International*, 16(3), 296-302.
- Baulac, M., Byrnes, W., Williams, P., Borghs, S., Webster, E., De Backer, M., & Dedeken, P. (2017). Lacosamide and sodium channel-blocking antiepileptic drug cross-titration against levetiracetam background therapy. *Acta Neurologica Scandinavica*, 135(4), 434-441.
- Baulac, M., Coulbaut, S., Doty, P., McShea, C., De Backer, M., Bartolomei, F., & Vlaicu, M. (2017). Adjunctive lacosamide for focal epilepsy: an open-label trial evaluating the impact of flexible titration and dosing on safety and seizure outcomes. *Epileptic Disorders*, 19(2), 186-194.
- Beran, R. G., Berkovic, S. F., Dunagan, F. M., Vajda, F. J., Danta, G., Black, A. B., & Mackenzie, R. (1998). Double-blind, placebo-controlled, crossover study of lamotrigine in treatment-resistant generalised epilepsy. *Epilepsia*, 39(12), 1329-1333.
- Berdichevsky, Y., Saponjian, Y., Park, K.-I., Roach, B., Pouliot, W., Lu, K., Staley, K. (2016). Staged anticonvulsant screening for chronic epilepsy. *Annals of Clinical and Translational Neurology*, 3(12), 908-923.
- Berg, M., Welty, T. E., Gidal, B. E., Diaz, F. J., Krebill, R., Szaflarski, J. P., Privitera, M. D. (2017). Bioequivalence Between Generic and Branded Lamotrigine in People With Epilepsy: The EQUIGEN Randomized Clinical Trial. *JAMA Neurol*, 74(8), 919-926.
- Bonnett, L. J., Powell, G. A., Tudur Smith, C., & Marson, A. (2017). Risk of a seizure recurrence after a breakthrough seizure and the implications for driving: Further analysis of the standard versus new antiepileptic drugs (SANAD) randomised controlled trial. *BMJ Open*, 7(7).
- Brandt, C., Borghs, S., Elmoufti, S., Mueller, K., Townsend, R., & de la Loge, C. (2017). Health-related quality of life in double-blind Phase III studies of brivaracetam as adjunctive therapy of focal seizures: A pooled, post-hoc analysis. *Epilepsy and Behavior*, 69, 80-85.
- Cao, J., Li, D., Gao, S.-S., & Song, Y.-J. (2016). Evaluation of the efficacy of adjunctive lamotrigine in the treatment of epilepsy and depression with Meta-analysis. *Chinese Journal of Contemporary Neurology and Neurosurgery*, 16(1), 23-29.
- Chadwick, D., Leiderman, D. B., Sauermann, W., Alexander, J., & Garofalo, E. (1996). Gabapentin in generalized seizures. *Epilepsy Res*, 25(3), 191-197.
- Chaves, J., Breia, P., Pimentel, J., Pelejão, R., Carvalho, M., Mateus, P., Gala, A. (2017). Eslicarbazepine acetate as adjunctive therapy in clinical practice: ESLADOBA study. *Acta Neurologica Scandinavica*, 136(5), 407-413.
- Devinsky, O., Cross, J. H., Laux, L., Marsh, E., Miller, I., Nabbout, R., Wright, S. (2017). Trial of cannabidiol for drug-resistant seizures in the dravet syndrome. *New England Journal of Medicine*, 376(21), 2011-2020.
- Devinsky, O., Marsh, E., Friedman, D., Thiele, E., Laux, L., Sullivan, J., Cilio, M. R. (2016). Cannabidiol in patients with treatment-resistant epilepsy: an open-label interventional trial. *The Lancet Neurology*, 15(3), 270-278.
- Elger, C. E., Hong, S. B., Brandt, C., Mancione, L., Han, J., & Strohmaier, C. (2017). BGG492 as an adjunctive treatment in patients with partial-onset seizures: A 12-week, randomized, double-blind, placebo-controlled, phase II dose-titration study with an open-label extension. *Epilepsia*, 58(7), 1217-1226.
- Eriksson, A. S., Nergardh, A., & Hoppu, K. (1998). The efficacy of lamotrigine in children and adolescents with refractory generalized epilepsy: a randomized, double-blind, crossover study. *Epilepsia*, 39(5), 495-501.

- Foldvary-Schaefer, N., Neme-Mercante, S., Andrews, N., Bruton, M., Wang, L., Morrison, S., Grigg-Damberger, M. (2017). Wake up to sleep: The effects of lacosamide on daytime sleepiness in adults with epilepsy. *Epilepsy and Behavior*, 75, 176-182.
- Glauser, T. A., Holland, K., O'Brien, V. P., Keddache, M., Martin, L. J., Clark, P. O., Grabowski, G. (2017). Pharmacogenetics of antiepileptic drug efficacy in childhood absence epilepsy. *Ann Neurol*, 81(3), 444-453.
- Gillham, R. A., Blacklaw, J., McKee, P. J., & Brodie, M. J. (1993). Effect of vigabatrin on sedation and cognitive function in patients with refractory epilepsy. *J Neurol Neurosurg Psychiatry*, 56(12), 1271-1275.
- Glauser, T., Kluger, G., Sachdeo, R., Krauss, G., Perdomo, C., & Arroyo, S. (2008). Rufinamide for generalized seizures associated with Lennox-Gastaut syndrome. *Neurology*, 70(21), 1950-1958.
- Gofshteyn, J., Wilfong, A., Devinsky, O., Bluvstein, J., Charuta, J., Ciliberto, M., Marsh, E. (2017). Cannabidiol as a Potential Treatment for Febrile Infection-Related Epilepsy Syndrome (FIREs) in the Acute and Chronic Phases. *Journal of child neurology*, 32(1), 35-40.
- Goyal, M., Lal, V., Sandal, R., Modi, M., Sachdeva, J., & Kharbanda, P. (2016). A comparative blinded study of cognitive effects of carbamazepine, valproate, lamotrigine and levetiracetam in adult epileptic patients. *Annals of indian academy of neurology. Conference: 24th annual conference of the indian academy of neurology, IANCON 2016. India. Conference start: 20161110. Conference end: 20161113*, 19(6), S93.
- Hanin, A., Tubiana, R., Bedouet, M., Gales, A., Zahr, N., Liou, A., & Dupont, S. (2017). Rates of serum level determinations of antiepileptic drugs in accord with guidelines: A clinical study at a tertiary center. *Revue Neurologique*.
- Kerr, M. P., Baker, G. A., & Brodie, M. J. (2005). A randomized, double-blind, placebo-controlled trial of topiramate in adults with epilepsy and intellectual disability: impact on seizures, severity, and quality of life. *Epilepsy Behav*, 7(3), 472-480.
- Kessler, S. K., Shinnar, S., Cnaan, A., Dlugos, D., Conry, J., Hirtz, D. G., Glauser, T. A. (2017). Pretreatment seizure semiology in childhood absence epilepsy. *Neurology*, 89(7), 673-679.
- Kim, J. H., Lee, S. K., Loesch, C., Namgoong, K., Lee, H. W., & Hong, S. B. (2017). Comparison of levetiracetam and oxcarbazepine monotherapy among Korean patients with newly diagnosed focal epilepsy: A long-term, randomized, open-label trial. *Epilepsia*, 58(4), e70-e74.
- Kothare, S., Kluger, G., Sachdeo, R., Williams, B., Olhaye, O., Perdomo, C., & Bibbiani, F. (2017). Dosing considerations for rufinamide in patients with Lennox–Gastaut syndrome: Phase III trial results and real-world clinical data. *Seizure*, 47, 25-33.
- Kroep, S., Bouwmeester, W., Dimova, S., Zhang, Y., Borghs, S., & Charokopou, M. (2017). Relative efficacy and safety of lacosamide as monotherapy for adults with newly diagnosed focal seizures: a network meta-analysis of randomized clinical trials. *Value in health. Conference: ISPOR 22nd annual international meeting. United states*, 20(5), A341.
- Kwan, P., Trinka, E., Van Paesschen, W., Rektor, I., Johnson, M. E., & Lu, S. (2014). Adjunctive brivaracetam for uncontrolled focal and generalized epilepsies: results of a phase III, double-blind, randomized, placebo-controlled, flexible-dose trial. *Epilepsia*, 55(1), 38-46.
- Liu, M. J., Su, X. J., Md, X. Y. S., Wu, G. F., Zhang, Y. Q., Gao, L., Zou, L. P. (2017). Clinical features of benign epilepsy of childhood with centrotemporal spikes in chinese children. *Medicine*, 96(4), e5623.
- Loiseau, P., Hardenberg, J. P., Pestre, M., Guyot, M., Schechter, P. J., & Tell, G. P. (1986). Double-blind, placebo-controlled study of vigabatrin (gamma-vinyl GABA) in drug-resistant epilepsy. *Epilepsia*, 27(2), 115-120.
- Lowrie, M., Thomson, S., Bessant, C., Sparkes, A., Harvey, R. J., & Garosi, L. (2017). Levetiracetam in the management of feline audiogenic reflex seizures: a randomised, controlled, open-label study. *J Feline Med Surg*, 19(2), 200-206.
- Luna, D., Dulac, O., Pajot, N., & Beaumont, D. (1989). Vigabatrin in the treatment of childhood epilepsies: A single-blind placebo-controlled study. *Epilepsia*, 30(4), 430-437.
- McGinty, R., & Costello, D. (2017). Long-term lacosamide retention-Real-world experience at a tertiary epilepsy center in Ireland. *Epilepsy & behavior*, 68, 141-145.
- McShea, C., Isojarvi, J., Doty, P., Berkovic, S., Kiley, M., O'Brien, T., Smith, P. (2016). Long-term adjunctive lacosamide treatment in patients with partial-onset seizures. *Acta Neurologica Scandinavica*, 133(2), 136-144.
- Michelucci, R., Cavaciocchi, P., Riguzzi, P., Passarelli, D., Parmeggiani, L., Santangelo, M., Tassinari, C. A. (1992). Single-blind, placebo-controlled dose-modification study of vigabatrin in refractory epileptic patients. *Journal of Epilepsy*, 5(4), 248-252.
- Misra, U. K., & Kalita, J. (2016). A comparison of four antiepileptic drugs in status epilepticus: experience from India. *Int J Neurosci*, 126(11), 1013-1019.
- Motte, J., Trevathan, E., Arvidsson, J. F. V., Barrera, M. N., Mullens, E. L., & Manasco, P. (1997). Lamotrigine for

- generalized seizures associated with the Lennox-Gastaut syndrome. *New England Journal of Medicine*, 337(25), 1807-1812.
- Mukuria, C., Young, T., Keetharuth, A., Borghs, S., & Brazier, J. (2017). Sensitivity and responsiveness of the EQ-5D-3L in patients with uncontrolled focal seizures: an analysis of Phase III trials of adjunctive brivaracetam. *Qual Life Res*, 26(3), 749-759.
- Nickels, K. (2017). Cannabidiol in patients with intractable epilepsy due to TSC: A possible medication but not a miracle. *Epilepsy Currents*, 17(2), 91-92.
- Noachtar, S., Andermann, E., Meyvisch, P., Andermann, F., Gough, W. B., & Schiemann-Delgado, J. (2008). Levetiracetam for the treatment of idiopathic generalized epilepsy with myoclonic seizures. *Neurology*, 70(8), 607-616.
- Pack, A. M. (2016). Perampanel: Another choice for patients with idiopathic generalized epilepsy who have tonic-clonic seizures. *Epilepsy Currents*, 16(1), 27-28.
- Palhagen, S., Canger, R., Henriksen, O., van Parys, J. A., Riviere, M. E., & Karolchik, M. A. (2001). Rufinamide: a double-blind, placebo-controlled proof of principle trial in patients with epilepsy. *Epilepsy Res*, 43(2), 115-124.
- Peigné, S., Chhun, S., Tod, M., Rey, E., Rodrigues, C., Chiron, C., Jullien, V. (2017). Population Pharmacokinetics of Stiripentol in Paediatric Patients with Dravet Syndrome Treated with Stiripentol, Valproate and Clobazam Combination Therapy. *Clinical Pharmacokinetics*, 1-10.
- Perez, J., Chiron, C., Musial, C., Rey, E., Blehaut, H., D'Athis, P., Dulac, O. (1999). Stiripentol: Efficacy and tolerability in children with epilepsy. *Epilepsia*, 40(11), 1618-1626.
- Premoli, I., Biondi, A., Carlesso, S., Rivolta, D., & Richardson, M. P. (2017). Lamotrigine and levetiracetam exert a similar modulation of TMS-evoked EEG potentials. *Epilepsia*, 58(1), 42-50.
- Qu, J., Lu, S. H., Lu, Z. L., Xu, P., Xiang, D. X., & Qu, Q. (2017). Pharmacogenetic and case-control study on potassium channel related gene variants and genetic generalized epilepsy. *Medicine (United States)*, 96(26).
- Reynolds, E. H., Ring, H. A., Farr, I. N., Heller, A. J., & Elwes, R. D. (1991). Open, double-blind and long-term study of vigabatrin in chronic epilepsy. *Epilepsia*, 32(4), 530-538.
- Rudzinski, L., Vlez-Ruiz, N., Gedelman, E., Mauricio, E., Shih, J., & Karakis, I. (2016). New antiepileptic drugs: focus on ezogabine, clobazam, and perampanel. *Journal of investigative medicine*, 64(6), 1087-1101.
- Sander, J. W., Patsalos, P. N., Oxley, J. R., Hamilton, M. J., & Yuen, W. C. (1990). A randomised double-blind placebo-controlled add-on trial of lamotrigine in patients with severe epilepsy. *Epilepsy Res*, 6(3), 221-226.
- Schmidt, D. (2016). Drug treatment strategies for epilepsy revisited: starting early or late? One drug or several drugs? *Epileptic Disorders*, 18(4), 356-366.
- Schoemaker, R., Wade, J., & Stockis, A. (2017). Brivaracetam population pharmacokinetics in children with epilepsy aged 1 month to 16 years. *European journal of clinical pharmacology*, 1-7.
- Schoemaker, R., Wade, J. R., & Stockis, A. (2017). Extrapolation of a Brivaracetam Exposure-Response Model from Adults to Children with Focal Seizures. *Clinical Pharmacokinetics*, 1-12.
- Schoenberg, M. R., Rum, R. S., Osborn, K. E., & Werz, M. A. (2017). A randomized, double-blind, placebo-controlled crossover study of the effects of levetiracetam on cognition, mood, and balance in healthy older adults. *Epilepsia*, 58(9), 1566-1574.
- Sedighi, M., Asadi, F., Moradian, N., Vakiliyamini, M., & Moradian, M. (2016). Efficacy and safety of levetiracetam in the management of seizures in neonates. *Neurosciences (Riyadh)*, 21(3), 232-235.
- Shih, J., Whitlock, J., Chimato, N., Vargas, E., Karceski, S., & Frank, R. (2016). Epilepsy treatment in adults and adolescents: expert opinion, 2016. *Epilepsy and behavior*. (no pagination), 2016, Date of Publication: September 02.
- Shinnar, R. C., Shinnar, S., Cnaan, A., Clark, P., Dlugos, D., Hirtz, D. G., Glauser, T. A. (2017). Pretreatment behavior and subsequent medication effects in childhood absence epilepsy. *Neurology*, 89(16), 1698-1706.
- Sidhu, H. S., Srinivas, R., & Sadhotra, A. (2017). Evaluate the effects of long-term valproic acid treatment on metabolic profiles in newly diagnosed or untreated female epileptic patients: A prospective study. *Seizure*, 48, 15-21.
- Sperling, M. R., Klein, P., & Tsai, J. (2017). Randomized, double-blind, placebo-controlled phase 2 study of ganaxolone as add-on therapy in adults with uncontrolled partial-onset seizures. *Epilepsia*, 58(4), 558-564.
- Steinig, I., von Podewils, F., Möddel, G., Bauer, S., Klein, K. M., Paule, E., Strzelczyk, A. (2017). Postmarketing experience with brivaracetam in the treatment of epilepsies: A multicenter cohort study from Germany. *Epilepsia*, 58(7), 1208-1216.
- Striano, P., & Belcastro, V. (2017). Update on pharmacotherapy of myoclonic seizures. Expert opinion on pharmacotherapy, 18(2), 187-193.

- Theitler, J., Brik, A., Shaniv, D., Berkovitch, M., & Gandelman-Marton, R. (2017). Antiepileptic Drug Treatment in Community-Dwelling Older Patients with Epilepsy: A Retrospective Observational Study of Old- Versus New-Generation Antiepileptic Drugs. *Drugs and Aging*, 1-9.
- Trevathan, E., Kerls, S. P., Hammer, A. E., Vuong, A., & Messenheimer, J. A. (2006). Lamotrigine adjunctive therapy among children and adolescents with primary generalized tonic-clonic seizures. *Pediatrics*, 118(2), e371-378.
- Uldall, P., Bulteau, C., Pedersen, S. A., Dulac, O., & Lyby, K. (2000). Tiagabine adjunctive therapy in children with refractory epilepsy: A single-blind dose escalating study. *Epilepsy Research*, 42(2-3), 159-168.
- Wechsler, R. T., Yates, S. L., Messenheimer, J., Leroy, R., Beller, C., & Doty, P. (2017). Lacosamide for uncontrolled primary generalized tonic-clonic seizures: An open-label pilot study with 59-week extension. *Epilepsy Research*, 130, 13-20.
- Xu, X. X., Liu, X. R., Fan, C. Y., Lai, J. X., Shi, Y. W., Yang, W., Liao, W. P. (2017). Functional Investigation of a GRIN2A Variant Associated with Rolandic Epilepsy. *Neuroscience Bulletin*, 1-10.
- Yamamoto, T., Shimizu, M., Sato, K., Kurata, A., & Numachi, Y. (2017). Long-term efficacy and safety of lamotrigine: Results from the open-label extension of the lamotrigine monotherapy study in adult Japanese patients with epilepsy. *Epilepsy and Seizure*, 9(1), 11-17.
- Yu, P., Zhou, D., Liao, W., Wang, X., Wang, Y., Wang, T., Hong, Z. (2017). An investigation of the characteristics of outpatients with epilepsy and antiepileptic drug utilization in a multicenter cross-sectional study in China. *Epilepsy & behavior*, 69, 126-132.
- Zhang, L., Zou, X., Li, S., & Li, H. (2016). Eslicarbazepine vs brivaracetam for adults with refractory partial seizures: a meta-Analysis. *Epilepsia. Conference: 12th european congress on epileptology. Czech republic. Conference start: 20160911. Conference end: 20160915*, 57, 181-182.

Anfallskuperande läkemedel (5)

- Appleton, R. E., Sweeney, A., Choonara, I., Robson, J., & Molyneux, E. (1993). Lorazepam vs diazepam in the acute treatment of epilepsy in children. *Seizure*, (2), 159.
- Appleton, R., Sweeney, A., Choonara, I., Robson, J., & Molyneux, E. (1995). Lorazepam versus diazepam in the acute treatment of epileptic seizures and status epilepticus. *Developmental medicine and child neurology*, 37(8), 682-688.
- Mahvelati, F., Tonekaboni, H., Javadzade, M., & Ghofrani, M. (2007). The efficacy of propofol and midazolam in treatment of refractory status epilepticus in children. *Iranian Journal of Medical Sciences*, 32(2), 74-79.
- Milligan, N. M., Dhillon, S., & Griffiths, A. (1984). A clinical trial of single dose rectal and oral administration of diazepam for the prevention of serial seizures in adult epileptic patients. *Journal of Neurology Neurosurgery and Psychiatry*, 47(3), 235-240.
- Malu CK, Kahamba DM, Walker TD, Mukampunga C, Musalu EM, Kokolomani J, et al. Efficacy of sublingual lorazepam versus intrarectal diazepam for prolonged convulsions in Sub-Saharan Africa. *J Child Neurol* 2014; 29: 89-902.
- Moretti, R., Julliand, S., Rinaldi, V. E., & Titomanlio, L. (2017). Buccal Midazolam Compared With Rectal Diazepam Reduces Seizure Duration in Children in the Outpatient Setting.

Anti-depressiva läkemedel (79)

- Alper K, Schwartz KA, Kolts RL, Khan A. Seizure incidence in psychopharmacological clinical trials: an analysis of Food and Drug Administration (FDA) summary basis of approval reports. *Biol Psychiatry* 2007; 62:345-54.
- Aman, M. G., Lam, K. S., & Collier-Crespin, A. (2003). Prevalence and patterns of use of psychoactive medicines among individuals with autism in the Autism Society of Ohio. *J Autism Dev Disord*, 33(5), 527-534.
- Andersen, B. B., Mikkelsen, M., Vesterager, A., Dam, M., Kristensen, H. B., Pedersen, B., Mengel, H. (1991). No influence of the antidepressant paroxetine on carbamazepine, valproate and phenytoin. *Epilepsy Res*, 10(2-3), 201-204.
- Bhatti, M., Dorris, P., & Mehndiratta, P. (2017). Impact of psychotropic drugs on seizure threshold. *Neurology*, 88(16).
- Bloechliger, M., Ceschi, A., Rüegg, S., Kupferschmidt, H., Kraehenbuehl, S., Jick, S. S., Bodmer, M. (2016). Risk of Seizures Associated with Antidepressant Use in Patients with Depressive Disorder: Follow-up Study with a Nested Case-Control Analysis Using the Clinical Practice Research Datalink. *Drug Safety*, 39(4), 307-321.
- Blumer, D. (1997). Antidepressant and double antidepressant treatment for the affective disorder of epilepsy. *Journal of Clinical Psychiatry*, 58(1), 3-11.
- Blumer, D., Montouris, G., Davies, K., Wyler, A., Phillips, B., & Hermann, B. (2002). Suicide in epilepsy:

- Psychopathology, pathogenesis, and prevention. *Epilepsy and Behavior*, 3(3 I), 232-241.
- Blöchliger, M., Ceschi, A., Rüegg, S., Kupferschmidt, H., Krähenbühl, S., Jick, S., Bodmer, M. (2015). Risk of seizures associated with antidepressant use in patients with depressive disorder-a nested case-control analysis. *European Psychiatry*, 30, 186.
- Brandt, A., Godfred, R., Parikh, M., Haltiner, A., Fortier, N., Caylor, L., Doherty, M. (2014). Do antidepressants make seizures more or less likely in focal epilepsy patients undergoing elective eeg monitoring? *Neurology*, 82(10).
- Bryan, G. E., Marsh, E. E., Jabbari, B., & Gunderson, C. H. (1983). Incidence in seizures in patients receiving tricyclic and tetracyclic antidepressants. *Annals of Neurology*, 14(1), P159.
- Buchanan, N. (2001). Medications which may lower seizure threshold. *Australian Prescriber*, 24(1), 8.
- Burley, D., Jukes, A., & Steen, J. (1978). Maprotiline hydrochloride and grand-mal seizures. *Br Med J*, 2(6146), 1230.
- Caracci, G., & Decina, P. (1991). Fluoxetine and Prolonged Seizure. *Convuls Ther*, 7(2), 145-147.
- Cole, J. O., Hartmann, E., & Brigham, P. (1980). L-Tryptophan: Clinical studies. *McLean Hospital Journal*, 5(1), 37-71.
- Davison, K. (1965). EEG activation after intravenous amitriptyline. *Electroencephalography and clinical neurophysiology*, 19(3), 298-300.
- Dessain, E. C., Schatzberg, A. F., Woods, B. T., & Cole, J. O. (1986). Maprotiline treatment in depression. A perspective on seizures. *Arch Gen Psychiatry*, 43(1), 86-90.
- Dobesberger, J., Ristic, A. J., Walser, G., Höfler, J., Unterberger, I., & Trinka, E. (2014). Selective serotonin reuptake inhibitors prolong seizures - Preliminary results from an observational study. *Clinical Neurology and Neurosurgery*, 120, 89-92.
- Downs, J., Giust, J., & Dunn, D. W. (2017). Considerations for ADHD in the child with epilepsy and the child with migraine. *Expert Review of Neurotherapeutics*, 17(9), 861-869.
- Duncan, D., & Taylor, D. (1995). Which is the safest antidepressant to use in epilepsy? *Psychiatric Bulletin*, 19(6), 355-357.
- Dunn, D. W., Besag, F., Caplan, R., Aldenkamp, A., Gobbi, G., & Sillanpää, M. (2016). Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): Anxiety, depression and childhood epilepsy. *Epileptic Disorders*, 18, S24-S30.
- Elmassry, H. A., Nagy, N., Samy, N., & Eid, M. (2015). Cross section comparative study for comorbid migraine and major depression in epileptic patients: Demographic, clinical, electroencephalography and therapeutic implications. *Neuropsychiatric Electrophysiology*, 1(1).
- Fogel, B. S., & Haltzman, S. (1987). Desipramine and phenytoin: a potential drug interaction of therapeutic relevance. *J Clin Psychiatry*, 48(9), 387-388.
- Fromm, G., & Wessel, H. (1977). Imipramine in absence and minor motor seizures: double-blind study and long-term follow-up. *Neurology*, 27(4), 370.
- Fromm, G., Wessel, H., Glass, J., Alvin, J., & Horn, G. (1978). Imipramine in absence and myoclonic-astatic seizures. *Neurology*, 28(9 Pt 1), 953-957.
- Fromm, G. H., & Wessel, H. B. (1977). Imipramine in absence and minor motor seizures: Double-blind study and long-term follow-up. *Neurology*, 27(4), 370.
- Fromm, G. H., Wessel, H. B., & Glass, J. D. (1978). Imipramine in absence and myoclonic-astatic seizures. *Neurology*, 28(9 I), 953-957.
- Gabb, M. G., & Barry, J. J. (2005). The link between mood disorders and epilepsy: Why is it important to diagnose and treat? *Advanced Studies in Medicine*, 5(6 C), S572-S578.
- Gross, A., Devinsky, O., Westbrook, L. E., Wharton, A. H., & Alper, K. (2000). Psychotropic medication use in patients with epilepsy: Effect on seizure frequency. *Journal of Neuropsychiatry and Clinical Neurosciences*, 12(4), 458-464.
- Hall, M. J., & Russell, R. I. (1978). Maprotiline hydrochloride and grand-mal seizures. *Br Med J*, 2(6142), 961.
- Harmant, J., van Rijckevorsel-Harmant, K., de Barsy, T., & Hendrickx, B. (1990). Fluvoxamine: an antidepressant with low (or no) epileptogenic effect. *Lancet*, 336(8711), 386.
- Hill, T., Coupland, C., Morriss, R., Arthur, A., Moore, M., & Hippisley-Cox, J. (2015). Antidepressant use and risk of epilepsy and seizures in people aged 20 to 64 years: cohort study using a primary care database. *BMC Psychiatry*, 15, 315.
- Hitchings, A. W. (2016). Drugs that lower the seizure threshold. *Adverse Drug Reaction Bulletin*, 298(1), 1151-1154.
- Kanner, A. M., Kozak, A. M., & Frey, M. (2000). The Use of Sertraline in Patients with Epilepsy: Is It Safe? *Epilepsy Behav*, 1(2), 100-105.

- Karnaze, D. S., & Weiss, G. (1992). Tricyclic antidepressants may cause seizure exacerbation. *American Family Physician*, 45(3), 1384.
- Karouni, M., Arulthas, S., Larsson, P. G., Rytter, E., Johannessen, S. I., & Johannessen Landmark, C. (2010). Psychiatric comorbidity in patients with epilepsy: A population-based study. *European Journal of Clinical Pharmacology*, 66(11), 1151-1160.
- Karouni, M., Henning, O., Larsson, P. G., Johannessen, S. I., & Johannessen Landmark, C. (2013). Pharmacological treatment of psychiatric comorbidity in patients with refractory epilepsy. *Epilepsy and Behavior*, 29(1), 77-81.
- Kerr, M. P., Mensah, S., Besag, F., De Toffol, B., Ettinger, A., Kanemoto, K., Wilson, S. J. (2011). International consensus clinical practice statements for the treatment of neuropsychiatric conditions associated with epilepsy. *Epilepsia*, 52(11), 2133-2138.
- Kuhn, K. U., Quednow, B. B., Thiel, M., Falkai, P., Maier, W., & Elger, C. E. (2003). Antidepressive treatment in patients with temporal lobe epilepsy and major depression: a prospective study with three different antidepressants. *Epilepsy Behav*, 4(6), 674-679.
- Kumlien, E., & Lundberg, P. O. (2010). Seizure risk associated with neuroactive drugs: data from the WHO adverse drug reactions database. *Seizure*, 19(2), 69-73.
- Kühn, K., Quednow, B., Thiel, M., Falkai, P., Maier, W., & Elger, C. (2003). Antidepressive treatment in patients with temporal lobe epilepsy and major depression: a prospective study with three different antidepressants. *Epilepsy Behav*, 4(6), 674-679.
- Li, W., & Ma, D.-R. (2005). A randomized controlled trial to evaluate the efficacy of paroxetine and doxepin in treating epileptic patients with depression. *Chinese journal of clinical rehabilitation*, 9(12), 20-21.
- Lundberg, P., Strandell, J., & Kumlien, E. (2009). Seizure risk associated with neuroactive drugs: A report from WHO collaboration centre for international drug monitoring. *Epilepsia*, 50, 199-200.
- Maguire, M. J., Weston, J., Singh, J., & Marson, A. G. (2014). Antidepressants for people with epilepsy and depression. *Cochrane Database Syst Rev*(12), Cd010682.
- Montilla, V. G., Pérez, A. V., Paredes, G. A., & Vera, M. (2014). Relative effectiveness of escitalopram versus Sertraline in patients with epilepsy and depression receiving anticonvulsant oxcarbazepine monotherapy. Instituto Autónomo Hospital Universitario de Los Andes, Mérida, Venezuela. December 2012-July 2013. *Epileptic Disorders*, 16, 44-45.
- Novak, P. H., Ekins-Daukes, S., Simpson, C. R., Milne, R. M., Helms, P., & McLay, J. S. (2005). Acute drug prescribing to children on chronic antiepilepsy therapy and the potential for adverse drug interactions in primary care. *British Journal of Clinical Pharmacology*, 59(6), 712-717.
- Ojemann, L. M., Friel, P. N., Trejo, W. J., & Dudley, D. L. (1983). Effect of doxepin on seizure frequency in depressed epileptic patients. *Neurology*, 33(5), 646-648.
- Ojemann, L. M., Friel, P. N., Trejo, W. R., & Dudley, D. L. (1982). Doxepin treatment for the depressed epileptic patient. *Neurology*, 32(4 II), A 226.
- Ojong, M., & Allen, S. N. (2012). Treatment of depression in patients with epilepsy. In (Vol. 37, pp. 29-32).
- Okazaki, M., Adachi, N., Ito, M., Watanabe, M., Watanabe, Y., Kato, M., & Onuma, T. (2011). One-year seizure prognosis in epilepsy patients treated with antidepressants. *Epilepsy and Behavior*, 22(2), 331-335.
- Orjuela-Rojas, J., Martínez-Juárez, I., Ruiz-Chow, A., & Crail-Melendez, D. (2015). Treatment of depression in patients with temporal lobe epilepsy: A pilot study of cognitive behavioral therapy vs. selective serotonin reuptake inhibitors. *Epilepsy Behav*, 51, 176-181.
- Orjuela-Rojas, J. M., Martínez-Juárez, I. E., Ruiz-Chow, A., & Crail-Melendez, D. (2015). Treatment of depression in patients with temporal lobe epilepsy: A pilot study of cognitive behavioral therapy vs. selective serotonin reuptake inhibitors. *Epilepsy and Behavior*, 51, 176-181.
- Osman, A., Seri, S., & Cavanna, A. E. (2016). Clinical characteristics of patients with epilepsy in a specialist neuropsychiatry service. *Epilepsy and Behavior*, 58, 44-47.
- Pantovic, M., Milovanovic, D., Pantovic, V., Ravanic, B., Ravanic, J., Jovic, S., Djukic Dejanovic, S. (2009). Amelioration of major depressive disorder in epilepsy with sertraline D. Ravanic1. *Journal of the Neurological Sciences*, 285, S117-S118.
- Payandemehr, B., Ghasemi, M., & Dehpour, A. R. (2015). Citalopram as a good candidate for treatment of depression in patients with epilepsy. *Epilepsy Behav*, 44, 96-97.
- Pollock, B. G., Perel, J. M., & Shostak, M. (1986). Understanding the response lag to tricyclics. I. Application of pulse-loading regimens with intravenous clomipramine. *Psychopharmacology Bulletin*, 22(1), 214-219.
- Popkin, M. K., Callies, A. L., & Mackenzie, T. B. (1985). The outcome of antidepressant use in the medically ill. *Arch Gen Psychiatry*, 42(12), 1160-1163.
- Popli, A. P., Kando, J. C., Pillay, S. S., Tohen, M., & Cole, J. O. (1995). Occurrence of seizures related to

- psychotropic medication among psychiatric inpatients. *Psychiatric Services*, 46(5), 486-488.
- Radmanovic, B., Djukic Dejanovic, S., Ravanic, D., Mihajlovic, G., Pantovic, M., Janjic, V., Ilic, M. (2012). Treatment of major depressive disorder in epilepsy. *International Journal of Neuropsychopharmacology*, 15, 154.
- Ravanic, D., Jovanovic, M., Janjic, V., Mihajlovic, G., Djukic Dejanovic, S., Petrovic, D., Radmanovic, B. (2010). Amelioration of major depressive disorder in epilepsy with escitalopram. *International Journal of Neuropsychopharmacology*, 13, 160.
- Robertson, M., & Trimble, M. (1985). The treatment of depression in patients with epilepsy. A double-blind trial. *Journal of Affective Disorders*, 9(2), 127-136.
- Robertson, M. M., & Trimble, M. R. (1985). The treatment of depression in patients with epilepsy. A double-blind trial. *Journal of Affective Disorders*, 9(2), 127-136.
- Romanowicz, M., & Bostwick, J. M. (2012). Epileptic and depressed. *Current Psychiatry*, 11(5), 45-49.
- Sparshatt, A., Whiskey, E., & Taylor, D. (2008). Valproate as prophylaxis for clozapine-induced seizures: Survey of practice. *Psychiatric Bulletin*, 32(7), 262-265.
- Specchio, L. M., Iudice, A., Specchio, N., La Neve, A., Spinelli, A., Galli, R., Murri, L. (2004). Citalopram as treatment of depression in patients with epilepsy. *Clinical Neuropharmacology*, 27(3), 133-136.
- Spina, E., Avenoso, A., Pollicino, A., Caputi, A., Fazio, A., & Pisani, F. (1993). Carbamazepine coadministration with fluoxetine or fluvoxamine. *Therapeutic drug monitoring*, 15(3), 247-250.
- Steinert, T., & Froscher, W. (2017). Epileptic Seizures Under Antidepressive Drug Treatment: Systematic Review. *Pharmacopsychiatry*.
- Swedish Council on Health Technology, A. (2004). SBU Systematic Review Summaries. In *Treatment of Depression: A Systematic Review*. Stockholm: Swedish Council on Health Technology Assessment (SBU)
- Tedrus, G. M. A. S., Fonseca, L. C., Augusto, M. N., & Trindade, F. S. (2016). Major depressive episode, cognition, and epilepsy. *Epilepsy and Behavior*, 64, 219-223.
- Thompson, A. W., Miller, J. W., Katon, W., Chaytor, N., & Ciechanowski, P. (2009). Sociodemographic and clinical factors associated with depression in epilepsy. *Epilepsy and Behavior*, 14(4), 655-660.
- Trimble, M. (1984). Epilepsy, antidepressants, and the role of nomifensine. *Journal of Clinical Psychiatry*, 45(4ii), 39-42.
- Trimble, M. R. (1984). Epilepsy, antidepressants, and the role of nomifensine. *Journal of Clinical Psychiatry*, 45(4 II), 39-42.
- Turner, A. L., & Perry, M. S. (2017). Outside the box: Medications worth considering when traditional antiepileptic drugs have failed. *Seizure*, 50, 173-185.
- Van Cott, A. C., & Pugh, M. J. (2012). Potential anti-epileptic drug interaction with psychotropic medication in older adults with epilepsy. *Epilepsy Currents*, 12(1).
- Warner, N. M., Gwinn, R. P., & Doherty, M. J. (2016). Individualizing therapies with responsive epilepsy neurostimulation - A mirtazapine case study of hippocampal excitability. *Epilepsy Behav Case Rep*, 6, 70-72.
- Wen, X., Meador, K. J., Loring, D. W., Eisenschenk, S., Segal, R., & Hartzema, A. G. (2010). Is antiepileptic drug use related to depression and suicidal ideation among patients with epilepsy? *Epilepsy and Behavior*, 19(3), 494-500.
- Wiesmann, U. C., & Baker, G. A. (2013). Self-reported feelings of anger and aggression towards others in patients on levetiracetam: Data from the UK antiepileptic drug register. *BMJ Open*, 3(3).
- Wroblewski, B. A., Leary, J. M., Phelan, A. M., Whyte, J., & Manning, K. (1992). Methylphenidate and seizure frequency in brain injured patients with seizure disorders. *Journal of Clinical Psychiatry*, 53(3), 86-89.
- Zhu, S., Luo, L., & Gui, Y. (2004). Short-term efficacy of venlafaxine treating the depression in epilepsy patients. *Chinese Journal of Rehabilitation*, 19(2), 100-101.
- Zou, X., Zhou, J., & Long, Y. (2013). Efficacy and safety of antidepressants for depression in people with epilepsy: A systematic review. *Epilepsia*, 54, 67-68.

Anti-psykotiska läkemedel (19)

- Bhatti, M., Dorris, P., & Mehndiratta, P. (2017). Impact of psychotropic drugs on seizure threshold. *Neurology*, 88(16).
- Biemann, P., Levac, T., & Gagnon, M. A. (1978). Clonazepam: Its efficacy in association with phenytoin and phenobarbital in mental patients with generalized major motor seizures. *International Journal of Clinical Pharmacology Therapy and Toxicology*, 16(6), 268-273.
- Boon, M., & Ter Mors, B. (2014). Pharmacological treatment of interictal psychoses in patients with epilepsy, a systematic review. *Brain Injury*, 28(5-6), 646.

- Davison, K. (1965). EEG activation after intravenous amitriptyline. *Electroencephalography and clinical neurophysiology*, 19(3), 298-300.
- Eapen, V., & Gururaj, A. (2005). An open clinical trial of risperidone in children with seizures and hyperactivity. *Journal of Pediatric Neurology*, 3(1), 31-34.
- Erwin, C., Gerber, C., Morrison, S., & James, J. (1973). Lithium carbonate and convulsive disorders. *Archives of general psychiatry*, 28(5), 646-648.
- Farooq, S., & Sherin, A. (2008). Interventions for psychotic symptoms concomitant with epilepsy. *Cochrane Database Syst Rev*(4), Cd006118.
- Farooq, S., & Sherin, A. (2015). Interventions for psychotic symptoms concomitant with epilepsy. *Cochrane Database Syst Rev*(12), Cd006118.
- Gonzalez-Heydrich, J., Pandina, G. J., Fleisher, C. A., Hsin, O., Raches, D., Bourgeois, B. F., & Biederman, J. (2004). No seizure exacerbation from risperidone in youth with comorbid epilepsy and psychiatric disorders: A case series. *Journal of Child and Adolescent Psychopharmacology*, 14(2), 295-310.
- Kamm, I., & Mandel, A. (1967). Thioridazine in the treatment of behavior disorders in epileptics. *Diseases of the nervous system*, 28(1), 46-48.
- Nielsen, J., Correll, C. U., Manu, P., & Kane, J. M. (2013). Termination of clozapine treatment due to medical reasons: when is it warranted and how can it be avoided? *J Clin Psychiatry*, 74(6), 603-613; quiz 613.
- Okazaki M, Adachi N, Akanuma N, Hara K, Ito M, Kato M, et al. Do antipsychotic drugs increase seizure frequency in epilepsy patients? *Eur Neuropsychopharmacol* 2014; 24:1738-44.
- Ravanic, D., Djukic Dejanovic, S., Ristic Ignjatovic, D., Illic, M., Borovcanin, M., Babic, S., & Jovanovic, M. (2013). Treatment of psychotic episode in epilepsy-olanzapine vs risperidone. *European Psychiatry*, 28.
- Rettig, J. (1956). Chlorpromazine and Mysoline in the control of convulsive epilepsy in mentally deficient patients. *Journal of nervous and mental disease*, 124, 601-611.
- Robertson, M. M., & Trimble, M. R. (1985). The treatment of depression in patients with epilepsy. A double-blind trial. *Journal of Affective Disorders*, 9(2), 127-136.
- Sansoy, O. M., & Whorton, D. (1966). Chlordiazepoxide hydrochloride in treatment of convulsive disorders. A three-year retrospective study. *Rocky Mountain medical journal*, 63(12), 56-58.
- Spina, E., Avenoso, A., Pollicino, A. M., Caputi, A. P., Fazio, A., & Pisani, F. (1993). Carbamazepine coadministration with fluoxetine or fluvoxamine. *Ther Drug Monit*, 15(3), 247-250.
- Stoudemire, A., Nelson, A., & Houp, J. L. (1983). Interictal schizophrenia-like psychoses in temporal lobe epilepsy. *Psychosomatics*, 24(4), 331-333, 337-339.
- Zou, X., Zhou, J., & Long, Y. (2013). Efficacy and safety of antidepressants for depression in people with epilepsy: A systematic review. *Epilepsia*, 54, 67-68.

Centralstimulerande läkemedel (35)

- Adams, J., Alipio-Jocson, V., Inoyama, K., Bartlett, V., Sandhu, S., Oso, J., Meador, K. (2016). Effects of methylphenidate on cognition, quality of life, and seizure frequency in adults with epilepsy: A 1-month open-label trial. *Journal of Neuropsychiatry and Clinical Neurosciences*, 28(3), e33.
- Adams, J., Alipio-Jocson, V., Inoyama, K., Bartlett, V., Sandhu, S., Oso, J., Meador, K. (2017). Methylphenidate, cognition, and epilepsy: a double-blind, placebo-controlled, single-dose study. *Neurology*, 88(5), 470-476.
- Adams, J., Alipio-Jocson, V., Inoyama, K., Bartlett, V., Sandhu, S., Oso, J., Meador, K. J. (2017). Methylphenidate, cognition, and epilepsy: A 1-month open-label trial. *Epilepsia*.
- Artsy, E., McCarthy, D. C., Hurwitz, S., M, K. P., Dworetzky, B. A., & Lee, J. W. (2012). Use of modafinil in patients with epilepsy. *Epilepsy and Behavior*, 23(4), 405-408.
- Brunklaus, A., Dorris, L., & Zuberi, S. M. (2013). The impact of methylphenidate on seizure frequency and severity in children with attention-deficit-hyperactivity disorder and difficult-to-treat epilepsies. *Dev Med Child Neurol*, 55(10), 966-967.
- Caplan, R. (2017). ADHD in pediatric epilepsy: Fact or fiction? *Epilepsy Currents*, 17(2), 93-95.
- Downs, J., Giust, J., & Dunn, D. W. (2017). Considerations for ADHD in the child with epilepsy and the child with migraine. *Expert Rev Neurother*, 17(9), 861-869.
- Fosi, T., Lax-Pericall, M. T., Scott, R. C., Neville, B. G., & Aylett, S. E. (2013). Methylphenidate treatment of attention deficit hyperactivity disorder in young people with learning disability and difficult-to-treat epilepsy: Evidence of clinical benefit. *Epilepsia*, 54(12), 2071-2081.
- Gonzalez-Heydrich, J., Hsin, O., Gumlak, S., Kimball, K., Rober, A., Azeem, M. W., Biederman, J. (2014). Comparing stimulant effects in youth with ADHD symptoms and epilepsy. *Epilepsy Behav*, 36, 102-107.
- Gonzalez-Heydrich, J., Weiss, M., Connolly, M., Wambera, K., Jan, J. E., Plioplys, S., Kratochvil, C. J. (2006).

- Pharmacological management of a youth with ADHD and a seizure disorder. *J Am Acad Child Adolesc Psychiatry*, 45(12), 1527-1532.
- Gross, A., Devinsky, O., Westbrook, L. E., Wharton, A. H., & Alper, K. (2000). Psychotropic medication use in patients with epilepsy: Effect on seizure frequency. *Journal of Neuropsychiatry and Clinical Neurosciences*, 12(4), 458-464.
- Gross-Tsur, V., Manor, O., Meere, J., Joseph, A., & Shalev, R. (1997). Epilepsy and attention deficit hyperactivity disorder: Is methylphenidate safe and effective? *J Pediatr*, 130(1), 40-44.
- Gross-Tsur, V., Manor, O., van der Meere, J., Joseph, A., & Shalev, R. S. (1997). Epilepsy and attention deficit hyperactivity disorder: Is methylphenidate safe and effective? *J Pediatr*, 130(4), 670-674.
- Gucuyener, K., Erdemoglu, A. K., Senol, S., Serdaroglu, A., Soysal, S., & Kockar, A. I. (2003). Use of methylphenidate for attention-deficit hyperactivity disorder in patients with epilepsy or electroencephalographic abnormalities. *J Child Neurol*, 18(2), 109-112.
- Jesse, A., Alipio-Jocson, V., Inoyama, K., Bartlett, V., Sandhu, S., Oso, J., Meador, K. (2016). Methylphenidate's effects on cognition in patients with epilepsy: A randomized, double-blind, placebo-controlled, crossover, single-dose study. *Neurology*, 86(16).
- Kalil Neto, F., & Nunes, M. L. (2017). Evaluation of sleep organization in patients with attention deficit hyperactivity disorder (ADHD) and ADHD as a comorbidity of epilepsy. *Sleep Med*, 33, 91-96.
- Kanazawa, O. (2014). Reappraisal of abnormal EEG findings in children with ADHD: On the relationship between ADHD and epileptiform discharges. *Epilepsy and Behavior*, 41, 251-256.
- Koneski, J. A., Casella, E. B., Agertt, F., & Ferreira, M. G. (2011). Efficacy and safety of methylphenidate in treating ADHD symptoms in children and adolescents with uncontrolled seizures: a Brazilian sample study and literature review. *Epilepsy Behav*, 21(3), 228-232.
- McAfee, A. T., Holdridge, K. C., Johannes, C. B., Hornbuckle, K., & Walker, A. M. (2008). The effect of pharmacotherapy for attention deficit hyperactivity disorder on risk of seizures in pediatric patients as assessed in an insurance claims database. *Curr Drug Saf*, 3(2), 123-131.
- McAfee, A. T., Landon, J., Jones, M., Bangs, M. E., Acharya, N., Hornbuckle, K., & Wong, J. (2013). A cohort study of the risk of seizures in a pediatric population treated with atomoxetine or stimulant medications. *Pharmacoepidemiol Drug Saf*, 22(4), 386-393.
- Mercier, C., Roche, S., Gaillard, S., Kassai, B., Arzimanoglou, A., Herbillon, V., Rheims, S. (2016). Partial validation of a French version of the ADHD-rating scale IV on a French population of children with ADHD and epilepsy. Factorial structure, reliability, and responsiveness. *Epilepsy Behav*, 58, 1-6.
- Millichap, J. G., Millichap, J. J., & Stack, C. V. (2011). Utility of the electroencephalogram in attention deficit hyperactivity disorder. *Clin EEG Neurosci*, 42(3), 180-184.
- Moore, J. L., McAuley, J. W., Long, L., & Bornstein, R. (2002). An evaluation of the effects of methylphenidate on outcomes in adult epilepsy patients. *Epilepsy and Behavior*, 3(1), 92-95.
- Mulas, F., Roca, P., Ros-Cervera, G., Gandía-Benetó, R., & Ortiz-Sánchez, P. (2014). Pharmacological management of attention deficit hyperactivity disorder with methylphenidate and atomoxetine within a context of epilepsy. *Revista de Neurologia*, 58(SUPPL. 1), S43-S49.
- Radziuk, A. L., Kieling, R. R., Santos, K., Rotert, R., Bastos, F., & Palmini, A. L. (2015). Methylphenidate improves the quality of life of children and adolescents with ADHD and difficult-to-treat epilepsies. *Epilepsy Behav*, 46, 215-220.
- Ravi, M., & Ickowicz, A. (2016). Epilepsy, Attention-Deficit/Hyperactivity Disorder and Methylphenidate: Critical Examination of Guiding Evidence. *J Can Acad Child Adolesc Psychiatry*, 25(1), 50-58.
- Reed, V. A., Buitelaar, J. K., Anand, E., Day, K. A., Treuer, T., Upadhyaya, H. P., Savill, N. C. (2016). The Safety of Atomoxetine for the Treatment of Children and Adolescents with Attention-Deficit/Hyperactivity Disorder: A Comprehensive Review of Over a Decade of Research. *CNS Drugs*, 30(7), 603-628.
- Rheims, S., Herbillon, V., Villeneuve, N., Auvin, S., Napuri, S., Cancès, C., Barthez, M. A. (2016). ADHD in childhood epilepsy: Clinical determinants of severity and of the response to methylphenidate. *Epilepsia*, 57(7), 1069-1077.
- Santos K, Palmini A, Radziuk AL, Rotert R, Bastos F, Booij L, et al. The impact of methylphenidate on seizure frequency and severity in children with attention-deficit-hyperactivity disorder and difficult-to-treat epilepsies. *Dev Med Child Neurol* 2013; 55:654-60.
- Socanski, D., Aurlien, D., Herigstad, A., Thomsen, P. H., & Larsen, T. K. (2015). Attention deficit/hyperactivity disorder and interictal epileptiform discharges: Is it safe to use methylphenidate? *Seizure*, 25, 80-83.
- Torres, A., Whitney, J., Rao, S., Tilley, C., Lobel, R., & Gonzalez-Heydrich, J. (2011). Tolerability of atomoxetine for treatment of pediatric attention-deficit/hyperactivity disorder in the context of epilepsy. *Epilepsy Behav*, 20(1), 95-102.

- van der Feltz-Cornelis, C. M., & Aldenkamp, A. P. (2006). Corrigendum to "Effectiveness and safety of methylphenidate in adult attention deficit hyperactivity disorder in patients with epilepsy: An open treatment trial" *Epilepsy and Behavior*, 9(3), 542.
- van der Feltz-Cornelis, C. M., & Aldenkamp, A. P. (2006). Effectiveness and safety of methylphenidate in adult attention deficit hyperactivity disorder in patients with epilepsy: An open treatment trial. *Epilepsy and Behavior*, 8(3), 659-662.
- Wroblewski, B. A., Leary, J. M., Phelan, A. M., Whyte, J., & Manning, K. (1992). Methylphenidate and seizure frequency in brain injured patients with seizure disorders. *J Clin Psychiatry*, 53(3), 86-89.
- Yoo, H. K., Park, S., Wang, H. R., Lee, J. S., Kim, K., Paik, K. W., Ko, T. S. (2009). Effect of methylphenidate on the quality of life in children with epilepsy and attention deficit hyperactivity disorder: and open-label study using an osmotic-controlled release oral delivery system. *Epileptic Disord*, 11(4), 301-308.

Kirurgis behandling

Resektiv kirurgi (33)

- Bjellvi, J., Flink, R., Rydenhag, B., & Malmgren, K. (2015). Complications of epilepsy surgery in Sweden 1996-2010: a prospective, population-based study. *J Neurosurg*, 122(3), 519-525.
- Burkhardt, J. K., Neidert, M. C., Woernle, C. M., Bernays, R. L., Grunwald, T., & Wulf, M. A. (2013). Excessively elevated C-reactive protein after surgery for temporal lobe epilepsy. *Clin Neurol Neurosurg*, 115(8), 1245-1249.
- Cendes, F., Castellano, G., Pereira, A. R., Pereira, F. R., Yasuda, C. L., Tedeschi, H., Mantovani Guerreiro, C. A. (2010). Dynamic changes in white and gray matter volume are associated with outcome of surgical treatment in temporal lobe epilepsy. *Neuroimage*, 49(1), 71-79.
- Costa, J. C., Nunes, M. L., Almeida, A. G., & Palmini, A. L. (2010). Incidence of SUDEP in a cohort of patients with refractory epilepsy: the role of surgery and lesion localization. *Arq Neuropsiquiatr*, 68(6), 898-902.
- Ding, P., Liang, S., Zhang, S., Zhang, J., Hu, X., & Yu, X. (2016). Resective surgery combined with corpus callosotomy for children with non-focal lesional Lennox-Gastaut syndrome. *Acta Neurochir (Wien)*, 158(11), 2177-2184.
- Elger, C. E., Reuber, M., Helmstaedter, C., Kurthen, M., & Lux, S. (2003). Chronic epilepsy and cognition: A longitudinal study in temporal lobe epilepsy. *Annals of Neurology*, 54(4), 425-432.
- Health Quality. Ontario Epilepsy surgery: an evidence summary. (2012). *Ont Health Technol Assess Ser*, 12(17), 1-28
- Ismail, R., Mikati, M. A., Rahi, A. C., Comair, Y., & Faour, R. (2004). Effects of epilepsy surgery on quality of life: a controlled study in a Middle Eastern population. *Epilepsy Behav*, 5(1), 72-80.
- Jobst BC, Cascino GD. Resective epilepsy surgery for drug-resistant focal epilepsy: a review. *JAMA*. 2015 Jan 20;313(3):285-93.
- Johnston, J. M., Bertrand, M. E., Smyth, M. D., Limbrick, D. D., Dorward, I. G., & Titus, J. B. (2011). Extratemporal, nonlesional epilepsy in children: postsurgical clinical and neurocognitive outcomes. *J Neurosurg Pediatr*, 7(2), 179-188.
- Kelly, K., Smith, M. L., Olds, J., Whiting, S., Snyder, T., Elliott, I. M., & Kadis, D. S. (2011). Self-reported symptoms of psychological well-being in young adults who underwent resective epilepsy surgery in childhood. *Epilepsia*, 52(5), 891-899.
- Kumlien, E., Doss, R. C., & Gates, J. R. (2002). Treatment outcome in patients with mesial temporal sclerosis. *Seizure*, 11(7), 413-417.
- Lach, L., Elliott, I. M., & Lou Smith, M. (2006). Memory outcome after pediatric epilepsy surgery: objective and subjective perspectives. *Child Neuropsychol*, 12(3), 151-164.
- Loddenkemper, T., An, S., & Sanchez Fernandez, I. (2015). Pediatric refractory epilepsy: A decision analysis comparing medical versus surgical treatment. *Epilepsia*, 56(2), 263-272.
- Nilsson, L., Ahlbom, A., Tomson, T., & Farahmand, B. Y. (2003). Mortality in a population-based cohort of epilepsy surgery patients. *Epilepsia*, 44(4), 575-581.
- Otsuki, T., Kim, H. D., Luan, G., Inoue, Y., Baba, H., Oguni, H., . . . Sugai, K. (2016). Surgical versus medical treatment for children with epileptic encephalopathy in infancy and early childhood: Results of an international multicenter cohort study in Far-East Asia (the FACE study). *Brain Dev*, 38(5), 449-460.
- Puwanant, A., Alshekhlee, A., Jehi, L., & Koubeissi, M. Z. (2009). In-hospital complications of epilepsy surgery: a six-year nationwide experience. *Br J Neurosurg*, 23(5), 524-529.
- Radhakrishnan, K., & Rao, M. B. (2000). Is epilepsy surgery possible in countries with limited resources? *Epilepsia*, 41, S31-34.

- Radhakrishnan, K., Sarma, P. S., Abraham, M., Radhakrishnan, A., Kesavadas, C., & Dash, G. K. (2012). An audit of the presurgical evaluation and patient selection for extratemporal resective epilepsy surgery in a resource-poor country. *Seizure*, 21(5), 361-366.
- Shamseddine, A., Mikati, M. A., Sinno, D., Ataya, N., Comair, Y., Rahi, A., El-Banna, D. (2010). Quality of life after surgery for intractable partial epilepsy in children: a cohort study with controls. *Epilepsy Res*, 90(3), 207-213.
- Sinha, S., Duncan, J. S., Smith, S. J., Sander, J. W., Bell, G. S., Walker, M. C., Tisi, J. (2010). Premature mortality in refractory partial epilepsy: does surgical treatment make a difference? *J Neurol Neurosurg Psychiatry*, 81(7), 716-718.
- Smith, M. L., Kerr, E., & Law, N. (2015). Evaluation of behavioral outcomes in children 1 year after epilepsy surgery. *Epilepsia*, 56(10), 1605-1614.
- Smith, M. L., Lach, L., & Elliott, I. M. (2004). Cognitive, psychosocial, and family function one year after pediatric epilepsy surgery. *Epilepsia*, 45(6), 650-660.
- Smith, M. L., Lach, L., Elliott, I. M., & Kadis, D. S. (2008). Psychosocial outcomes in children two years after epilepsy surgery: has anything changed? *Epilepsia*, 49(4), 634-641.
- Smith, M. L., Olds, J., Lach, L., Whiting, S., McCleary, L., Snyder, T., Kadis, D. S. (2012). Quality of life in young adults who underwent resective surgery for epilepsy in childhood. *Epilepsia*, 53(9), 1577-1586.
- Smith, M. L., & Puka, K. (2016). Where are they now? Psychosocial, educational, and vocational outcomes after epilepsy surgery in childhood. *Epilepsia*, 57(4), 574-581.
- Sperling, M. R., Nei, M., Asadi-Pooya, A. A., & Barshow, S. (2016). A reappraisal of mortality after epilepsy surgery. *Neurology*, 86(21), 1938-1944.
- Stavem K, Bjørnaes H, Langmoen IA. Long-term seizures and quality of life after epilepsy surgery compared with matched controls. *Neurosurgery*. 2008 Feb;62(2):326-34; discussion 334-5.
- Tekcan, A., Aydemir, N., Canbeyli, R., & Ozkara, C. (2004). Changes in quality of life and self-perspective related to surgery in patients with temporal lobe epilepsy. *Epilepsy Behav*, 5(5), 735-742.
- Tracy, J., Evans, J., Skidmore, C., Sharan, A., Sperling, M., Pustina, D., & Doucet, G. (2014). Distinct types of white matter changes are observed after anterior temporal lobectomy in epilepsy. *PLoS One*, 9(8), e104211.
- Vickrey, B. G., Le, M., Kraemer, S., Rausch, R., Pietras, C. J., & Passaro, E. A. (2003). Early and late cognitive changes following temporal lobe surgery for epilepsy. *Neurology*, 60(6), 951-959.
- West, S., Nolan, S. J., Cotton, J., Gandhi, S., Weston, J., Sudan, A., Newton, R. (2015). Surgery for epilepsy. *Cochrane Database Syst Rev*(7), Cd010541.
- Wiebe, S., Eliasziw, M., Dykeman, J., Parrent, A., & Fiest, K. (2013). Impact of seizure frequency on improvement in quality of life in medically and surgically treated patients with epilepsy: Results from a randomized controlled trial. *Epilepsia*, 54, 287.
- Yasuda CL, Tedeschi H, Oliveira EL, Ribas GC, Costa AL, Cardoso TA, Montenegro MA, Guerreiro CA, Guerreiro MM, Li LM, Cendes F. Comparison of short-term outcome between surgical and clinical treatment in temporal lobe epilepsy: a prospective study. *Seizure*. 2006 Jan;15(1):35-40. Epub 2005 Dec 5.

[Resektiv kirurgi tidsfaktor \(62\)](#)

- Aberastury, M., Comas, B., Garcia, M., Besocke, A., Ciraolo, C., Agosta, G., & Silva, W. (2016). Epilepsy surgery in children and adolescents: Report on 43 cases. *Arch Argent Pediatr*, 114(5), 458-463.
- Adry, R. A., Meguins, L. C., da Silva Junior, S. C., Pereira, C. U., de Araujo Filho, G. M., & Marques, L. H. (2016). Factors predicting the outcome following surgical treatment of mesial temporal epilepsy due to mesial temporal sclerosis. *Acta Neurochir (Wien)*, 158(12), 2355-2363.
- Ansari, S. F., Tubbs, R. S., Terry, C. L., & Cohen-Gadol, A. A. (2010). Surgery for extratemporal nonlesional epilepsy in adults: an outcome meta-analysis. [S (review) (TN) X (KWR)]. *Acta Neurochir (Wien)*, 152(8), 1299-1305.
- Aronica, E., Leenstra, S., van Veelen, C. W., van Rijen, P. C., Hulsebos, T. J., Tersmette, A. C., . . . Troost, D. (2001). Glioneuronal tumors and medically intractable epilepsy: a clinical study with long-term follow-up of seizure outcome after surgery. [X (tidsjmf görs men i cancerpatienter) (TN)]. *Epilepsy Res*, 43(3), 179-191.
- Aull-Watschinger, S., Pataraia, E., Czech, T., & Baumgartner, C. (2008). Outcome predictors for surgical treatment of temporal lobe epilepsy with hippocampal sclerosis. [C (onset jmf inte) (TN) X - inkludera (omvänt OR) (KWR)]. *Epilepsia*, 49(8), 1308-1316.
- Barba, C., Mai, R., Grisotto, L., Gozzo, F., Pellacani, S., Tassi, L., . . . Guerrini, R. (2017). Unilobar surgery for symptomatic epileptic spasms. *Ann Clin Transl Neurol*, 4(1), 36-45.
- Behdad, A., Limbrick, D. D., Jr., Bertrand, M. E., & Smyth, M. D. (2009). Epilepsy surgery in children with seizures arising from the rolandic cortex. *Epilepsia*, 50(6), 1450-1461.

- Binder, D. K., Von Lehe, M., Kral, T., Bien, C. G., Urbach, H., Schramm, J., & Clusmann, H. (2008). Surgical treatment of occipital lobe epilepsy. [X (KI) (TN)]. *J Neurosurg*, 109(1), 57-69.
- Boesebeck, F., Janszky, J., Kellinghaus, C., May, T., & Ebner, A. (2007). Presurgical seizure frequency and tumoral etiology predict the outcome after extratemporal epilepsy surgery. *J Neurol*, 254(8), 996-999.
- Burneo, J. G., Black, L., Martin, R., Devinsky, O., Pacia, S., Faught, E., . . . Kuzniecky, R. I. (2006). Race/ethnicity, sex, and socioeconomic status as predictors of outcome after surgery for temporal lobe epilepsy. [O (jmf mellan raser men finns data på tid) (TN) X (OR finns, men ngt otydligt) (KWR)]. *Arch Neurol*, 63(8), 1106-1110.
- Cats, E. A., Kho, K. H., Van Nieuwenhuizen, O., Van Veelen, C. W., Gosselaar, P. H., & Van Rijen, P. C. (2007). Seizure freedom after functional hemispherectomy and a possible role for the insular cortex: the Dutch experience. *J Neurosurg*, 107(4), 275-280.
- Chen, H. H., Chen, C., Hung, S. C., Liang, S. Y., Lin, S. C., Hsu, T. R., . . . Shih, Y. H. (2014). Cognitive and epilepsy outcomes after epilepsy surgery caused by focal cortical dysplasia in children: early intervention maybe better. *Childs Nerv Syst*, 30(11), 1885-1895.
- Chugani, H. T., Ilyas, M., Kumar, A., Juhasz, C., Kupsky, W. J., Sood, S., & Asano, E. (2015). Surgical treatment for refractory epileptic spasms: The Detroit series. [X (retrospektiv) (TN)]. *Epilepsia*, 56(12), 1941-1949.
- Dammann, P., Wrede, K., Jabbarli, R., Neuschulte, S., Menzler, K., Zhu, Y., . . . Sure, U. (2016). Outcome after conservative management or surgical treatment for new-onset epilepsy in cerebral cavernous malformation. *J Neurosurg*, 0, 1-9.
- Elsharkawy, A. E., Alabbasi, A. H., Pannek, H., Oppel, F., Schulz, R., Hoppe, M., . . . Ebner, A. (2009). Long-term outcome after temporal lobe epilepsy surgery in 434 consecutive adult patients. *J Neurosurg*, 110(6), 1135-1146.
- Elsharkawy, A. E., Alabbasi, A. H., Pannek, H., Schulz, R., Hoppe, M., Pahs, G., . . . Ebner, A. (2008). Outcome of frontal lobe epilepsy surgery in adults. *Epilepsy Res*, 81(2), 97-106.
- Elsharkawy, A. E., El-Ghandour, N. M., Oppel, F., Pannek, H., Schulz, R., Hoppe, M., . . . Ebner, A. (2009). Long-term outcome of lesional posterior cortical epilepsy surgery in adults. *J Neurol Neurosurg Psychiatry*, 80(7), 773-780.
- Englot, D. J., Berger, M. S., Barbaro, N. M., & Chang, E. F. (2011). Predictors of seizure freedom after resection of supratentorial low-grade gliomas. A review. [S (KI) (review) (TN) X (analys och siffror finns) (KWR)]. *J Neurosurg*, 115(2), 240-244.
- Englot, D. J., Berger, M. S., Barbaro, N. M., & Chang, E. F. (2012). Factors associated with seizure freedom in the surgical resection of glioneuronal tumors. [X (TN)]. *Epilepsia*, 53(1), 51-57.
- Englot, D. J., Han, S. J., Lawton, M. T., & Chang, E. F. (2011). Predictors of seizure freedom in the surgical treatment of supratentorial cavernous malformations. *J Neurosurg*, 115(6), 1169-1174.
- Garcia Gracia, C., Yardi, R., Kattan, M. W., Nair, D., Gupta, A., Najm, I., . . . Jehi, L. (2015). Seizure freedom score: a new simple method to predict success of epilepsy surgery. *Epilepsia*, 56(3), 359-365.
- Grivas, A., Schramm, J., Kral, T., von Lehe, M., Helmstaedter, C., Elger, C. E., & Clusmann, H. (2006). Surgical treatment for refractory temporal lobe epilepsy in the elderly: seizure outcome and neuropsychological sequels compared with a younger cohort. [X (TN)]. *Epilepsia*, 47(8), 1364-1372.
- Hu, W. H., Ge, M., Zhang, K., Meng, F. G., & Zhang, J. G. (2012). Seizure outcome with surgical management of epileptogenic ganglioglioma: a study of 55 patients. *Acta Neurochir (Wien)*, 154(5), 855-861.
- Huang, C., Li, H., Chen, M., Si, Y., & Lei, D. (2014). Factors associated with preoperative and postoperative epileptic seizure in patients with cerebral ganglioglioma. *Pak J Med Sci*, 30(2), 245-249.
- Iida, K., Kagawa, K., Kartagiri, M., Hashizume, A., & Kurisu, K. (2015). Appropriating surgical intervention in mesial temporal-lobe epilepsy (MTLE) patients: Comparison of parameters influencing timing of surgery in patients with epilepsies with different evidence levels for surgery. *Epilepsy Currents*, 15, 355-356.
- Jayalakshmi, S., Vooturi, S., Vadapalli, R., Somayajula, S., Madigubba, S., & Panigrahi, M. (2015). Outcome of surgery for temporal lobe epilepsy in adults - A cohort study. [X (KI) (TN)]. *Int J Surg*, 0.
- Jayalakshmi, S., Vooturi, S., Vadapalli, R., Somayajula, S., Madigubba, S., & Panigrahi, M. (2016). Outcome of surgery for temporal lobe epilepsy in adults - A cohort study. *Int J Surg*, 36, 443-447.
- Kanchanatawan, B., Limothai, C., Srikiwilaikul, T., & Maes, M. (2014). Clinical predictors of 2-year outcome of resective epilepsy surgery in adults with refractory epilepsy: a cohort study. [X (visa olika Engel gr efter kirurgi) (TN)]. *BMJ Open*, 4(4), e004852.
- Khoury, J. S., Winokur, R. S., Tracy, J. I., & Sperling, M. R. (2005). Predicting seizure frequency after epilepsy surgery. [X (TN)]. *Epilepsy Res*, 67(3), 89-99.
- Kim, J. H., Lee, J. M., Kang, E., Kim, J. S., Song, I. C., & Chung, C. K. (2010). Functional reorganization associated with semantic language processing in temporal lobe epilepsy patients after anterior temporal lobectomy : a longitudinal functional magnetic resonance image study. [X? (kan ev räkna ut från tabell) (TN)]. *J Korean Neurosurg Soc*, 47(1), 17-25.

- Kim, Y. D., Heo, K., Park, S. C., Huh, K., Chang, J. W., Choi, J. U., . . . Lee, B. I. (2005). Antiepileptic drug withdrawal after successful surgery for intractable temporal lobe epilepsy. *Epilepsia*, 46(2), 251-257.
- Krsek, P., Jahodova, A., Kyncl, M., Kudr, M., Komarek, V., Jezdik, P., . . . Duchowny, M. (2013). Predictors of seizure-free outcome after epilepsy surgery for pediatric tuberous sclerosis complex. *Epilepsia*, 54(11), 1913-1921.
- Kumar, A., Valentin, A., Humayon, D., Longbottom, A. L., Jimenez-Jimenez, D., Mullatti, N., . . . Alarcon, G. (2013). Preoperative estimation of seizure control after resective surgery for the treatment of epilepsy. [X? (ingen info and AED resistens, Fig 1 inga siffror där datan finns) (TN)]. *Seizure*, 22(10), 818-826.
- Kumlien, E., Doss, R. C., & Gates, J. R. (2002). Treatment outcome in patients with mesial temporal sclerosis. *Seizure*, 11(7), 413-417.
- Kwon, C. S., Sheth, S. A., Walcott, B. P., Neal, J., Eskandar, E. N., & Ogilvy, C. S. (2013). Long-term seizure outcomes following resection of supratentorial cavernous malformations. *Clin Neurol Neurosurg*, 115(11), 2377-2381.
- Liava, A., Francione, S., Tassi, L., Lo Russo, G., Cossu, M., Mai, R., . . . Dalla Bernardina, B. (2012). Individually tailored extratemporal epilepsy surgery in children: anatomo-electro-clinical features and outcome predictors in a population of 53 cases. *Epilepsy Behav*, 25(1), 68-80.
- Lowe, N. M., Eldridge, P., Varma, T., & Wieschmann, U. C. (2010). The duration of temporal lobe epilepsy and seizure outcome after epilepsy surgery. [X (TN)]. *Seizure*, 19(5), 261-263.
- Maehara, T., Inaji, M., & Matsuura, M. (2013). Surgical effects of focus resection for patients with intractable epilepsy. [X (TN)]. *Neurol Med Chir (Tokyo)*, 53(5), 281-286.
- Meguins, L. C., Adry, R. A., Silva-Junior, S. C., Araujo Filho, G. M., & Marques, L. H. (2015). Shorter epilepsy duration is associated with better seizure outcome in temporal lobe epilepsy surgery. *Arq Neuropsiquiatr*, 73(3), 212-217.
- Mueller, C. A., Scorzin, J., von Lehe, M., Fimmers, R., Helmstaedter, C., Zentner, J., . . . Schramm, J. (2012). Seizure outcome 1 year after temporal lobe epilepsy: an analysis of MR volumetric and clinical parameters. [X (TN)]. *Acta Neurochir (Wien)*, 154(8), 1327-1336.
- Parsa, A. T., Tihan, T., Fang, S., Barry, J. J., Han, S. J., Barbaro, N. M., . . . Chang, E. F. (2011). Early surgical intervention in adult patients with ganglioglioma is associated with improved clinical seizure outcomes. [X (KWR)]. *J Clin Neurosci*, 18(1), 29-33.
- Pati, S., Deep, A., Troester, M. M., Kossoff, E. H., & Ng, Y. T. (2013). Lennox-Gastaut syndrome symptomatic to hypothalamic hamartoma: evolution and long-term outcome following surgery. [X (TN)]. *Pediatr Neurol*, 49(1), 25-30.
- Pinheiro-Martins, A. P., Bianchin, M. M., Velasco, T. R., Terra, V. C., Araujo, D., Jr., Wichert-Ana, L., . . . Sakamoto, A. C. (2012). Independent predictors and a prognostic model for surgical outcome in refractory frontal lobe epilepsy. *Epilepsy Res*, 99(1), 55-63.
- Radhakrishnan, A., Menon, R., Menon, D., Singh, A., Radhakrishnan, N., Vilanilam, G., . . . Thomas, S. V. (2016). Early resective surgery causes favorable seizure outcome in malformations of cortical development. [X (TN)]. *Epilepsy Res*, 124, 1-11.
- Ramirez-Molina, J. L., Di Giacomo, R., Mariani, V., Deleo, F., Cardinale, F., Uscategui-Daccarett, A. M., . . . Tassi, L. (2017). Surgical outcomes in two different age groups with Focal Cortical Dysplasia type II: Any real difference? *Epilepsy Behav*, 70, 45-49.
- Schramm, J., Kral, T., Grunwald, T., & Blumcke, I. (2001). Surgical treatment for neocortical temporal lobe epilepsy: clinical and surgical aspects and seizure outcome. *J Neurosurg*, 94(1), 33-42.
- Schulz, R., Hoppe, M., Boesebeck, F., Gyimesi, C., Pannek, H. W., Woermann, F. G., . . . Ebner, A. (2011). Analysis of reoperation in mesial temporal lobe epilepsy with hippocampal sclerosis. [X? (KI) (ev räkna från tabell) (TN)]. *Neurosurgery*, 68(1), 89-97; discussion 97.
- Siegel, A. M., Cascino, G. D., Meyer, F. B., McClelland, R. L., So, E. L., Marsh, W. R., . . . Sharbrough, F. W. (2004). Resective reoperation for failed epilepsy surgery: seizure outcome in 64 patients. *Neurology*, 63(12), 2298-2302.
- Sirven, J. I., Malamut, B. L., O'Connor, M. J., & Sperling, M. R. (2000). Temporal lobectomy outcome in older versus younger adults. [X? (KI) (kanske går att anv) (TN)]. *Neurology*, 54(11), 2166-2170.
- Smyth, M. D., Limbrick, D. D., Jr., Ojemann, J. G., Zempel, J., Robinson, S., O'Brien, D. F., . . . Park, T. S. (2007). Outcome following surgery for temporal lobe epilepsy with hippocampal involvement in preadolescent children: emphasis on mesial temporal sclerosis. *J Neurosurg*, 106(3), 205-210.
- Souza-Oliveira, C., Escorsi-Rosset, S., Bianchin, M. M., Terra, V. C., Wichert-Ana, L., Machado, H. R., & Sakamoto, A. C. (2009). Comparative role of neuropsychological testing in the presurgical evaluation of children with medically intractable epilepsies. [X (TN)]. *Childs Nerv Syst*, 25(7), 875-880.
- Spencer, S. S., Schramm, J., Wyler, A., O'Connor, M., Orbach, D., Krauss, G., . . . Westerveld, M. (2002). Multiple subpial transection for intractable partial epilepsy: an international meta-analysis. [X? (metaanalys, beskriver i text men inga uträkningar?) (TN)]. *Epilepsia*, 43(2), 141-145.

- Stavrou, I., Baumgartner, C., Frischer, J. M., Trattnig, S., & Knosp, E. (2008). Long-term seizure control after resection of supratentorial cavernomas: a retrospective single-center study in 53 patients. *Neurosurgery*, 63(5), 888-896; discussion 897.
- Tigaran, S., Cascino, G. D., McClelland, R. L., So, E. L., & Richard Marsh, W. (2003). Acute postoperative seizures after frontal lobe cortical resection for intractable partial epilepsy. *Epilepsia*, 44(6), 831-835.
- Vale, F. L., Effio, E., Arredondo, N., Bozorg, A., Wong, K., Martinez, C., . . . Benbadis, S. R. (2012). Efficacy of temporal lobe surgery for epilepsy in patients with negative MRI for mesial temporal lobe sclerosis. [X (TN)]. *J Clin Neurosci*, 19(1), 101-106.
- Vale, F. L., Vivas, A. C., Manwaring, J., Schoenberg, M. R., & Benbadis, S. R. (2015). Temporal lobe epilepsy and cavernous malformations: surgical strategies and long-term outcomes. *Acta Neurochir (Wien)*, 157(11), 1887-1895; discussion 1895.
- Wang, C., Yu, X., Shrestha, S., Qian, C., Wang, L., & Chen, G. (2016). A predicted model for postoperative seizure outcomes after the surgical resection of supratentorial cavernous malformations. *Medicine (Baltimore)*, 95(26), e4078.
- Wieser, H. G., Ortega, M., Friedman, A., & Yonekawa, Y. (2003). Long-term seizure outcomes following amygdalohippocampectomy. *J Neurosurg*, 98(4), 751-763.
- Wiesemann, U. C., Larkin, D., Varma, T., & Eldridge, P. (2008). Predictors of outcome after temporal lobectomy for refractory temporal lobe epilepsy. [X (TN)]. *Acta Neurol Scand*, 118(5), 306-312.
- von der Breie, C., Kuczaty, S., & von Lehe, M. (2014). Surgical management and long-term outcome of pediatric patients with different subtypes of epilepsy associated with cerebral cavernous malformations. *J Neurosurg Pediatr*, 13(6), 699-705.
- Wong, C. H., Bleasel, A., Wen, L., Eberl, S., Byth, K., Fulham, M., . . . Mohamed, A. (2010). The topography and significance of extratemporal hypometabolism in refractory mesial temporal lobe epilepsy examined by FDG-PET. [X (TN)]. *Epilepsia*, 51(8), 1365-1373.
- Yeon, J. Y., Kim, J. S., Choi, S. J., Seo, D. W., Hong, S. B., & Hong, S. C. (2009). Supratentorial cavernous angiomas presenting with seizures: surgical outcomes in 60 consecutive patients. [X (TN)]. *Seizure*, 18(1), 14-20.

Annan behandling

Ketogen kost (121)

- Akhondian, J., Moayyedpour, A., Raouf Ziai, M., Khaje Dalui, M., & Hosseini, H. (2009). The effect of thymoquinone on pediatric intractable seizures. *Epilepsia*, 50, 95.
- Akhondian, J., Parsa, A., & Rakhshande, H. (2007). The effect of Nigella sativa L. (black cumin seed) on intractable pediatric seizures. *Medical Science Monitor*, 13(12), CR555-CR559.
- Al Khayat, H. A., Awadalla, M. M., Al Wakad, A., & Marzook, Z. A. (2010). Polyunsaturated fatty acids in children with idiopathic intractable epilepsy: Serum levels and therapeutic response. *Journal of Pediatric Neurology*, 8(2), 175-185.
- Alshehri, A. A. A. A., Bokhari, R. F., Kojan, S., Farwana, M., & Murad, M. (2014). Intravenous versus non-intravenous benzodiazepines for the abortion of seizures: A systematic review and meta-analysis of randomized controlled trials. *Canadian Journal of Neurological Sciences*, 41, S26-S27.
- Anthony, J., & Rush, R. (1998). A randomised controlled trial of intravenous magnesium sulphate versus placebo. *Br J Obstet Gynaecol*, 105(7), 810.
- Araya-Quintanilla, F., Celis-Rosati, A., Rodriguez-Leiva, C., Silva-Navarro, C., Silva-Pinto, Y., & Toro-Jeria, B. (2016). [Effectiveness of a ketogenic diet in children with refractory epilepsy: a systematic review]. *Revista de Neurologia*, 62(10), 439-448.
- Auvin, S. (2010). Ketogenic diet in infantile spasms: Time for new perspectives. *Future Neurology*, 5(5), 653-656.
- Auvin, S. (2016). Non-pharmacological medical treatment in pediatric epilepsies. *Revue Neurologique*, 172(3), 182-185.
- Avula, S., Parikh, S., Demarest, S., Kurz, J., & Gropman, A. (2014). Treatment of mitochondrial disorders. *Current Treatment Options in Neurology*, 16(6).
- Bahassan, N. A., & Jan, M. M. (2006). Ketogenic diet: Update and application. *Neurosciences*, 11(4), 235-240.
- Barzegar, M., Irandoust, P., & Mameghani, M. E. (2010). A modified Atkins diet for intractable childhood epilepsy. *Iranian Journal of Child Neurology*, 4(3), 15-20.
- Bergqvist, A. G., Schall, J. I., Gallagher, P. R., Cnaan, A., & Stallings, V. A. (2005). Fasting versus gradual initiation of the ketogenic diet: a prospective, randomized clinical trial of efficacy. *Epilepsia*, 46(11), 1810-1819.

- Bergqvist, A. G. C., Schall, J. I., & Stallings, V. A. (2007). Vitamin D status in children with intractable epilepsy, and impact of the ketogenic diet. *Epilepsia*, 48(1), 66-71.
- Brigo, F., & Igwe, S. C. (2015). Antiepileptic drugs for the treatment of infants with severe myoclonic epilepsy. *Cochrane Database of Systematic Reviews*, 10, CD010483.
- Camfield, P., & Camfield, C. (2003). What is new in paediatric epilepsy? *Paediatrics and Child Health*, 8(9), 557-560.
- Caraballo, R. H., Cersósimo, R. O., Sakr, D., Cresta, A., Escobal, N., & Fejerman, N. (2006). Ketogenic diet in patients with myoclonic-astatic epilepsy. *Epileptic Disorders*, 8(2), 151-155.
- Casas-Fernandez, C., Martínez-Bermejo, A., Rufo-Campos, M., Smeyers-Dura, P., Alarcón-Martínez, H., Ibáñez-Mico, S., Campos-Castello, J. (2011). Efficacy and tolerability of oral lacosamide (LCM) as adjunctive therapy in 130 pediatric patients with pharmacoresistant focal epilepsy. *Epilepsia*, 52, 75.
- Cervenka, M. C., Terao, N. N., Bosarge, J. L., Henry, B. J., Klees, A. A., Morrison, P. F., & Kossoff, E. H. (2012). Email management of the modified Atkins diet for adults with epilepsy is feasible and effective. *Epilepsia*, 53(4), 728-732.
- Chung, S., Hogan, R., Holmay, M., Yu, V., Nguyen, H., Blatt, I., & Halvorsen, M. (2016). Development of unique dosage form and differentiating features of USL255, Qudexy® XR (topiramate) extended-release capsules. *Neurology*, 86(16).
- Cole, A. J. (2003). Evaluating the efficacy of antiepileptic treatments: Introduction. *Epilepsia*, 44, 1.
- Coppola, G., Epifanio, G., Auricchio, G., Federico, R. R., Resicato, G., & Pascotto, A. (2006). Plasma free carnitine in epilepsy children, adolescents and young adults treated with old and new antiepileptic drugs with or without ketogenic diet. *Brain and Development*, 28(6), 358-365.
- Coppola, G., Veggiotti, P., Cusmai, R., Bertoli, S., Cardinali, S., Dionisi-Vici, C., Pascotto, A. (2002). The ketogenic diet in children, adolescents and young adults with refractory epilepsy: An Italian multicentric experience. *Epilepsy Res*, 48(3), 221-227.
- Coppola, G., Verrotti, A., D'Aniello, A., Arcieri, S., Operto, F. F., Della Corte, R., Pascotto, A. (2010). Valproic acid and phenobarbital blood levels during the first month of treatment with the ketogenic diet. *Acta Neurol Scand*, 122(4), 303-307.
- DeGiorgio, C. M., Miller, P. R., Harper, R., Gornbein, J., Schrader, L., Soss, J., & Meymandi, S. (2015). Fish oil (n-3 fatty acids) in drug resistant epilepsy: a randomised placebo-controlled crossover study. *J Neurol Neurosurg Psychiatry*, 86(1), 65-70.
- de Kinderen RJ, Lambrechts DA, Wijnen BF, Postulart D, Aldenkamp AP, Majoe MH, et al. An economic evaluation of the ketogenic diet versus care as usual in children and adolescents with intractable epilepsy: An interim analysis. *Epilepsia* 2016; 57:41-50.
- DiMario Jr, F. J., & Holland, J. (2002). The ketogenic diet: A review of the experience at Connecticut Children's Medical Center. *Pediatric Neurology*, 26(4), 288-292.
- Dotson, R., Dickinson, L., Kang, H., & Dasheiff, R. (1985). Effect of simultaneously ingested milk on phenytoin bioavailability. *Neurology*, 35(10), 1526-1527.
- Dressler, A., Klaus, M. T., Muehlebner, A., Moerzinger, M., Pahs, G., Benninger, F., & Feucht, M. (2011). Lateralizing signs in infantile spasms: Differences in video and EEG analysis. *Epilepsia*, 52, 5.
- Dressler, A., Trimmel-Schwahafer, P., Reithofer, E., Groppel, G., Muhlebner, A., Samuel, S., Feucht, M. (2015). The ketogenic diet versus ACTH in the treatment of infantile spasms: A prospective randomised study. *European Journal of Paediatric Neurology*, 19, S17.
- Dunn, D. W., Johnson, C. S., Fastenau, P. S., Byars, A. W., DeGraww, T. J., Perkins, S. M., & Austin, J. K. (2011). Are AEDs associated with suicidal ideation? *Epilepsy Currents*, 11(1).
- El-Rashidy, O. F., Nassar, M. F., Abdel-Hamid, I. A., Shatla, R. H., Abdel-Hamid, M. H., Gabr, S. S., Shaaban, S. Y. (2013). Modified Atkins diet vs classic ketogenic formula in intractable epilepsy. *Acta Neurol Scand*, 128(6), 402-408.
- Eros, E., Geher, P., Gomor, B., & Czeizel, A. E. (1998). Epileptogenic activity of folic acid after drug induces SLE (folic acid and epilepsy). *Eur J Obstet Gynecol Reprod Biol*, 80(1), 75-78.
- Garg, S. K., Kumar, N., Bhargava, V. K., & Prabhakar, S. K. (1998). Effect of grapefruit juice on carbamazepine bioavailability in patients with epilepsy. *Clin Pharmacol Ther*, 64(3), 286-288.
- Ghofrani, M., & Akhondian, J. (2010). Intractable epilepsy in children. *Iranian Journal of Child Neurology*, 4(3), 7-14.
- Gibberd, F. B., Nicholls, A., & Wright, M. G. (1981). The influence of folic acid on the frequency of epileptic attacks. *Eur J Clin Pharmacol*, 19(1), 57-60.
- Giovannelli, F., Zaccara, G., Cincotta, M., Loiacono, G., & Verrotti, A. (2015). Characterization of the adverse events profile of placebo-treated patients in randomized controlled trials on drug-resistant focal epilepsies.

- Journal of Neurology*, 262(6), 1401-1406.
- Grant, R. H., & Stores, O. P. (1970). Folic acid in folate-deficient patients with epilepsy. *Br Med J*, 4(5736), 644-648.
- Gulati, S., & Chakrabarty, B. (2014). Dietary therapy in childhood epilepsy, an overview. *International Journal of Epilepsy*, 1(1), 27-35.
- Hamed, S. A. (2010). The rationale for neuroprotection in epilepsy: Steps forward for new therapeutic and preventive strategies. *Journal of Integrative Neuroscience*, 9(1), 65-102.
- Hancock, E. C., Osborne, J. P., & Edwards, S. W. (2008). Treatment of infantile spasms. *Cochrane Database of Systematic Reviews*, 0(4).
- Hancock, E. C., Osborne, J. P., & Edwards, S. W. (2009). Treatment of infantile spasms. *Cochrane Database of Systematic Reviews*, 0(4).
- Herzog, A. G. (2015). Catamenial epilepsy: current concepts of definition, prevalence, pathophysiology and treatment. *Zeitschrift fur Epileptologie*, 28(4), 295-303.
- Hsu, W., Sing, C., He, Y., Worsley, A., Wong, I., & Chan, E. (2013). Systematic review and meta-analysis of the safety of perampanel in the treatment of partial-onset epilepsy. *Drug Safety*, 36(9), 854-855.
- Hulshof, H. M., Carpay, J. A., Jansen, F. E., & Braun, K. P. J. (2014). The modified atkins diet in patients with refractory epilepsy and severe intellectual disability-design of a randomized controlled trial. *Epilepsia*, 55, 205.
- Jackson, C. F., Makin, S. M., Marson, A. G., & Kerr, M. (2015). Non-pharmacological interventions for people with epilepsy and intellectual disabilities. *Cochrane Database of Systematic Reviews*, 9, CD005502.
- Jeeja, M. C., Jayakrishnan, T., Narayanan, P. V., Kumar, M. S., Thejus, T., & Anilakumari, V. P. (2014). Folic acid supplementation on homocysteine levels in children taking antiepileptic drugs: A randomized controlled trial. *J Pharmacol Pharmacother*, 5(2), 93-99.
- Jefferson, J. W. (2009). Herbal psychopharmacology: Show me the data. *Psychopharm Rev.*, 44, 25-32.
- Jekovec-Vrhovsček, M., Kocijančič, A., & Preželj, J. (2000). Effect of vitamin D and calcium on bone mineral density in children with CP and epilepsy in full-time care. *Developmental Medicine and Child Neurology*, 42(6), 403-405.
- Johnson, J., Brittain, S., Keeler, E., Louro, D., & Baroldi, P. (2012). Oxcarbazepine extended-release (SPN-804) therapy in patients with refractory partial seizures: Review of neurologic and ophthalmologic adverse events. *Neurology*, 78(1).
- Kang, H. C., Lee, Y. J., Lee, J. S., Lee, E. J., Eom, S., You, S. J., & Kim, H. D. (2011). Comparison of short-versus long-term ketogenic diet for intractable infantile spasms. *Epilepsia*, 52(4), 781-787.
- Kannan, L., Gulati, S., Agarwala, A., Kabra, M., Tripathi, M., & Pandey, R. M. (2013). Efficacy of lowglycemic index diet therapy in children with refractory epilepsy: A randomized controlled trial. *Epilepsia*, 54, 229.
- Kim, J. A., Yoon, J. R., Lee, E., Lee, J. S., Kim, H. D., & Kang, H. C. (2015). Comparison of efficacy between a modified atkins diet and a classic ketogenic diet in childhood intractable epilepsy. *Epilepsy Currents*, 15, 95-96.
- King-Stephens, D., Mirro, E., Weber, P., Laxer, K., Van Ness, P., Salanova, V., Morrell, M. (2014). Lateralization of temporal lobe epilepsy with long-term ambulatory intracranial monitoring using the RNS™ system: Experience in 82 patients. *Epilepsy Currents*, 14, 459-460.
- Klein, P., Janousek, J., Barber, A., & Weissberger, R. (2010). Ketogenic diet treatment in adults with refractory epilepsy. *Epilepsy and Behavior*, 19(4), 575-579.
- Klepper, J., Scheffer, H., Leiendoeker, B., Gertsen, E., Binder, S., Leferink, M., Willemse, M. A. (2005). Seizure control and acceptance of the ketogenic diet in GLUT1 deficiency syndrome: A 2- to 5-year follow-up of 15 children enrolled prospectively. *Neuropediatrics*, 36(5), 302-308.
- Kossoff, E. (2013). Ghee whiz! The growing evidence for the benefits of the modified atkins diet. *Epilepsy Currents*, 13(5), 211-212.
- Kossoff, E. H., McGrogan, J. R., Bluml, R. M., Pillas, D. J., Rubenstein, J. E., & Vining, E. P. (2006). A modified Atkins diet is effective for the treatment of intractable pediatric epilepsy. *Epilepsia*, 47(2), 421-424.
- Kossoff, E. H., Rowley, H., Sinha, S. R., & Vining, E. P. G. (2008). A prospective study of the modified Atkins diet for intractable epilepsy in adults. *Epilepsia*, 49(2), 316-319.
- Kossoff, E. H., Turner, Z., Bluml, R. M., Pyzik, P. L., & Vining, E. P. (2007). A randomized, crossover comparison of daily carbohydrate limits using the modified Atkins diet. *Epilepsy Behav*, 10(3), 432-436.
- Kossoff, E. H., Turner, Z., Bluml, R. M., Pyzik, P. L., & Vining, E. P. G. (2007). A randomized, crossover comparison of daily carbohydrate limits using the modified Atkins diet. *Epilepsy and Behavior*, 10(3), 432-436.
- Kraemer, G., Trinka, E., & Werhahn, K. J. (2009). STEP ONE: Study on the treatment of elderly patients with

- older and newer anti-epileptic drugs. - Interim report on recruitment and drop-out rates. *Epilepsia*, 50, 62.
- Krämer, G., Werhahn, K. J., & Trinka, E. (2009). Step one: Study on the treatment of elderly patients with older and newer anti-epileptic drugs, interim report on recruitment and drop-out rates. *Journal of the Neurological Sciences*, 285, S256.
- Lambrechts, D., Kinderen, R. J. A., Vles, H. S. H., Louw, A. J., Aldenkamp, A. P., & Majoe, M. J. M. (2015). The MCT-ketogenic diet as a treatment option in refractory childhood epilepsy: A prospective study with 2-year follow-up. *Epilepsy & Behavior*, 51, 261-266.
- Lang, N., Lange, M., Schmitt, F. C., Bös, M., Weber, Y., Evers, S., Stephani, U. (2016). Intravenous lacosamide in clinical practice-Results from an independent registry. *Seizure*, 39, 5-9.
- Liu, L., & Henry, J. C. (2009). New-onset partial epilepsy in adults. *Current Treatment Options in Neurology*, 11(4), 242-252.
- Loewen, G., Mansbach, H., Shin, P., Leyco, A., Abou-Khalil, B., Armstrong, R., Rosenfeld, W. (2009). Steady-state pharmacokinetics of retigabine as adjunctive therapy in patients with epilepsy. *Epilepsia*, 50, 113.
- Lucarelli, S., Frediani, T., Lastrucci, G., Viscido, G., Federici, T., Spalice, A., Cucchiara, S. (2012). Efficacy of the cow milk proteins free diet in children suffering from rolandic epilepsy. *Digestive and Liver Disease*, 44, S252.
- Ma, L., Yung, A., Yau, E., & Kwong, K. (2010). Clinical guidelines on management of prolonged seizures, serial seizures and convulsive status epilepticus in children. *Hong Kong Journal of Paediatrics*, 15(1), 52-83.
- Mani, J. (2008). Temporal lobe epilepsy in children. *Journal of Pediatric Neurosciences*, 3(1), 2-6.
- Mann, D., Liu, J., Chew, M., Bockbrader, H., Alvey, C. W., Zegarac, E., Pitman, V. (2014). A placebo-controlled, escalating dose, multiple dose study to evaluate the safety, tolerability and pharmacokinetics of pregabalin in pediatric patients with partial onset seizures. *Epilepsy Currents*, 14, 437-438.
- Marsh, E. B., Freeman, J. M., Kossoff, E. H., Vining, E. P. G., Rubenstein, J. E., Pyzik, P. L., & Hemingway, C. (2006). The outcome of children with intractable seizures: A 3- to 6-year follow-up of 67 children who remained on the ketogenic diet less than one year. *Epilepsia*, 47(2), 425-430.
- Martin, K., Jackson, C. F., Levy, R. G., & Cooper, P. N. (2016). Ketogenic diet and other dietary treatments for epilepsy. *Cochrane Database of Systematic Reviews*, 2, CD001903.
- Miranda, M. J., Mortensen, M., Povlsen, J. H., Nielsen, H., & Beniczky, S. (2011). Danish study of a Modified Atkins diet for medically intractable epilepsy in children: Can we achieve the same results as with the classical ketogenic diet? *Seizure*, 20(2), 151-155.
- Müller, A., Helbig, I., Jansen, C., Bast, T., Guerrini, R., Jähn, J., Kluger, G. (2016). Retrospective evaluation of low long-term efficacy of antiepileptic drugs and ketogenic diet in 39 patients with CDKL5-related epilepsy. *European Journal of Paediatric Neurology*, 20(1), 147-151.
- Nair, D. R. (2016). Management of drug-resistant epilepsy. *CONTINUUM Lifelong Learning in Neurology*, 22(1), 157-172.
- Nathan, J. K., Purandare, A. S., Parekh, Z. B., & Manohar, H. V. (2009). Ketogenic diet in Indian children with uncontrolled epilepsy. *Indian Pediatrics*, 46(8), 669-673.
- Neal, E. G., Chaffe, H., Schwartz, R. H., Lawson, M. S., Edwards, N., Fitzsimmons, G., Cross, J. H. (2009). A randomized trial of classical and medium-chain triglyceride ketogenic diets in the treatment of childhood epilepsy. *Epilepsia*, 50(5), 1109-1117.
- Ng, Y., Glauser, T., Ferreira, J., Olhaye, O., Williams, B., Perdomo, C., & Bibbiani, F. (2016). Response durability analyses from a rufinamide pivotal trial in lennox-gastaut syndrome (LGS). *Neurology*, 86(16).
- Norris, J. W., & Pratt, R. F. (1971). A controlled study of folic acid in epilepsy. A controlled study of folic acid in epilepsy. *Neurology*, 21(6), 659-664.
- O'Callaghan, F. J. K., Edwards, S., Hancock, E., Johnson, A., Kennedy, C., Lux, A., Osborne, J. P. (2015). The International Collaborative Infantile Spasms Study (ICISS) comparing hormonal therapies (prednisolone or tetracosactide depot) and vigabatrin versus hormonal therapies alone in the treatment of infantile spasms: Early clinical outcome. *European Journal of Paediatric Neurology*, 19, S16-S17.
- Oguni, M., Inoue, N., Takahata, K., & Koseki, M. (2011). Is modified atkins diet therapy applicable for adults with intractable epilepsy patients? - a short trial in adult volunteers. *Epilepsy and Seizure*, 4(1), 7-14.
- Pellock, J., Haut, S., & Seinfeld, S. (2014). Benzodiazepine use for emergency treatment of seizures: A review. *Epilepsy Currents*, 14, 179.
- Pires, M. E., Illea, A., Bourel, E., Bellavoline, V., Merdariu, D., Berquin, P., & Auvin, S. (2013). Ketogenic diet for infantile spasms refractory to first-line treatments: An open prospective study. *Epilepsy Res*, 105(1), 189-194.
- Raju, K. N., Gulati, S., Kabra, M., Agarwala, A., Sharma, S., Pandey, R. M., & Kalra, V. (2011). Efficacy of 4:1 (classic) versus 2.5:1 ketogenic ratio diets in refractory epilepsy in young children: a randomized open

- labeled study. *Epilepsy Res*, 96(1-2), 96-100.
- Raju, K. N. V., Gulati, S., Kabra, M., Agarwala, A., Sharma, S., Pandey, R. M., & Kalra, V. (2011). Efficacy of 4:1 (classic) versus 2.5:1 ketogenic ratio diets in refractory epilepsy in young children: A randomized open labeled study. *Epilepsy Res*, 96(1), 96-100.
- Ralston, A. J., Snaith, R. P., & Hinley, J. B. (1970). Effects of folic acid on fit-frequency and behaviour in epileptics on anticonvulsants. *Lancet*, 1(7652), 867-868.
- Reda, D. M. A., Abd-El-Fatah, N. K., Omar, T. E. S. I., & Darwish, O. A. H. (2015). Fish oil intake and seizure control in children with medically resistant epilepsy. *North American Journal of Medical Sciences*, 7(7), 317-321.
- Rieger, J., Bähr, O., Maurer, G. D., Hattingen, E., Franz, K., Brucker, D., Steinbach, J. P. (2014). ERGO: A pilot study of ketogenic diet in recurrent glioblastoma. *International Journal of Oncology*, 44(6), 1843-1852.
- Rowan, A. J., Kelly, K. M., & Birnbaum, A. K. (2005). Epilepsy later in life: Managing the unique problems of seizures in the elderly. *Geriatrics*, 60(5), 3-20.
- Saengsuwan, J., Boonyaleepan, S., Tiamkao, S., & Integrated Epilepsy, G. (2015). Diet, exercise, sleep, sexual activity, and perceived stress in people with epilepsy in NE Thailand. *Epilepsy & Behavior*, 45, 39-43.
- Saneto, R. P., & Anderson, G. D. (2009). Onset of action and seizure control in Lennox-Gastaut syndrome: Focus on rufinamide. *Therapeutics and Clinical Risk Management*, 5(1), 271-280.
- Saxena, V. S., & Nadkarni, V. V. (2011). Nonpharmacological treatment of epilepsy. *Annals of Indian Academy of Neurology*, 14(3), 148-152.
- Seo, J. H., Lee, Y. M., Lee, J. S., Kang, H. C., & Kim, H. D. (2007). Efficacy and tolerability of the ketogenic diet according to lipid:nonlipid ratios--comparison of 3:1 with 4:1 diet. *Epilepsia*, 48(4), 801-805.
- Sharma, M., Gulati, S., & Choudhary, A. (2012). Treatment of mitochondrial disorders. *Journal of Pediatric Neurology*, 10(4), 235-245.
- Sharma, S., Jain, P., Gulati, S., Sankhyan, N., & Agarwala, A. (2015). Use of the modified atkins diet in lennox gastaut syndrome. *Journal of Child Neurology*, 30(5), 576-579.
- Sharma, S., Sankhyan, N., Gulati, S., & Agarwala, A. (2012). Use of the modified Atkins diet for the treatment of refractory childhood epilepsy: A randomized controlled trial. *Developmental Medicine and Child Neurology*, 54, 99.
- Shawki, M., El Wakeel, L., Shatla, R., El-Saeed, G., Ibrahim, S., & Badary, O. (2013). The clinical outcome of adjuvant therapy with black seed oil on intractable paediatric seizures: a pilot study. *Epileptic Disord*, 15(3), 295-301.
- Singh, M. K., Gulati, S., Aggarwal, A., Chakrabarty, B., Pandey, R. M., Tripathi, M., & Kabra, M. (2015). Efficacy of modified atkins diet versus ketogenic diet in children with refractory epilepsy aged 1 year to 18 years: A randomized controlled trial (RCT). *Epilepsia*, 56, 245.
- Sivaswamy, L., Acsadi, G., & Jiang, H. (2009). Treatment of seizures in childhood. *Therapy*, 6(1), 41-50.
- Somale, A. K., & Kondekar, S. V. (2012). Ketogenic diet as an effective alternative therapy in childhood refractory epilepsy: A systematic review. *Epilepsia*, 53, 17.
- Spilioti, M., Pavlou, E., Gogou, M., Katsanika, I., Papadopoulou-Alataki, E., Grafakou, O., Evangelou, A. (2016). Valproate effect on ketosis in children under ketogenic diet. *European Journal of Paediatric Neurology*, 0.
- Stern, J. M. (2006). Overview of treatment guidelines for epilepsy. *Current Treatment Options in Neurology*, 8(4), 280-288.
- Stern, J. M. (2009). Overview of evaluation and treatment guidelines for epilepsy. *Current Treatment Options in Neurology*, 11(4), 273-284.
- Svoronos, A., Kilbride, R. D., Mendoza, L., Szaflarski, J. P., Carpenter, A., Claassen, J., & Hirsch, L. J. (2012). Non-traditional therapies for prolonged refractory status epilepticus: A multicenter review. *Epilepsy Currents*, 12(1).
- Taub, K. S., Kessler, S. K., & Bergqvist, A. G. (2014). Risk of seizure recurrence after achieving initial seizure freedom on the ketogenic diet. *Epilepsia*, 55(4), 579-583.
- Tenney, J. R., & Glauser, T. A. (2013). The current state of absence epilepsy: Can we have your attention? *Epilepsy Currents*, 13(3), 135-140.
- Van Dam, V. S., & Korff, C. M. (2013). Dravet syndrome: An update. *Schweizer Archiv fur Neurologie und Psychiatrie*, 164(5), 153-157.
- van Rijckevorsel, K. (2008). Treatment of Lennox-Gastaut syndrome: Overview and recent findings. *Neuropsychiatric Disease and Treatment*, 4(6), 1001-1019.
- Van Sweden, B. (1988). VPA syrup in childhood epilepsy. Results of an international clinical multicentre trial. *Acta Neurologica Belgica*, 88(3), 152-162.
- Verhaert, K., Boyd, S., Neal, E. G., Chaffe, H., Schwartz, R. H., Lawson, M. S., Cross, J. H. (2009). Evolution of

- electrographic characteristics and correlation with clinical changes in children on a ketogenic diet. *European Journal of Paediatric Neurology*, 13, S73-S74.
- Vestergaard, P. (2015). Effects of antiepileptic drugs on bone health and growth potential in children with epilepsy. *Paediatric Drugs*, 17(2), 141-150.
- Yerra, S. R., & Kilpatrick, C. (2006). Pregnancy in women with epilepsy. *Medicine Today*, 7(2), 16-23.
- Yoon, J. R., Lee, E., Dong Kim, H., Soo Lee, J., & Chul Kang, H. (2014). Comparison of ketogenic diet and modified atkins diet in children with epilepsy : A randomized controlled trial. *Epilepsy Currents*, 14, 401.
- Yuen, A. W., Flugel, D., Poepel, A., Bell, G. S., Peacock, J. L., & Sander, J. W. (2012). Non-randomized open trial of eicosapentaenoic acid (EPA), an omega-3 fatty acid, in ten people with chronic epilepsy. *Epilepsy Behav*, 23(3), 370-372.
- Yuen, A. W., Sander, J. W., Fluegel, D., Patsalos, P. N., Bell, G. S., Johnson, T., & Koepp, M. J. (2005). Omega-3 fatty acid supplementation in patients with chronic epilepsy: a randomized trial. *Epilepsy Behav*, 7(2), 253-258.
- Zaccara, G., Giovannelli, F., Cincotta, M., Loiacono, G., & Verrotti, A. (2015). Adverse events of placebo-treated, drug-resistant, focal epileptic patients in randomized controlled trials: a systematic review. *Journal of Neurology*, 262(3), 501-515.
- Zare, M., Ghazvini, M. R., Yazdi, H. R., Nezhad, V. S., & Chitsaz, A. (2008). Ineffectiveness of folic acid supplementation against phenytoin-induced decrease in salivary immunoglobulin A concentration of epileptic patients. *Eur Neurol*, 59(6), 299-301.
- Zare, M., Okhovat, A. A., Esmaillzadeh, A., Mahvary, J., Najafi, M. R., & Saadatnia, M. (2015). Modified atkins diet in adult patients with refractory epilepsy: A controlled randomized clinical trial. *Epilepsia*, 56, 12.
- Zare, M., Okhovat, A. A., Najafi, M. R., Mahvary, G., & Esmaeel Zada, A. (2011). The effect of the modified Atkins diet for adults with intractable epilepsy. *Epilepsia*, 52, 39.
- Zare, M., Okhovat, A. A., Najafi, M. R., Mahvary, G., & Esmaeel Zada, A. (2011). Effects of modified Atkin's diet in adults with refractory epilepsy. *European Journal of Neurology*, 18, 458.
- Zhou, R., Jacobson, C., Weng, J., Cheng, E., Lay, J., Hung, P., Hussain, S. A. (2015). Potential efficacy of cannabidiol for treatment of refractory infantile spasms and lennox gastaut syndrome. *Epilepsy Currents*, 15, 360-361.

DBS (27)

- Badawy, R. A., Vogrin, S. J., Lai, A., & Cook, M. J. (2013). The cortical excitability profile of temporal lobe epilepsy. *Epilepsia*, 54(11), 1942-1949.
- Bauer, S., Baier, H., Baumgartner, C., Bohlmann, K., Fauser, S., Graf, W., Hamer, H. (2016). Transcutaneous Vagus Nerve Stimulation (tVNS) for Treatment of Drug-Resistant Epilepsy: a Randomized, Double-Blind Clinical Trial (cMPsE02). *Brain stimulation*, 9(3), 356-363.
- Bauer, S., Baier, H., Baumgartner, C., Bohlmann, K., Fauser, S., Graf, W., Hamer, H. M. (2016). Transcutaneous Vagus Nerve Stimulation (tVNS) for Treatment of Drug-Resistant Epilepsy: A Randomized, Double-Blind Clinical Trial (cMPsE02). *Brain Stimul*, 9(3), 356-363.
- Chang, B., & Xu, J. (2017). Deep brain stimulation for refractory temporal lobe epilepsy: a systematic review and meta-analysis with an emphasis on alleviation of seizure frequency outcome. *Childs Nerv Syst*.
- Cukiert, A., Cukiert, C., Burattini, J., Mariani, P., & Bezerra, D. (2017). Seizure outcome after hippocampal deep brain stimulation in patients with refractory temporal lobe epilepsy: a prospective, controlled, randomized, double-blind study. *Epilepsia*, 58(10), 1728-1733.
- Fisher RS, Uematsu S, Krauss GL, Cysyk BJ, McPherson R, Lesser RP, et al. Placebo-controlled pilot study of centromedian thalamic stimulation in treatment of intractable seizures. *Epilepsia* 1992; 33:841-51.
- Goldberg-Stern, H., Yaacobi, E., Phillip, M., & de Vries, L. (2014). Endocrine effects of valproic acid therapy in girls with epilepsy: a prospective study. *Eur J Paediatr Neurol*, 18(6), 759-765.
- Gonen, O. M., Gandlerman-Marton, R., Kipervasser, S., & Neufeld, M. Y. (2015). The prognosis of refractory epilepsy patients rejected from epilepsy surgery. *Acta Neurol Scand*, 131(1), 58-62.
- Hansen, N., Chaib, L., Derner, M., Hampel, K. G., Elger, C. E., Surges, R., Fell, J. (2017). Memory encoding-related anterior hippocampal potentials are modulated by deep brain stimulation of the entorhinal area. *Hippocampus*.
- Heck, C. N., King-Stephens, D., Massey, A. D., Nair, D. R., Jobst, B. C., Barkley, G. L., Morrell, M. J. (2014). Two-year seizure reduction in adults with medically intractable partial onset epilepsy treated with responsive neurostimulation: final results of the RNS System Pivotal trial. *Epilepsia*, 55(3), 432-441.
- King-Stephens, D., Mirro, E., Weber, P. B., Laxer, K. D., Ness, P. C., Salanova, V., Morrell, M. J. (2015).

- Lateralization of mesial temporal lobe epilepsy with chronic ambulatory electrocorticography. *Epilepsia*, 56(6), 959-967.
- Lane, M. A., Kahlenberg, C. A., Li, Z., Kulandaival, K., Secore, K. L., Thadani, V. M., Jobst, B. C. (2017). The implantation effect: delay in seizure occurrence with implantation of intracranial electrodes. *Acta Neurol Scand*, 135(1), 115-121.
- Lim, S. N., Lee, C. Y., Lee, S. T., Tu, P. H., Chang, B. L., Lee, C. H., Wu, T. (2016). Low and High Frequency Hippocampal Stimulation for Drug-Resistant Mesial Temporal Lobe Epilepsy. *Neuromodulation*, 19(4), 365-372.
- Lin, L. C., Lee, M. W., Wei, R. C., Mok, H. K., & Yang, R. C. (2014). Mozart K.448 listening decreased seizure recurrence and epileptiform discharges in children with first unprovoked seizures: a randomized controlled study. *BMC Complement Altern Med*, 14, 17.
- Liu, A., Bryant, A., Jefferson, A., Friedman, D., Minhas, P., Barnard, S., Schachter, S. (2016). Exploring the efficacy of a 5-day course of transcranial direct current stimulation (TDCS) on depression and memory function in patients with well-controlled temporal lobe epilepsy. *Epilepsy Behav*, 55, 11-20.
- McLachlan RS, Pigott S, Tellez-Zenteno JF, Wiebe S, Parrent A. Bilateral hippocampal stimulation for intractable temporal lobe epilepsy: impact on seizures and memory. *Epilepsia* 2010; 51:304-7.
- Otsuki, T., Kim, H. D., Luan, G., Inoue, Y., Baba, H., Oguni, H., Sugai, K. (2016). Surgical versus medical treatment for children with epileptic encephalopathy in infancy and early childhood: Results of an international multicenter cohort study in Far-East Asia (the FACE study). *Brain Dev*, 38(5), 449-460.
- Rong, P., Liu, A., Zhang, J., Wang, Y., He, W., Yang, A., Zhu, B. (2014). Transcutaneous vagus nerve stimulation for refractory epilepsy: a randomized controlled trial. *Clin Sci (Lond)*, 0.
- Salimpour, H., Kazerooni, S., Seyedabadi, M., Nabipour, I., Nemati, R., Iranpour, D., & Assadi, M. (2013). Antiepileptic treatment is associated with bone loss: difference in drug type and region of interest. *J Nucl Med Technol*, 41(3), 208-211.
- Schneider, U. C., Bohlmann, K., Vajkoczy, P., & Straub, H. B. (2015). Implantation of a new Vagus Nerve Stimulation (VNS) Therapy generator, AspireSR: considerations and recommendations during implantation and replacement surgery-comparison to a traditional system. *Acta neurochirurgica*, 157(4), 721-728.
- Seynaeve, L., Devroye, A., Dupont, P., & Paesschen, W. (2016). Randomized crossover sham-controlled clinical trial of targeted low-frequency transcranial magnetic stimulation comparing a figure-8 and a round coil to treat refractory neocortical epilepsy. *Epilepsia*, 57(1), 141-150.
- Sprengers, M., Vonck, K., Carrette, E., Marson, A. G., & Boon, P. (2017). Deep brain and cortical stimulation for epilepsy. *Cochrane Database Syst Rev*, 7, Cd008497.
- Tellez-Zenteno JF, McLachlan RS, Parrent A, Kubu CS, Wiebe S. Hippocampal electrical stimulation in mesial temporal lobe epilepsy. *Neurology* 2006; 66:1490-4.
- Troster, A., Meador, K., Irwin, C., & Fisher, R. (2017). Memory and mood outcomes after anterior thalamic stimulation for refractory partial epilepsy. *Seizure*, 45, 133-141.
- Velasco F, Velasco M, Jimenez F, Velasco AL, Brito F, Rise M, et al. Predictors in the treatment of difficult-to-control seizures by electrical stimulation of the centromedian thalamic nucleus. *Neurosurgery* 2000; 47:295-304; discussion 304-5.
- Xiaotian, X., Hengzhong, Z., Yao, X., Zhipan, Z., Daoliang, X., & Yumei, W. (2013). Effects of antiepileptic drugs on reproductive endocrine function, sexual function and sperm parameters in Chinese Han men with epilepsy. *J Clin Neurosci*, 20(11), 1492-1497.
- Yilmaz, U., Yilmaz, T. S., Akinci, G., Korkmaz, H. A., & Tekgul, H. (2014). The effect of antiepileptic drugs on thyroid function in children. *Seizure*, 23(1), 29-35.

VNS (121)

- Aalbers, M. W., Rijkers, K., Klinkenberg, S., Majoie, M., & Cornips, E. M. (2015). Vagus nerve stimulation lead removal or replacement: surgical technique, institutional experience, and literature overview. *Acta Neurochir (Wien)*, 157(11), 1917-1924.
- Abou-Khalil, B., Hirsch, E., Leroy, R. F., Shaikh, S., & Hall, S. (2011). Efficacy of retigabine as adjunctive therapy in two randomized trials in adults with drugresistant partial-onset seizures: Completers population analysis. *Epilepsy Currents*, 11(1).
- Abubakr, A., & Wambacq, I. (2008). Long-term outcome of vagus nerve stimulation therapy in patients with refractory epilepsy. *J Clin Neurosci*, 15(2), 127-129.
- Aburahma, S. K., Alzoubi, F. Q., Hammouri, H. M., & Masri, A. (2015). Vagus nerve stimulation therapy in a developing country: a long term follow up study and cost utility analysis. *Seizure*, 25, 167-172.

- Aguilar-Amat, M., Abenza, M., & Ivañez, V. (2009). Vagus nerve stimulation for drug resistant epilepsy in adults: A follow up study. *Epilepsia*, 50, 34-35.
- Ali, II, Pirzada, N. A., Kanjwal, Y., Wannamaker, B., Medhkour, A., Koltz, M. T., & Vaughn, B. V. (2004). Complete heart block with ventricular asystole during left vagus nerve stimulation for epilepsy. *Epilepsy Behav*, 5(5), 768-771.
- Anghelescu, C., Butoianu, N., Tarta-Arsene, O., Leanca, M., Dica, A., Brehar, F., Craiu, D. (2014). Could vagus nerve stimulation serve as first line of treatment for epilepsy in RETT syndrome? *Epilepsia*, 55, 145.
- Argentoni-Baldochi, M., Forster, C., Baise, C., Cukiert, C., Mello, V., Cukiert, A., Mariani, P. (2011). Vagus nerve stimulation (VNS) in children with refractory secondary generalized or multifocal epilepsy who were not candidates for cortical resection. *Epilepsy Currents*, 11(1).
- Arts, W. F., & Geerts, A. T. (2009). When to start drug treatment for childhood epilepsy: the clinical-epidemiological evidence. *Eur J Paediatr Neurol*, 13(2), 93-101.
- Bauer, S., Baier, H., Baumgartner, C., Bohlmann, K., Fauser, S., Graf, W., Hamer, H. (2016). Transcutaneous Vagus Nerve Stimulation (tVNS) for Treatment of Drug-Resistant Epilepsy: a Randomized, Double-Blind Clinical Trial (cMPsE02). *Brain stimulation*, 9(3), 356-363.
- Bauman, J. A., Ridgway, E. B., Devinsky, O., & Doyle, W. K. (2006). Subpectoral implantation of the vagus nerve stimulator. *Neurosurgery*, 58(4), ONS-322-325; discussion ONS-325-326.
- Ben-Menachem, E. (1998). Vagus nerve stimulation for treatment of seizures? *Arch Neurol*, 55(2), 231-232.
- Ben-Menachem, E. (2001). Vagus nerve stimulation, side effects, and long-term safety. *J Clin Neurophysiol*, 18(5), 415-418.
- Ben-Menachem, E. (2002). Vagus-nerve stimulation for the treatment of epilepsy. *Lancet Neurol*, 1(8), 477-482.
- Ben-Menachem, E. (2010). Quality of life and the antidepressant effects of vagus nerve stimulation (VNS) in patients with epilepsy. *Epilepsy and Behavior*, 17(4), 587.
- Ben-Menachem, E., & French, J. A. (2005). VNS Therapy versus the latest antiepileptic drug. *Epileptic Disord*, 7, S22-26.
- Bernard, E. J., Passannante, A. N., Mann, B., Lannon, S., & Vaughn, B. V. (2002). Insertion of vagal nerve stimulator using local and regional anesthesia. *Surg Neurol*, 57(2), 94-98.
- Bernstein, A. L., & Hess, T. (2007). Vagus nerve stimulation therapy for pharmacoresistant epilepsy: effect on health care utilization. *Epilepsy Behav*, 10(1), 134-137.
- Boon, P., De Herdt, V., Vonck, K., & Van Roost, D. (2007). Clinical experience with vagus nerve stimulation and deep brain stimulation in epilepsy. *Acta Neurochir Suppl*, 97, 273-280.
- Boon, P., Vonck, K., De Reuck, J., & Caemaert, J. (2001). Vagus nerve stimulation for refractory epilepsy. *Seizure*, 10(6), 456-458; quiz 458-460.
- Boon, P., Vonck, K., de Reuck, J., & Caemaert, J. (2002). Vagus nerve stimulation for refractory epilepsy. *Seizure*, 11, 448-455.
- Boon, P. A. J. M. (2015). Neurostimulation in epilepsy. *European Journal of Neurology*, 22, 867.
- Bremer, A., Eriksson, A. S., Roste, G. K., Knudtzen, B., & Nakken, K. O. (2006). [Vagal nerve stimulation in children with drug-resistant epilepsy]. *Tidsskr Nor Laegeforen*, 126(7), 896-898.
- Bunch, S., DeGiorgio, C. M., Krahl, S., Britton, J., Green, P., Lancman, M., Heck, C. N. (2007). Vagus nerve stimulation for epilepsy: is output current correlated with acute response? *Acta Neurol Scand*, 116(4), 217-220.
- Cersosimo, R. O., Bartuluchi, M., Fortini, S., Soraru, A., Pomata, H., & Caraballo, R. H. (2011). Vagus nerve stimulation: effectiveness and tolerability in 64 paediatric patients with refractory epilepsies. *Epileptic Disord*, 13(4), 382-388.
- Chadwick, D. (2001). Vagal-nerve stimulation for epilepsy. *Lancet*, 357(9270), 1726-1727.
- Chambers, A., & Bowen, J. M. (2013). Electrical stimulation for drug-resistant epilepsy: An evidence-based analysis. *Ontario Health Technology Assessment Series*, 13(18).
- Chang, E. F., & Barbaro, N. M. (2010). Nonresective epilepsy surgery. *Epilepsia*, 51, 87-89.
- Couch, J. D., Gilman, A. M., & Doyle, W. K. (2016). Long-term Expectations of Vagus Nerve Stimulation: A Look at Battery Replacement and Revision Surgery. *Neurosurgery*, 78(1), 42-46.
- Cukiert, A., Cukiert, C. M., Burattini, J. A., Lima, A. M., Forster, C. R., Baise, C., & Argentoni-Baldochi, M. (2013). Long-term outcome after callosotomy or vagus nerve stimulation in consecutive prospective cohorts of children with Lennox-Gastaut or Lennox-like syndrome and non-specific MRI findings. *Seizure*, 22(5), 396-400.
- Cukiert, A., Cukiert, C., Burattini, J., Mariani, P., & Bezerra, D. (2017). Seizure outcome after hippocampal deep brain stimulation in patients with refractory temporal lobe epilepsy: a prospective, controlled, randomized, double-blind study. *Epilepsia*, 58(10), 1728-1733.

- De Ferrari, G. M., & Schwartz, P. J. (2011). Vagus nerve stimulation: From pre-clinical to clinical application: Challenges and future directions. *Heart Failure Reviews*, 16(2), 195-203.
- DeGiorgio, C. M., Schachter, S. C., Handforth, A., Salinsky, M., Thompson, J., Uthman, B., Heck, C. (2000). Prospective long-term study of vagus nerve stimulation for the treatment of refractory seizures. *Epilepsia*, 41(9), 1195-1200.
- Dugan, P., & Devinsky, O. (2013). Epilepsy: Guidelines on vagus nerve stimulation for epilepsy. *Nat Rev Neurol*, 9(11), 611-612.
- Elliott, R. E., Carlson, C., Kalhorn, S. P., Moshel, Y. A., Weiner, H. L., Devinsky, O., & Doyle, W. K. (2009). Refractory epilepsy in tuberous sclerosis: vagus nerve stimulation with or without subsequent resective surgery. *Epilepsy Behav*, 16(3), 454-460.
- Elliott, R. E., Morsi, A., Geller, E. B., Carlson, C. C., Devinsky, O., & Doyle, W. K. (2011). Impact of failed intracranial epilepsy surgery on the effectiveness of subsequent vagus nerve stimulation. *Neurosurgery*, 69(6), 1210-1217.
- Elliott, R. E., Morsi, A., Kalhorn, S. P., Marcus, J., Sellin, J., Kang, M., Doyle, W. K. (2011). Vagus nerve stimulation in 436 consecutive patients with treatment-resistant epilepsy: long-term outcomes and predictors of response. *Epilepsy Behav*, 20(1), 57-63.
- Elliott, R. E., Morsi, A., Tanweer, O., Grobelny, B., Geller, E., Carlson, C., Doyle, W. K. (2011). Efficacy of vagus nerve stimulation over time: review of 65 consecutive patients with treatment-resistant epilepsy treated with VNS > 10 years. *Epilepsy Behav*, 20(3), 478-483.
- Elliott, R. E., Rodgers, S. D., Bassani, L., Morsi, A., Geller, E. B., Carlson, C., Doyle, W. K. (2011). Vagus nerve stimulation for children with treatment-resistant epilepsy: a consecutive series of 141 cases. *J Neurosurg Pediatr*, 7(5), 491-500.
- Englot, D. J., Rolston, J. D., Wang, D. D., Hassnain, K. H., Gordon, C. M., & Chang, E. F. (2012). Efficacy of vagus nerve stimulation in posttraumatic versus nontraumatic epilepsy. *J Neurosurg*, 117(5), 970-977.
- Ferlisi, M., & Shorvon, S. (2012). The outcome of therapies in refractory and super-refractory convulsive status epilepticus and recommendations for therapy. *Brain*, 135, 2314-2328.
- FineSmith, R. B., Zampella, E., & Devinsky, O. (1999). Vagal nerve stimulator: a new approach to medically refractory epilepsy. *N J Med*, 96(6), 37-40.
- Frost, M., Gates, J., Helmers, S. L., Wheless, J. W., Levisohn, P., Tardo, C., & Conry, J. A. (2001). Vagus nerve stimulation in children with refractory seizures associated with Lennox-Gastaut syndrome. *Epilepsia*, 42(9), 1148-1152.
- Funnel, M., & Ranta, A. (2012). Vagal nerve stimulation in New Zealand: improvement of seizure control following the implantation of a vagal nerve stimulator. *N Z Med J*, 125(1361), 109-111.
- Gatzonis, S. D., Georgaculias, N., Singounas, E., Jenkins, A., Stamboulis, E., & Siafakas, A. (1999). Elimination of oxcarbazepine-induced oculogyric crisis following vagus nerve stimulation. *Neurology*, 52(9), 1918-1919.
- Ghaemi, K., Elsharkawy, A. E., Schulz, R., Hoppe, M., Polster, T., Pannek, H., & Ebner, A. (2010). Vagus nerve stimulation: outcome and predictors of seizure freedom in long-term follow-up. *Seizure*, 19(5), 264-268.
- Guberman, A. (2004). Vagus nerve stimulation in the treatment of epilepsy. *Cmaj*, 171(10), 1165-1166.
- Hallbook, T., Lundgren, J., Stjernqvist, K., Blennow, G., Stromblad, L. G., & Rosen, I. (2005). Vagus nerve stimulation in 15 children with therapy resistant epilepsy; its impact on cognition, quality of life, behaviour and mood. *Seizure*, 14(7), 504-513.
- Harvey, A. S. (2003). Topiramate: Potential trade-offs between efficacy and tolerability in the treatment of epilepsy. *Journal of Paediatrics and Child Health*, 39(6), 414-415.
- Hatton, K. W., McLarney, J. T., Pittman, T., & Fahy, B. G. (2006). Vagal nerve stimulation: Overview and implications for anesthesiologists. *Anesthesia and Analgesia*, 103(5), 1241-1249.
- He, W., Wang, X. Y., Zhou, L., Li, Z. M., Jing, X. H., Lv, Z. L., Zhu, B. (2015). Transcutaneous auricular vagus nerve stimulation for pediatric epilepsy: study protocol for a randomized controlled trial. *Trials*, 16, 371.
- Healy, S., Lang, J., Te Water Naude, J., Gibbon, F., & Leach, P. (2013). Vagal nerve stimulation in children under 12 years old with medically intractable epilepsy. *Childs Nerv Syst*, 29(11), 2095-2099.
- Heberle, C., Berquin, P., Larnicol, N., & Wallois, F. (2002). Vagus nerve stimulation in a case of epilepsy with CSWSS: respiratory side effects during sleep. *Epilepsia*, 43(10), 1268-1270.
- Heck, C. N. (2011). Long term follow-up of the RNS(TM) system in adults with medically intractable partial onset seizures. *Epilepsy Currents*, 11(1).
- Helmers, S. L., Duh, M. S., Guerin, A., Sarda, S. P., Samuelson, T. M., Bunker, M. T., Faught, E. (2012). Clinical outcomes, quality of life, and costs associated with implantation of vagus nerve stimulation therapy in pediatric patients with drug-resistant epilepsy. *Eur J Paediatr Neurol*, 16(5), 449-458.
- Helmers, S. L., Griesemer, D. A., Dean, J. C., Sanchez, J. D., Labar, D., Murphy, J. V., Morris, G. L., 3rd. (2003).

- Observations on the use of vagus nerve stimulation earlier in the course of pharmacoresistant epilepsy: patients with seizures for six years or less. *Neurologist*, 9(3), 160-164.
- Helmers, S. L., Wheless, J. W., Frost, M., Gates, J., Levisohn, P., Tardo, C., Madsen, J. R. (2001). Vagus nerve stimulation therapy in pediatric patients with refractory epilepsy: retrospective study. *J Child Neurol*, 16(11), 843-848.
- Herman, S. T., & Pedley, T. A. (1998). New options for the treatment of epilepsy. *Jama*, 280(8), 693-694.
- Howland, R. H. (2006). What is vagus nerve stimulation? *J Psychosoc Nurs Ment Health Serv*, 44(8), 11-14.
- Kabir, S. M., Rajaraman, C., Rittey, C., Zaki, H. S., Kemeny, A. A., & McMullan, J. (2009). Vagus nerve stimulation in children with intractable epilepsy: indications, complications and outcome. *Childs Nerv Syst*, 25(9), 1097-1100.
- Kalanithi PS, Arrigo RT, Tran P, Gephart MH, Shuer L, Fisher R, et al. Rehospitalization and emergency department use rates before and after vagus nerve stimulation for epilepsy: use of state databases to provide longitudinal data across multiple clinical settings. *Neuromodulation* 2014; 17:60-4; discussion 64-5.
- Kahlow, H., & Olivecrona, M. (2013). Complications of vagal nerve stimulation for drug-resistant epilepsy: a single center longitudinal study of 143 patients. *Seizure*, 22(10), 827-33.
- Kapur, R., Meador, K. J., Loring, D. W., Kanner, A. M., & Morrell, M. J. (2014). Quality of life (QOL) and mood in adults with medically intractable partial onset seizures treated with responsive cortical stimulation. *Annals of Neurology*, 76, S31.
- Kellinghaus, C., & Loddenkemper, T. (2006). Double-blind, randomized controlled study of bilateral cerebellar stimulation [1]. *Epilepsia*, 47(7), 1247.
- Kelts, G., O'Connor, P. D., Hussey, R. W., & Maturo, S. (2015). An electrical cause of stridor: pediatric vagal nerve stimulators. *Int J Pediatr Otorhinolaryngol*, 79(2), 251-253.
- Khurana, D. S., Reumann, M., Hobdell, E. F., Neff, S., Valencia, I., Legido, A., & Kothare, S. V. (2007). Vagus nerve stimulation in children with refractory epilepsy: unusual complications and relationship to sleep-disordered breathing. *Childs Nerv Syst*, 23(11), 1309-1312.
- Kossoff, E. H., Pyzik, P. L., Rubenstein, J. E., Bergqvist, A. G., Buchhalter, J. R., Donner, E. J., Wheless, J. W. (2007). Combined ketogenic diet and vagus nerve stimulation: rational polytherapy? *Epilepsia*, 48(1), 77-81.
- Kostov, K., Kostov, H., & Tauboll, E. (2009). Long-term vagus nerve stimulation in the treatment of Lennox-Gastaut syndrome. *Epilepsy Behav*, 16(2), 321-324.
- Koutroumanidis, M., Hennessy, M. J., Binnie, C. D., & Polkey, C. E. (2000). Aggravation of partial epilepsy and emergence of new seizure type during treatment with VNS. *Neurology*, 55(6), 892-893.
- Labar, D. (2004). Vagus nerve stimulation for epilepsy. *Clinical Neuroscience Research*, 4(1), 81-87.
- Labar, D., & Ponticello, L. (2003). Persistent antiepileptic effects after vagus nerve stimulation ends? *Neurology*, 61(12), 1818.
- Labar, D. R. (2002). Antiepileptic drug use during the first 12 months of vagus nerve stimulation therapy: a registry study. *Neurology*, 59(6), S38-43.
- Marson, A. G., Maguire, M., & Ramaratnam, S. (2009). Epilepsy. *BMJ Clin Evid*, 2009.
- McLachlan, R. S. (1997). Vagus nerve stimulation for intractable epilepsy: a review. *J Clin Neurophysiol*, 14(5), 358-368.
- Menascu, S., Kremer, U., Schiller, Y., Blatt, I., Watemberg, N., Boxer, M., Ben-Zeev, B. (2013). The Israeli retrospective multicenter open-label study evaluating vagus nerve stimulation efficacy in children and adults. *Israel Medical Association Journal*, 15(11), 673-677.
- Mesa, T., Rios, L., Chicharro, A., & Muñoz, M. (2011). Induction of psychogenic nonepileptic seizures: Comparison between simultaneous hyperventilation and photic stimulus versus IV saline placebo. *Epilepsia*, 52, 104.
- Morris Iii, G. L. (2003). A retrospective analysis of the effects of magnet-activated stimulation in conjunction with vagus nerve stimulation therapy. *Epilepsy and Behavior*, 4(6), 740-745.
- Murphy, J. V., & Patil, A. A. (2005). Improving the lives of patients with medically refractory epilepsy by electrical stimulation of the nervous system. *Expert Rev Med Devices*, 2(2), 175-189.
- Nagarajan, L., Walsh, P., Gregory, P., & Lee, M. (2002). VNS therapy in clinical practice in children with refractory epilepsy. *Acta Neurol Scand*, 105(1), 13-17.
- Nakken, K. O., Henriksen, O., Roste, G. K., & Lossius, R. (2003). Vagal nerve stimulation--the Norwegian experience. *Seizure*, 12(1), 37-41.
- Neufeld, M. Y., & Korczyn, A. D. (1996). New drugs and vagal stimulation for treatment of epilepsy--the Israeli experience. *Neurol Neurochir Pol*, 30, 113-120.
- Orosz, I., Hoffmann, J., Kerling, F., Polster, T., Hoppe, M., König, I., Sperner, J. (2009). Is transcranial direct

- current stimulation (tDCS) predictive for vagus nerve stimulation (VNS) therapy outcome? *Epilepsia*, 50, 65.
- Orosz, I., Hoffmann, J., Panzer, A., König, I., & Sperner, J. (2009). Is prediction of Vagus Nerve Stimulation (VNS) therapy outcome possible? *Epilepsia*, 50, 163-164.
- Orosz, I., Hoffmann, J., Panzer, A., König, I., & Sperner, J. (2009). Prediction of vagus nerve stimulation outcome in patients with pharmacoresistant epilepsy. *Epilepsia*, 50, 25-26.
- Parhizgar, F., Nugent, K., & Raj, R. (2011). Obstructive sleep apnea and respiratory complications associated with vagus nerve stimulators. *J Clin Sleep Med*, 7(4), 401-407.
- Park, Y. D. (2003). The effects of vagus nerve stimulation therapy on patients with intractable seizures and either Landau-Kleffner syndrome or autism. *Epilepsy Behav*, 4(3), 286-290.
- Pearl, P. L., Conry, J. A., Yaun, A., Taylor, J. L., Heffron, A. M., Sigman, M., Gaillard, W. D. (2008). Misidentification of vagus nerve stimulator for intravenous access and other major adverse events. *Pediatr Neurol*, 38(4), 248-251.
- Privitera, M. D., Welty, T. E., Ficker, D. M., & Welge, J. (2002). Vagus nerve stimulation for partial seizures. *Cochrane Database Syst Rev*, 0(1), Cd002896.
- Privitera, M. D., Welty, T. T. E., Ficker, D. D. M., & Welge, J. (2009). Vagus nerve stimulation for partial seizures. *Cochrane Database of Systematic Reviews*, 0(4).
- Ramsay, E., Kuzniecky, R., Rosenfeld, W., & Mirza, W. (1993). Acute side effects of vagus nerve stimulation (VNS) in 113 refractory epilepsy patients. *Epilepsia*, 34, 193.
- Rao, M. B. (2013). Surgery for drug-resistant focal epilepsy. *J Assoc Physicians India*, 61(8), 55-57.
- Robinson, R. (2011). Children who have experienced >5 aeds are more likely to respond to vagal nerve stimulation (VNS) than trials of further AEDS. *Epilepsia*, 52, 74.
- Robinson, R. O. (2011). Vagal nerve stimulation is more likely to reduce seizure frequency and severity than trials of further anti-epileptic drugs in children who have already had more than five anti-epileptic drugs. *Developmental Medicine and Child Neurology*, 53, 15.
- Rolston, J. D., Englot, D. J., Wang, D. D., Shih, T., & Chang, E. F. (2012). Comparison of seizure control outcomes and the safety of vagus nerve, thalamic deep brain, and responsive neurostimulation: evidence from randomized controlled trials. *Neurosurg Focus*, 32(3), E14.
- Rydenhag, B., Silander, H., & Ben-Menachem, E. (2013). Preliminary experience with a new system for vagus nerve stimulation. *Epilepsy Currents*, 13, 61.
- Ryvlin, P. (2003). Beyond pharmacotherapy: surgical management. *Epilepsia*, 44, 23-28.
- Ryvlin, P., Gilliam, F., Nguyen, D., Colicchio, G., Pizzanelli, C., Marroso, F., Perucca, E. (2011). Randomized controlled trial shows quality-of-life benefits with VNS added-on to best medical therapy versus best medical therapy alone in adults with drug-resistant epilepsy. *Epilepsia*, 52, 83.
- Ryvlin, P., Gilliam, F. G., Nguyen, D. K., Colicchio, G., Iudice, A., Tinuper, P., Perucca, E. (2015). Erratum: The long-term effect of vagus nerve stimulation on quality of life in patients with pharmacoresistant focal epilepsy: The PuLSE (Open Prospective Randomized Long-term Effectiveness) trial (*Epilepsia* (2014) 55(893-900)). *Epilepsia*, 56(6), 983.
- Ryzi, M., Brazdil, M., Novak, Z., Chrastina, J., Oslejskova, H., Rektor, I., & Kuba, R. (2013). Long-term vagus nerve stimulation in children with focal epilepsy. *Acta Neurol Scand*, 127(5), 316-322.
- Sakas, D. E., Korfiatis, S., Nicholson, C. L., Panourias, I. G., Georgakoulias, N., Gatzonis, S., & Jenkins, A. (2007). Vagus nerve stimulation for intractable epilepsy: outcome in two series combining 90 patients. *Acta Neurochir Suppl*, 97, 287-291.
- Sanossian, N., & Haut, S. (2002). Chronic diarrhea associated with vagal nerve stimulation. *Neurology*, 58(2), 330.
- Schachter, S. C. (2002). Vagus nerve stimulation therapy summary: five years after FDA approval. *Neurology*, 59(6), S15-20.
- Schachter, S. C. (2006). Therapeutic effects of vagus nerve stimulation in epilepsy and implications for sudden unexpected death in epilepsy. *Clin Auton Res*, 16(1), 29-32.
- Schallert, G., Foster, J., Lindquist, N., & Murphy, J. V. (1998). Chronic stimulation of the left vagal nerve in children: effect on swallowing. *Epilepsia*, 39(10), 1113-1114.
- Shankar, R., Olotu, V. O., Cole, N., Sullivan, H., & Jory, C. (2013). Case report: vagal nerve stimulation and late onset asystole. *Seizure*, 22(4), 312-314.
- Sheth, R. D., & Stafstrom, C. E. (2002). Intractable pediatric epilepsy: Vagal nerve stimulation and the ketogenic diet. *Neurologic Clinics*, 20(4), 1183-1194.
- Shields, W. D. (2004). Management of epilepsy in mentally retarded children using the newer antiepileptic drugs, vagus nerve stimulation, and surgery. *J Child Neurol*, 19, S58-64.
- Shiozawa, P., da Silva, M. E., de Carvalho, T. C., Cordeiro, Q., Brunoni, A. R., & Fregni, F. (2014). Transcutaneous

- vagus and trigeminal nerve stimulation for neuropsychiatric disorders: A systematic review. *Arquivos de Neuro-Psiquiatria*, 72(7), 542-547.
- Shorvon, S., & Ferlisi, M. (2011). The treatment of super-refractory status epilepticus: a critical review of available therapies and a clinical treatment protocol. *Brain*, 134, 2802-2818.
- Singh, A., Dalal, K., & Tripathi, M. (2007). Indirect vagus nerve stimulation and reflexology for management of epilepsy. *Epilepsia*, 48, 84.
- Smyth, M. D., Tubbs, R. S., Bebin, E. M., Grabb, P. A., & Blount, J. P. (2003). Complications of chronic vagus nerve stimulation for epilepsy in children. *J Neurosurg*, 99(3), 500-503.
- Spanaki, M. V., Allen, L. S., Mueller, W. M., & Morris, G. L., 3rd. (2004). Vagus nerve stimulation therapy: 5-year or greater outcome at a university-based epilepsy center. *Seizure*, 13(8), 587-590.
- Spuck, S., Tronnier, V., Orosz, I., Schonweiler, R., Sepehrnia, A., Nowak, G., & Sperner, J. (2010). Operative and technical complications of vagus nerve stimulator implantation. *Neurosurgery*, 67(2), 489-494.
- Terra, V. C., Furlanetti, L. L., Nunes, A. A., Thomé, U., Nisyama, M. A., Sakamoto, A. C., & Machado, H. R. (2014). Vagus nerve stimulation in pediatric patients: Is it really worthwhile? *Epilepsy and Behavior*, 31, 329-333.
- Thompson, E. M., Wozniak, S. E., Roberts, C. M., Kao, A., Anderson, V. C., & Selden, N. R. (2012). Vagus nerve stimulation for partial and generalized epilepsy from infancy to adolescence. *J Neurosurg Pediatr*, 10(3), 200-205.
- Uthman, B. M., Reichl, A. M., Dean, J. C., Eisenschenk, S., Gilmore, R., Reid, S., Wilder, B. J. (2004). Effectiveness of vagus nerve stimulation in epilepsy patients: a 12-year observation. *Neurology*, 63(6), 1124-1126.
- Wheless, J. W., & Maggio, V. (2002). Vagus nerve stimulation therapy in patients younger than 18 years. *Neurology*, 59(6), S21-25.
- Vonck, K., Dedeurwaerdere, S., De Groote, L., Thadani, V., Claeys, P., Gossiaux, F., Boon, P. (2005). Generator replacement in epilepsy patients treated with vagus nerve stimulation. *Seizure*, 14(2), 89-99.
- Wu, C., & Sharan, A. D. (2013). Neurostimulation for the treatment of epilepsy: a review of current surgical interventions. *Neuromodulation*, 16(1), 10-24; discussion 24.
- Zalvan, C., Sulica, L., Wolf, S., Cohen, J., Gonzalez-Yanes, O., & Blitzer, A. (2003). Laryngopharyngeal dysfunction from the implant vagal nerve stimulator. *Laryngoscope*, 113(2), 221-225.
- Zumsteg, D., Jenny, D., & Wieser, H. G. (2000). Vocal cord adduction during vagus nerve stimulation for treatment of epilepsy. *Neurology*, 54(6), 1388-1389.

Hälsoekonomi


- Aburahma SK, Alzoubi FQ, Hammouri HM, Masri A. Vagus nerve stimulation therapy in a developing country: a long term follow up study and cost utility analysis. *Seizure* 2015;25:167-72.
- Al-Qudah AA, Abu-Sheik S, Tamimi AF. Diagnostic value of short duration outpatient video electroencephalographic monitoring. *Pediatr Neurol* 1999;21:622-5.
- Beghi E, Frigeni B, Beghi M, De Compadri P, Garattini L. A review of the costs of managing childhood epilepsy. *PharmacoEconomics* 2005;23:27-45.
- Ben-Menachem E, French JA. VNS Therapy versus the latest antiepileptic drug. *Epileptic Disord* 2005;7 Suppl 1:S22-6.
- Ben-Menachem E, Hellstrom K, Verstappen D. Analysis of direct hospital costs before and 18 months after treatment with vagus nerve stimulation therapy in 43 patients. *Neurology* 2002;59:S44-7.
- Beran RG. Economic analysis of epilepsy care. *Epilepsia* 1999;40 Suppl 8:19-24.
- Boon P, D'Have M, Van Walleghem P, Michielsen G, Vonck K, Caemaert J, et al. Direct medical costs of refractory epilepsy incurred by three different treatment modalities: a prospective assessment. *Epilepsia* 2002;43:96-102.
- Boon P, Vonck K, D'Have M, O'Connor S, Vandekerckhove T, De Reuck J. Cost-benefit of vagus nerve stimulation for refractory epilepsy. *Acta Neurol Belg* 1999;99:275-80.
- Boon P, Vonck K, Reuck JD, Caemaert J. Vagus nerve stimulation for refractory epilepsy. *Seizure* 2001;10:448-455.
- Boon P, Vonck K, Vandekerckhove T, De Reuck J, Calliauw L. Vagus nerve stimulation for refractory epilepsy. *Neurosurgery Quarterly* 2001;11:66-72.
- Boon P, Vonck K, Vandekerckhove T, D'Have M, Nieuwenhuis L, Michielsen G, et al. Vagus nerve stimulation for medically refractory epilepsy; efficacy and cost-benefit analysis. *Acta Neurochir (Wien)* 1999;141:447-52; discussion 453.

- Bronen RA, Fulbright RK, Spencer SS, Spencer DD, Kim JH, Lange RC. Economic impact of replacing CT with MR imaging for refractory epilepsy. *Magn Reson Imaging* 1997;15:857-62.
- Camp C, Smithson WH, Bunker M, Burke T, Hughes D. Impact of vagus nerve stimulation on secondary care burden in children and adults with epilepsy: Review of routinely collected hospital data in England. *Epilepsy Behav* 2015;52:68-73.
- Chambers A, Bowen JM. Electrical stimulation for drug-resistant epilepsy: An evidence-based analysis. Ontario Health Technology Assessment Series 2013;13.
- Chayasirisobhon S, Griggs L, Westmoreland S, Kim CS. The usefulness of one to two hour video EEG monitoring in patients with refractory seizures. *Clinical EEG Electroencephalography* 1993;24:78-84.
- Chen J, Lei D. Surgery: a cost-effective option for drug-resistant epilepsy in China. *World Neurosurg* 2014;82:e375-6.
- Colombo F. Developing areas and their cost-benefit correlations: radiosurgery. *Childs Nerv Syst* 1995;11:29-32.
- Corabian P, Leggett P. Vagus nerve stimulation for refractory epilepsy (Structured abstract). In: Health Technology Assessment Database. Alberta Heritage Foundation for Medical Research (AHFMR); 2001. p 42.
- Dash D, Hernandez-Ronquillo L, Moien-Afshari F, Tellez-Zenteno JF. Ambulatory EEG: a cost-effective alternative to inpatient video-EEG in adult patients. *Epileptic Disord* 2012;14:290-7.
- de Kinderen RJ, Lambrechts DA, Wijnen BF, Postulart D, Aldenkamp AP, Majoe MH, et al. An economic evaluation of the ketogenic diet versus care as usual in children and adolescents with intractable epilepsy: An interim analysis. *Epilepsia* 2016;57:41-50.
- De Oliveira THF, Francisco AN, Demartini Junior Z, Stebel SL. The role of vagus nerve stimulation in refractory epilepsy. *Arquivos de Neuro-Psiquiatria* 2017;75:657-666.
- DellaBadia J, Jr., Bell WL, Keyes JW, Jr., Mathews VP, Glazier SS. Assessment and cost comparison of sleep-deprived EEG, MRI and PET in the prediction of surgical treatment for epilepsy. *Seizure* 2002;11:303-9.
- Elisevich K, Kasunic K, Hathaway SJ, Smith BJ, Stewart RW, Barkley GL. Clinical care pathways in a surgical epilepsy program: A comparison of costs by diagnostic related groups for program and nonprogram admissions. *Journal of Epilepsy* 1996;9:282-290.
- Epilepsy surgery: an evidence summary (Structured abstract). In: Health Technology Assessment Database. Medical Advisory Secretariat, Ontario Ministry of Health and Long-Term Care (MAS); 2012.
- Forbes R. Cost-utility of vagus nerve stimulation (VNS) therapy for medically refractory epilepsy--an update. *Seizure* 2008;17:387-88.
- Forbes RB, Macdonald S, Eljamal S, Roberts RC. Cost-utility analysis of vagus nerve stimulators for adults with medically refractory epilepsy. *Seizure* 2003;12:249-256.
- Gilbert DL, Pyzik PL, Vining EP, Freeman JM. Medication cost reduction in children on the ketogenic diet: data from a prospective study. *J Child Neurol* 1999;14:469-71.
- Guerreiro CA, Montenegro MA, Kobayashi E, Noronha AL, Guerreiro MM, Cendes F. Daytime outpatient versus inpatient video-EEG monitoring for presurgical evaluation in temporal lobe epilepsy. *J Clin Neurophysiol* 2002;19:204-8.
- Helmers SL, Duh MS, Guerin A, Sarda SP, Samuelson TM, Bunker MT, et al. Clinical and economic impact of vagus nerve stimulation therapy in patients with drug-resistant epilepsy. *Epilepsy Behav* 2011;22:370-5.
- Helmers SL, Duh MS, Guerin A, Sarda SP, Samuelson TM, Bunker MT, et al. Clinical outcomes, quality of life, and costs associated with implantation of vagus nerve stimulation therapy in pediatric patients with drug-resistant epilepsy. *Eur J Paediatr Neurol* 2012;16:449-58.
- Jarrar RG, Buchhalter JR. Therapeutics in pediatric epilepsy, Part 1: The new antiepileptic drugs and the ketogenic diet. *Mayo Clin Proc* 2003;78:359-70.
- Jennen P, Sabers A, Christensen J, Ibsen R, Kjellberg J. Socioeconomic evaluation of vagus stimulation: A controlled national study. *Seizure* 2016;42:15-19.
- Kalanithi PS, Arrigo RT, Tran P, Gephart MH, Shuer L, Fisher R, et al. Rehospitalization and emergency department use rates before and after vagus nerve stimulation for epilepsy: use of state databases to provide longitudinal data across multiple clinical settings. *Neuromodulation* 2014;17:60-4; discussion 64-5.
- Kayyali HR, Luniova A, Abdelmoity A. Ketogenic Diet Decreases Emergency Room Visits and Hospitalizations Related to Epilepsy. *Epilepsy Res Treat* 2016;2016:5873208.
- Keene D, Ventureyra ECG. Epilepsy surgery for 5- to 18-year old patients with medically refractory epilepsy - Is it cost efficient? *Child's Nervous System* 1999;15:52-55.
- King JT, Jr., Sperling MR, Justice AC, O'Connor MJ. A cost-effectiveness analysis of anterior temporal lobectomy for intractable temporal lobe epilepsy. *J Neurosurg* 1997;87:20-8.


- Kitwitee P, Unnwongse K, Srikiwilaikul T, Yadee T, Limwattananon C. Corrigendum to 'Cost-Utility of Video-Electroencephalography Monitoring Followed by Surgery in Adults with Drug-Resistant Focal Epilepsy in Thailand' [World Neurosurgery 98 (2017) 750-760]. World Neurosurgery 2017.
- Kitwitee P, Unnwongse K, Srikiwilaikul T, Yadee T, Limwattananon C. Cost-Utility of Video-Electroencephalography Monitoring Followed by Surgery in Adults with Drug-Resistant Focal Epilepsy in Thailand. World Neurosurg 2017;98:750-760.e3.
- Kuba R, Novak Z, Chrastina J, Pazourkova M, Hermanova M, Oslejskova H, et al. Comparing the effects of cortical resection and vagus nerve stimulation in patients with nonlesional extratemporal epilepsy. Epilepsy Behav 2013;28:474-80.
- Kuzniecky R, Devinsky O. Surgery Insight: Surgical management of epilepsy. Nature Clinical Practice Neurology 2007;3:673-681.
- Lachaine J, Laberge A, Bibeau J, Lambert-Obry V. Systematic review of economic evaluations of treatments in epilepsy. Value in Health 2015;18:A756-A757.
- Langfitt JT, Bronstein K. Use of surgical procedures in the treatment of epilepsy: Interventions and outcomes. Disease Management and Health Outcomes 1999;5:23-40.
- Langfitt JT, Holloway RG, McDermott MP, Messing S, Sarosky K, Berg AT, et al. Health care costs decline after successful epilepsy surgery. Neurology 2007;68:1290-8.
- Langfitt JT. Cost-effectiveness of anterotemporal lobectomy in medically intractable complex partial epilepsy. Epilepsia 1997;38:154-63.
- Lansingh VC, Carter MJ, Martens M. Global Cost-effectiveness of Cataract Surgery. Ophthalmology 2007;114:1670-1678.
- MacCracken KA, Scalisi JC. Development and evaluation of a ketogenic diet program. J Am Diet Assoc 1999;99:1554-8.
- Majoie HJ, Berfelo MW, Aldenkamp AP, Evers SM, Kessels AG, Renier WO. Vagus nerve stimulation in children with therapy-resistant epilepsy diagnosed as Lennox-Gastaut syndrome: clinical results, neuropsychological effects, and cost-effectiveness. J Clin Neurophysiol 2001;18:419-28.
- Malmgren K, Hedstrom A, Granqvist R, Malmgren H, Ben-Menachem E. Cost analysis of epilepsy surgery and of vigabatrin treatment in patients with refractory partial epilepsy. Epilepsy Res 1996;25:199-207.
- Mandel A, Ballew M, Pina-Garza JE, Stalmasek V, Clemens LH. Medical costs are reduced when children with intractable epilepsy are successfully treated with the ketogenic diet. J Am Diet Assoc 2002;102:396-8.
- Mansouri A, Aldakkan A, Kosicka MJ, Tarride JE, Valiante TA. Bridging the gap between evidence and practice for adults with medically refractory temporal lobe epilepsy: Is a change in funding policy needed to stimulate a shift in practice? Epilepsy Research and Treatment 2015;2015.
- Marras CE, Canevini MP, Colicchio G, Guerrini R, Rubboli G, Scerrati M, et al. Health Technology Assessment report on the presurgical evaluation and surgical treatment of drug-resistant epilepsy. Epilepsia 2013;54:49-58.
- Moien-Afshari F, Griebel R, Sadanand V, Vrbancic M, Hernandez-Ronquillo L, Lowry N, et al. Safety and yield of early cessation of AEDs in video-EEG telemetry and outcomes. Can J Neurol Sci 2009;36:587-92.
- Murtagh J, Foerster V, Warburton RN, Lentle BC, Wood RJ, Mensinkai S, et al. Clinical and cost effectiveness of CT and MRI for selected clinical disorders: results of two systematic reviews (Structured abstract). In: Health Technology Assessment Database. Canadian Agency for Drugs and Technologies in Health (CADTH); 2006. p 15.
- Murtagh J, Warburton RN, Foerster V, Lentle BC, Wood RJ, Mensinkai S, et al. CT and MRI for selected clinical disorders: a systematic review of economic evaluations (Structured abstract). In: Health Technology Assessment Database. Canadian Agency for Drugs and Technologies in Health (CADTH); 2006. p 96.
- Oldham MS, Horn PS, Tsevat J, Standridge S. Costs and Clinical Outcomes of Epilepsy Surgery in Children With Drug-Resistant Epilepsy. Pediatr Neurol 2015;53:216-20.
- Pichon RA, Augustovski F, Alcaraz A, Bardach A, Ferrante D, Garcia MS, et al. Vagus Nerve Stimulation for refractory epilepsy (Structured abstract). In: Health Technology Assessment Database. Institute for Clinical Effectiveness and Health Policy (IECS); 2006.
- Pichon RA, Augustovski F, Cernadas C, Ferrante D, Regueiro A, Garcia MS. Epilepsy surgery (Structured abstract). In: Health Technology Assessment Database; 2003.
- Pichon RA, Augustovski F, Garcia MS, Glujsovsky D, Lopez A, Rey-Ares L, et al. Vagus nerve stimulation for refractory epilepsy (Structured abstract). In: Health Technology Assessment Database. Institute for Clinical Effectiveness and Health Policy (IECS); 2011.

- Picot MC, Jaussent A, Neveu D, Kahane P, Crespel A, Gelisse P, et al. Cost-effectiveness analysis of epilepsy surgery in a controlled cohort of adult patients with intractable partial epilepsy: A 5-year follow-up study. *Epilepsia* 2016;57:1669-1679.
- Pillas D, Selai C. Economic aspects of epilepsy and antiepileptic treatment: A review of the literature. *Expert Review of Pharmacoeconomics and Outcomes Research* 2005;5:327-338.
- Rao MB, Radhakrishnan K. Is epilepsy surgery possible in countries with limited resources? *Epilepsia* 2000;41 Suppl 4:S31-4.
- Scherrmann J, Hoppe C, Kral T, Schramm J, Elger CE. Vagus nerve stimulation: clinical experience in a large patient series. *J Clin Neurophysiol* 2001;18:408-14.
- Schiltz NK, Kaiboriboon K, Koroukian SM, Singer ME, Love TE. Long-term reduction of health care costs and utilization after epilepsy surgery. *Epilepsia* 2016;57:316-24.
- Scott Perry M, Duchowny M. Surgical versus medical treatment for refractory epilepsy: Outcomes beyond seizure control. *Epilepsia* 2013;54:2060-2070.
- Si Y, Liu L, Fang JJ, Mu J, Hu J, Zhao LL, et al. Evaluation of the efficiency of inpatient 24-hour VEEG combined with MRI in consecutive patients with newly diagnosed epilepsies. *Epilepsy Behav* 2011;20:633-7.
- Silfvenius H, Dahlgren H, Jonsson E, Magnusson S, Marke L, Norlund A. Surgery for epilepsy (Structured abstract). In: Health Technology Assessment Database; 1991. p 95.
- Silfvenius H. Cost and cost-effectiveness of epilepsy surgery. *Epilepsia* 1999;40:32-39.
- Silfvenius H. Cost-benefit of epilepsy surgery. *Acta Neurol Belg* 1999;99:266-74.
- Surgery often most cost-effective option for intractable epilepsy. *Drugs and Therapy Perspectives* 1998;12:13-16.
- Tureczek IE, Fandino-Franky J, Wieser HG. Comparison of the epilepsy surgery programs in Cartagena, Colombia, and Zurich, Switzerland. *Epilepsia* 2000;41 Suppl 4:S35-40.
- van Hout B. Determining costs and cost-effectiveness of epilepsy. *Am J Manag Care* 1998;4:S501-9.
- Whiting P, Gupta R, Burch J, Mujica Mota RE, Wright K, Marson A, et al. A systematic review of the effectiveness and cost-effectiveness of neuroimaging assessments used to visualise the seizure focus in people with refractory epilepsy being considered for surgery. *Health Technology Assessment* 2006;10:iii-128.
- Whiting S, Donner E, RamachandranNair R, Grabowski J, Jetté N, Duque DR. Decreased health care utilization and health care costs in the inpatient and emergency department setting following initiation of ketogenic diet in pediatric patients: The experience in Ontario, Canada. *Epilepsy Research* 2017;131:51-57.
- Widjaja E, Li B, Medina LS. Diagnostic evaluation in patients with intractable epilepsy and normal findings on MRI: a decision analysis and cost-effectiveness study. *AJNR Am J Neuroradiol* 2013;34:1004-9, s1-2.
- Widjaja E, Li B, Schinkel CD, Puchalski Ritchie L, Weaver J, Snead OC, et al. Cost-effectiveness of pediatric epilepsy surgery compared to medical treatment in children with intractable epilepsy. *Epilepsy Res* 2011;94:61-8.
- Wiebe S, Gafni A, Blume WT, Girvin JP. An economic evaluation of surgery for temporal lobe epilepsy. *Journal of Epilepsy* 1995;8:227-235.

Flödesträd över litteraturgenomgången



Figur 1 Sökning och urval av litteratur. Exakt antal granskade artiklar i fulltext samt exakt antal exkluderade artiklar kan inte anges eftersom flera granskade och exkluderade artiklar varit aktuell för flera olika åtgärder.



Figur 2. Flödesschema över litteratursökning hälsoekonomi