

Behandling av rektusdiastas hos kvinnor/Treatment of diastasis recti in women, rapport 346 (2022)

Bilaga 2 Exkluderade artiklar och artiklar med hög risk för bias

Sammanställning av studier som exkluderas efter relevansgranskning i fulltext

Referens	Exklusionsorsak
Efficacy and Safety of Microfocused Ultrasound With Visualization in Abdominal Skin Laxity: A Randomized, Comparative Study. John Wiley & Sons, Inc.; 2020. p. 831-6.	Population
A.T K, Sahoo MR. Laparoscopic plication and mesh repair for diastasis recti: A case series. International Journal of Case Reports and Images. 2014;5(9):610–3. Available from: https://doi.org/10.5348/ijcri-201460-CS-10046 .	Studiedesign
Abd El Maksoud W, Abbas K. Onlay hernioplasty versus Rives–Stoppa repair for paraumbilical hernia associated with divarication of recti in diabetics in terms of recurrence and surgical site infection: a prospective randomized controlled trial. The Egyptian Journal of Surgery. 2018;37(4):453-9. Available from: https://doi.org/10.4103/ejs.ejs_42_18 .	Population
Abdel-Baki NA, Bessa SS, Abdel-Razek AH. Comparison of prosthetic mesh repair and tissue repair in the emergency management of incarcerated para-umbilical hernia: a prospective randomized study. Hernia. 2007;11(2):163-7.	Population
Abrahamson J, Gorman J. Pregnancy and ventral hernia repair. Hernia. 2000;4(4):187-91. Available from: https://doi.org/10.1007/BF01201063 .	Studiedesign
Acharry N, Kutty R. Abdominal exercise with bracing, a therapeutic efficacy in reducing diastasis-recti among postpartal females. Int J Physiother Res. 2015;3:999-1005. Available from: https://doi.org/10.16965/ijpr.2015.122 .	Studiedesign
Al-Basti HB, El-Khatib HA, Taha A, Sattar HA, Bener A. Intraabdominal pressure after full abdominoplasty in obese multiparous patients. Plast Reconstr Surg. 2004;113(7):2145-50; discussion 51-5. Available from: https://doi.org/10.1097/01.prs.0000122543.44977.46 .	Studiedesign
Al-Mansour MR, Wu J, Gagnon G, Knee A, Romanelli JR, Seymour NE. Linear versus volumetric CT analysis in predicting tension-free fascial closure in abdominal wall reconstruction. Hernia. 2021;25(1):91-8. Available from: https://doi.org/10.1007/s10029-020-02349-6 .	Studiedesign
al-Qattan MM. Abdominoplasty in multiparous women with severe musculoaponeurotic laxity. Br J Plast Surg. 1997;50(6):450-5. Available from: https://doi.org/10.1016/s0007-1226(97)90333-7 .	Studiedesign
Al-Shaham A. Neoumbilicoplasty is a useful adjuvant procedure in abdominoplasty. Can J Plast Surg. 2009;17(4):e20-3.	Studiedesign
Ansari S, Elmieh A, Alipour A. The effect of aquatic exercise on functional disability, flexibility and function of trunk muscles in postmenopausal	Population

Referens	Exklusionsorsak
women with chronic non-specific low back pain: randomized controlled trial. <i>Science & Sports</i> 2021 Jun;36(3):e103-e110. 2021.	
Appleton ND, Anderson KD, Hancock K, Scott MH, Walsh CJ. Initial UK experience with transversus abdominis muscle release for posterior components separation in abdominal wall reconstruction of large or complex ventral hernias: a combined approach by general and plastic surgeons. <i>Ann R Coll Surg Engl.</i> 2017;99(4):265-70. Available from: https://doi.org/10.1308/rcsann.2016.0241 .	Population
Armstrong P. Raising pelvic floor awareness among teenage girls. <i>Journal of the Association of Chartered Physiotherapists in Women's Health.</i> 2005(97):55-9.	Population
Asaadi M, Haramis HT. A simple technique for repair of rectus sheath defects. <i>Ann Plast Surg.</i> 1994;32(1):107-9. Available from: https://doi.org/10.1097/00000637-199401000-00017 .	Publikationstyp
Banerjee A, Mahalakshmi V, Baranitharan R. Effect of Antenatal Exercise Program with and without Abdominal Strengthening Exercises on Diastasis Rectus Abdominis - A Post Partum Follow Up. <i>Indian Journal of Physiotherapy and Occupational Therapy - An International Journal.</i> 2013;7(4):123-6. Available from: https://doi.org/10.5958/j.0973-5674.7.4.134 .	Population
Barchi LC, Franciss MY, Zilberstein B. Subcutaneous Videosurgery for Abdominal Wall Defects: A Prospective Observational Study. <i>J Laparoendosc Adv Surg Tech A.</i> 2019;29(4):523-30. Available from: https://doi.org/10.1089/lap.2018.0697 .	Population
Batchvarova Z, Leymarie N, Lepage C, Leyder P. Use of a submuscular resorbable mesh for correction of severe postpregnancy musculoaponeurotic laxity: an 11-year retrospective study. <i>Plast Reconstr Surg.</i> 2008;121(4):1240-8. Available from: https://doi.org/10.1097/01.prs.0000302456.02109.04 .	Studiedesign
Bellido Luque J, Bellido Luque A, Nogales Muñoz A, Sanchez Matamoros I, Lopez Duran LD, Oliva Mompean F. 26th International Congress of the European Association for Endoscopic Surgery (EAES), London, United Kingdom, 30 May-1 June 2018: Video Presentations. <i>Surg Endosc.</i> 2018;32(Suppl 2):361-429. Available from: https://doi.org/10.1007/s00464-018-6179-z .	Publikationstyp
Bellido Luque J, Bellido Luque A, Valdivia J, Suarez Grau JM, Gomez Menchero J, Garcia Moreno J, et al. Totally endoscopic surgery on diastasis recti associated with midline hernias. The advantages of a minimally invasive approach. Prospective cohort study. <i>Hernia.</i> 2015;19(3):493-501. Available from: https://doi.org/https://dx.doi.org/10.1007/s10029-014-1300-2 .	Studiedesign
Bellido Luque J, Suarez Gráu JM, García Moreno J, Bellido Luque A, Gomez Menchero J, Duran Ferreras I, et al. 25th International Congress of the European Association for Endoscopic Surgery (EAES) Frankfurt, Germany, 14-17 June 2017 Video Presentations. <i>Surg Endosc.</i> 2017;31(Suppl 2):190-259. Available from: https://doi.org/10.1007/s00464-017-5542-9 .	Publikationstyp

Referens	Exklusionsorsak
Benito-Ruiz J, de Cabo F. Ultrasonography: a useful tool for plastic surgeons. <i>Aesthetic Plast Surg.</i> 2014;38(3):561-71. Available from: https://doi.org/10.1007/s00266-014-0300-z .	Studiedesign
Bentzon N, Adamsen S. Hernia of the posterior rectus sheath: a new entity? <i>Eur J Surg.</i> 1995;161(3):215-6.	Studiedesign
Bhanot A. Laparoscopic abdominoplasty for divarication of recti. <i>Journal of Chinese Clinical Medicine.</i> 2009;4(12):703-5.	Studiedesign
Bittner R, Schwarz J. Endoscopic mini/less open sublay operation for treatment of primary and secondary ventral hernias of the abdominal wall. <i>European Surgery.</i> 2017;49(2):65-70. Available from: https://doi.org/10.1007/s10353-017-0472-2 .	Population
Blotta RM, Costa SDS, Trindade EN, Meurer L, Maciel-Trindade MR. Collagen I and III in women with diastasis recti. <i>Clinics (Sao Paulo).</i> 2018;73:e319. Available from: https://doi.org/10.6061/clinics/2018/e319 .	Studiedesign
Bowman K. Diastasis Recti: When the Abs Don't Come Together. <i>IDEA Fitness Journal.</i> 2013;10(5):28-31.	Publikationstyp
Brauman D, Capocci J. Liposuction abdominoplasty: an advanced body contouring technique. <i>Plast Reconstr Surg.</i> 2009;124(5):1685-95. Available from: https://doi.org/10.1097/PRS.0b013e3181b98c5d .	Population
Bruner TW, Salazar-Reyes H, Friedman JD. Umbilical hernia repair in conjunction with abdominoplasty: a surgical technique to maintain umbilical blood supply. <i>Aesthet Surg J.</i> 2009;29(4):333-4. Available from: https://doi.org/10.1016/j.asj.2009.02.012 .	Population
Bucaria V, Boccuzzi A. Autologous Dermal Mesh in Severe Postpregnancy Recti Muscles Diastasis. <i>Aesthetic Plast Surg.</i> 2021;45(1):198-211. Available from: https://doi.org/10.1007/s00266-020-01991-w .	Studiedesign
Caldeira AML, Carrion K, Jaulis J. Repair of the Severe Muscle Aponeurotic Abdominal Laxity with Alloplastic Mesh in Aesthetic Abdominoplasty. <i>Aesthetic Plast Surg.</i> 2018;42(4):1039-49. Available from: https://doi.org/10.1007/s00266-018-1101-6 .	Population
Candido G, Lo T, Janssen PA. Risk factors for diastasis of the recti abdominis. <i>Journal of the Association of Chartered Physiotherapists in Women's Health.</i> 2005(97):49-54.	Studiedesign
Capitano S. Laparoscopic transabdominal preperitoneal approach for umbilical hernia with rectus diastasis. <i>Asian J Endosc Surg.</i> 2017;10(3):334-5. Available from: https://doi.org/10.1111/ases.12365 .	Population
Cardaillac C, Vieillefosse S, Oppenheimer A, Joueidi Y, Thubert T, Deffieux X. Diastasis of the rectus abdominis muscles in postpartum: Concordance of patient and clinician evaluations, prevalence, associated pelvic floor symptoms and quality of life. <i>Eur J Obstet Gynecol Reprod Biol.</i> 2020;252:228-32. Available from: https://doi.org/10.1016/j.ejogrb.2020.06.038 .	Studiedesign
Cardenas Restrepo JC, Munoz Ahmed JA. New technique of plication for miniabdominoplasty. <i>Plast Reconstr Surg.</i> 2002;109(3):1170-7; discussion 8-90. Available from: https://doi.org/10.1097/00006534-200203000-00061 .	Studiedesign

Referens	Exklusionsorsak
Carrara A, Catarci M, Fabris L, Zuolo M, Pellecchia L, Moscatelli P, et al. Prospective observational study of abdominal wall reconstruction with THT technique in primary midline defects with diastasis recti: clinical and functional outcomes in 110 consecutive patients. <i>Surg Endosc.</i> 2021;35(9):5104-14. Available from: https://doi.org/10.1007/s00464-020-07997-4 .	Studiedesign
Carrara A, Lauro E, Fabris L, Frisini M, Rizzo S. Endo-laparoscopic reconstruction of the abdominal wall midline with linear stapler, the THT technique. Early results of the first case series. <i>Ann Med Surg (Lond).</i> 2019;38:1-7. Available from: https://doi.org/10.1016/j.amsu.2018.12.002 .	Studiedesign
Cavalli M, Bruni PG, Lombardo F, Morlacchi A, Andretto Amodeo C, Campanelli G. Original concepts in anatomy, abdominal-wall surgery, and component separation technique and strategy. <i>Hernia.</i> 2020;24(2):411-9. Available from: https://doi.org/10.1007/s10029-019-02030-7 .	Population
Champault GG, Catheline JM, Barrat C. Parietoscopic treatment of abdominal wall defects: A report of 15 cases. <i>Hernia.</i> 1999;3(1):15-8. Available from: https://doi.org/10.1007/bf01576734 .	Population
Chang CJ. Assessment of videoendoscopy-assisted abdominoplasty for diastasis recti patients. <i>Biomed J.</i> 2013;36(5):252-6. Available from: https://doi.org/10.4103/2319-4170.113374 .	Studiedesign
Cheesborough JE, Dumanian GA. Simultaneous prosthetic mesh abdominal wall reconstruction with abdominoplasty for ventral hernia and severe rectus diastasis repairs. <i>Plast Reconstr Surg.</i> 2015;135(1):268-76. Available from: https://doi.org/10.1097/PRS.0000000000000840 .	Studiedesign
Chiarello CM, Falzone LA, McCaslin KE, Patel MN, Ulery KR. The effects of an exercise program on diastasis recti abdominis in pregnant women [corrected] [published erratum appears in J WOMENS HEALTH PHYS THER 2005;29(3):76]. <i>Journal of Women's Health Physical Therapy.</i> 2005;29(1):11-6.	Population
Cintesun E, Incesu Cintesun FN, Aydogdu M, Bayramoglu D, Celik C. Effect of re-approximation of the rectus muscles on diastasis recti abdominis at cesarean section - a prospective cross-sectional study. <i>Ginekol Pol.</i> 2021;92(2):132-6. Available from: https://doi.org/10.5603/GP.a2020.0143 .	Population
Claus C, Cavazzola L, Malcher F. SubCutaneous OnLay endoscopic Approach (SCOLA) for midline ventral hernias associated with diastasis recti. <i>Hernia.</i> 2021;25(4):957-62. Available from: https://doi.org/10.1007/s10029-021-02465-x .	Publikationstyp
Claus CMP, Malcher F, Cavazzola LT, Furtado M, Morrell A, Azevedo M, et al. Subcutaneous Onlay Laparoscopic Approach (Scola) for Ventral Hernia and Rectus Abdominis Diastasis Repair: Technical Description and Initial Results. <i>Arq Bras Cir Dig.</i> 2018;31(4):e1399. Available from: https://doi.org/10.1590/0102-672020180001e1399 .	Population
Colwell AS, Kpodzo D, Gallico GG, 3rd. Low scar abdominoplasty with inferior positioning of the umbilicus. <i>Ann Plast Surg.</i> 2010;64(5):639-44. Available from: https://doi.org/10.1097/SAP.0b013e3181db759c .	Intervention

Referens	Exklusionsorsak
Core GB, Mizgala CL, Bowen JC, 3rd, Vasconez LO. Endoscopic abdominoplasty with repair of diastasis recti and abdominal wall hernia. <i>Clin Plast Surg.</i> 1995;22(4):707-22.	Studiedesign
Correa MA. Videoendoscopic subcutaneous techniques for aesthetic and reconstructive plastic surgery. <i>Plast Reconstr Surg.</i> 1995;96(2):446-53. Available from: https://doi.org/10.1097/00006534-199508000-00030 .	Studiedesign
Cossa JP, Ngo P, Pelissier E. Stapled VTEP (sVTEP), diastasis and the "swollen abdomen". <i>Surg Endosc.</i> 2021. Available from: https://doi.org/10.1007/s00464-021-08656-y .	Studiedesign
Criss CN, Petro CC, Krpata DM, Seafler CM, Lai N, Fiutem J, et al. Functional abdominal wall reconstruction improves core physiology and quality-of-life. <i>Surgery.</i> 2014;156(1):176-82. Available from: https://doi.org/10.1016/j.surg.2014.04.010 .	Studiedesign
Cuccomarino S, Bonomo LD, Apra F, Toscano A, Jannaci A. Preaponeurotic endoscopic repair (REPA) of diastasis recti: a single surgeon's experience. <i>Surg Endosc.</i> 2021. Available from: https://doi.org/10.1007/s00464-021-08405-1 .	Studiedesign
Dabb RW, Hall WW, Baroody M, Saba AA. Circumferential suction lipectomy of the trunk with anterior rectus fascia plication through a periumbilical incision: an alternative to conventional abdominoplasty. <i>Plast Reconstr Surg.</i> 2004;113(2):727-32; discussion 33-4. Available from: https://doi.org/10.1097/01.PRS.0000101508.44801.F6 .	Studiedesign
Dalal K, Kaur A, Mitra M. Correlation between Diastasis Rectus Abdominis and Lumbopelvic Pain and Dysfunction. <i>Indian Journal of Physiotherapy and Occupational Therapy - An International Journal.</i> 2014;8(1):210-4. Available from: https://doi.org/10.5958/j.0973-5674.8.1.040 .	Studiedesign
Danilla S. Rectus Abdominis Fat Transfer (RAFT) in Lipoabdominoplasty: A New Technique to Achieve Fitness Body Contour in Patients that Require Tummy Tuck. <i>Aesthetic Plast Surg.</i> 2017;41(6):1389-99. Available from: https://doi.org/10.1007/s00266-017-0909-9 .	Intervention
de Brito MJ, Nahas FX, Barbosa MV, Dini GM, Kimura AK, Farah AB, et al. Abdominoplasty and its effect on body image, self-esteem, and mental health. <i>Ann Plast Surg.</i> 2010;65(1):5-10. Available from: https://doi.org/10.1097/SAP.0b013e3181bc30f7 .	Population
de Castro EJP, Radwanski HN, Pitanguy I, Nahas F. Long-term ultrasonographic evaluation of midline aponeurotic plication during abdominoplasty. <i>Plast Reconstr Surg.</i> 2013;132(2):333-8. Available from: https://doi.org/10.1097/PRS.0b013e3182958ad2 .	Studiedesign
de León SC, Carrera IDC, González CADLH, González MS. Comparison of the effects of an exercise program on straight diastasis according to the pos-tpart period. <i>Medicina Naturista.</i> 2020;14(1):53-8.	Språk
De'Ath HD, Lovegrove RE, Javid M, Peter N, Magee TR, Galland RB. An assessment of between-recti distance and divarication in patients with and without abdominal aortic aneurysm. <i>Ann R Coll Surg Engl.</i> 2010;92(7):591-4. Available from: https://doi.org/10.1308/003588410X12771863937089 .	Population

Referens	Exklusionsorsak
Deering RE, Chumanov ES, Stiffler-Joachim MR, Heiderscheit BC. Exercise Program Reduces Inter-Recti Distance in Female Runners Up to 2 Years Postpartum. <i>Journal of Women's Health Physical Therapy</i> . 2020;44(1):9-18. Available from: https://doi.org/10.1097/jwh.0000000000000157 .	Studiedesign
Deering RE, Senefeld J, Pashibin T, Neumann DA, Cruz M, Hunter SK. Fatigability of the Lumbopelvic Stabilizing Muscles in Women 8 and 26 Weeks Postpartum. <i>J Womens Health Phys Therap</i> . 2018;42(3):128-38. Available from: https://doi.org/10.1097/JWH.0000000000000109 .	Studiedesign
Dessy LA, Mazzocchi M, Fallico N, Annibaletti T, Scuderi N. Association between abdominal separation and inguinal or crural hernias: our experience and surgical indications. <i>J Plast Surg Hand Surg</i> . 2013;47(3):209-12. Available from: https://doi.org/10.3109/2000656X.2012.742444 .	Utfall
Dewan PA. Rectus abdominis muscle flap technique for the management of large defects. <i>Br J Urol</i> . 1997;80(3):511.	Publikationstyp
Dey A, Malik VK, Kumar A, Vashishta V, Agarwal B. Posterior component separation by transversus abdominis release-current paradigm in management of large complex hernias: Our initial experience. <i>Current Medicine Research and Practice</i> . 2018;8(4):134-7. Available from: https://doi.org/10.1016/j.cmrp.2018.06.001 .	Population
di Diego JM. Endoscopic Lipoabdominoplasty. <i>Plast Reconstr Surg Glob Open</i> . 2021;9(1):e3346. Available from: https://doi.org/10.1097/GOX.0000000000003346 .	Studiedesign
Di Martino M, Nahas FX, Barbosa MVJ, Montecinos Ayaviri NA, Kimura AK, Barella SM, et al. Seroma in lipoabdominoplasty and abdominoplasty: a comparative study using ultrasound. <i>Plast Reconstr Surg</i> . 2010;126(5):1742-51. Available from: https://doi.org/10.1097/PRS.0b013e3181efa6c5 .	Population
Dong CT, Sreeramoju P, Pechman DM, Weithorn D, Camacho D, Malcher F. SubCutaneous OnLay endoscopic Approach (SCOLA) mesh repair for small midline ventral hernias with diastasis recti: An initial US experience. <i>Surg Endosc</i> . 2020. Available from: https://doi.org/10.1007/s00464-020-08134-x .	Studiedesign
Dudai M, Gilboa Ittah K. Intraoperative hypertonic saline irrigation preventing seroma formation and reducing drain secretion in extended endoscopic hernia and linea alba reconstruction glue. <i>Hernia</i> . 2019;23(6):1291-6. Available from: https://doi.org/10.1007/s10029-019-01956-2 .	Population
Dufour S, Bernard S, Murray-Davis B, Graham N. Establishing Expert-Based Recommendations for the Conservative Management of Pregnancy-Related Diastasis Rectus Abdominis: A Delphi Consensus Study. <i>Journal of Women's Health Physical Therapy</i> . 2019;43(2):73-81. Available from: https://doi.org/10.1097/jwh.0000000000000130 .	Publikationstyp
Dufour SP, Graham N, Bernard S, Murray-Davis B. Section on Women's Health Combined Sections Meeting Platforms. <i>Journal of Women's Health Physical Therapy</i> . 2018;42(1):32-6. Available from: https://doi.org/10.1097/jwh.0000000000000090 .	Publikationstyp

Referens	Exklusionsorsak
Dumanian GA, Cheesborough JE. Reply: Simultaneous Prosthetic Mesh Abdominal Wall Reconstruction with Abdominoplasty for Ventral Hernia and Severe Rectus Diastasis Repairs. <i>Plast Reconstr Surg.</i> 2016;137(1):254e-5e. Available from: https://doi.org/10.1097/PRS.0000000000001913 .	Publikationstyp
East JM. Mesh tuck repair of ventral hernias of the abdomen: a new, simplified technique for sublay herniorrhaphy. <i>West Indian Med J.</i> 2007;56(6):514-9.	Population
El-Mekawy HS, Eldeeb A, Lythy MAE, Elbegawy AF. Effect of Abdominal Exercises versus Abdominal Supporting Belt on Post-Partum Abdominal Efficiency and Rectus Separation. <i>International Journal of Medical and Health Sciences.</i> 2013;7:75-9. Available from: https://doi.org/10.5281/zenodo.1334606 .	Population
Elkhatib H, Buddhavarapu SR, Henna H, Kassem W. Abdominal musculoaponeuretic system: magnetic resonance imaging evaluation before and after vertical plication of rectus muscle diastasis in conjunction with lipoabdominoplasty. <i>Plast Reconstr Surg.</i> 2011;128(6):733e-40e. Available from: https://doi.org/10.1097/PRS.0b013e318230c8a1 .	Studiedesign
Farouk M, Askar A. Crossed abdominal wall plication in abdominoplasty. <i>Journal of Taibah University Medical Sciences.</i> 2013;8(2):115-9. Available from: https://doi.org/10.1016/j.jtumed.2013.07.003 .	Population
Felfel M, El Khoury M, Marboeuf Y, Strohl D, Menu Y. Incarcerated hernia through the posterior rectus sheath. <i>AJR Am J Roentgenol.</i> 2005;185(5):1185-6. Available from: https://doi.org/10.2214/AJR.04.0637 .	Studiedesign
Fernandes da Mota PG, Pascoal AG, Carita AI, Bo K. Prevalence and risk factors of diastasis recti abdominis from late pregnancy to 6 months postpartum, and relationship with lumbo-pelvic pain. <i>Man Ther.</i> 2015;20(1):200-5. Available from: https://doi.org/10.1016/j.math.2014.09.002 .	Studiedesign
Ferreira LM, Castilho HT, Hochberg J, Ardenghi M, Toledo SR, Cruz RG, et al. Triangular mattress suture in abdominal diastasis to prevent epigastric bulging. <i>Ann Plast Surg.</i> 2001;46(2):130-4. Available from: https://doi.org/10.1097/00000637-200102000-00007 .	Publikationstyp
Fiori F, Ferrara F, Gobatti D, Gentile D, Stella M. Surgical treatment of diastasis recti: the importance of an overall view of the problem. <i>Hernia.</i> 2021;25(4):871-82. Available from: https://doi.org/10.1007/s10029-020-02252-0 .	Studiedesign
Fiori F, Ferrara F, Gobatti D, Gentile D, Stella M. Surgical treatment of diastasis recti: the importance of an overall view of the problem. <i>Hernia.</i> 2021;25(4):871-82. Available from: https://doi.org/10.1007/s10029-020-02252-0 .	Studiedesign
Fuenmayor P, Lujan HJ, Plasencia G, Karmaker A, Mata W, Vecin N. Robotic-assisted ventral and incisional hernia repair with hernia defect closure and intraperitoneal onlay mesh (IPOM) experience. <i>J Robot Surg.</i>	Population

Referens	Exklusionsorsak
2020;14(5):695-701. Available from: https://doi.org/10.1007/s11701-019-01040-y .	
Gama LJM, Barbosa MVJ, Czapkowski A, Ajzen S, Ferreira LM, Nahas FX. Single-Layer Plication for Repair of Diastasis Recti: The Most Rapid and Efficient Technique. <i>Aesthet Surg J.</i> 2017;37(6):698-705. Available from: https://doi.org/10.1093/asj/sjw263 .	Population
Gandhi JA, Shinde P, Kothari B, Churiwala JJ, Banker A. Endoscopic Pre-aponeurotic Repair (EPAR) Technique with Meshplasty for Treatment of Ventral Hernia and Rectus Abdominis Diastasis. <i>Indian Journal of Surgery.</i> 2020. Available from: https://doi.org/10.1007/s12262-020-02189-9 .	Studiedesign
Gangi S, Sparacino T, Furci M, Basile F. Hernia of the posterior lamina of the rectus abdominis muscle sheath: report of a case. <i>Ann Ital Chir.</i> 2002;73(3):335-7.	Studiedesign
García-Pastor P, Torregrosa A, Sancho-Muriel J, Argüelles B, Bonafe S, Iserte J, et al. Oral Communications. <i>Hernia.</i> 2016;20 Suppl 2:139-258. Available from: https://doi.org/10.1007/s10029-016-1482-x .	Publikationstyp
García-Pastor P, Torregrosa-Gallud A, Sancho J, Lopez M, Blasco Piqueras R, Bonafé S, et al. 2018 International Hernia Congress : March 12-15, 2018 Miami, FL. <i>Hernia.</i> 2018;22(Suppl 1):1-204. Available from: https://doi.org/10.1007/s10029-018-1733-0 .	Publikationstyp
Garvey PB, Bailey CM, Baumann DP, Liu J, Butler CE. Violation of the rectus complex is not a contraindication to component separation for abdominal wall reconstruction. <i>J Am Coll Surg.</i> 2012;214(2):131-9. Available from: https://doi.org/10.1016/j.jamcollsurg.2011.10.015 .	Studiedesign
Giurgis M, Bendure L, Davenport DL, Roth JS. The endoscopic component separation technique for hernia repair results in reduced morbidity compared to the open component separation technique. <i>Hernia.</i> 2012;16(1):47-51. Available from: https://doi.org/10.1007/s10029-011-0866-1 .	Studiedesign
Gluppe SB, Engh ME, Bo K. Immediate Effect of Abdominal and Pelvic Floor Muscle Exercises on Interrecti Distance in Women With Diastasis Recti Abdominis Who Were Parous. <i>Phys Ther.</i> 2020;100(8):1372-83. Available from: https://doi.org/10.1093/ptj/pzaa070 .	Studiedesign
Gnaneswaran N, Perera M, Jenkin A, Lau H, Presley R. Ventral hernia repair with lateral component separation and onlay Biodesign graft. <i>European Journal of Plastic Surgery.</i> 2016;39(4):279-86. Available from: https://doi.org/10.1007/s00238-016-1188-4 .	Population
Gomez-Menchero J, Guadalajara Jurado JF, Suarez Grau JM, Bellido Luque JA, Garcia Moreno JL, Alarcon Del Agua I, et al. Laparoscopic intracorporeal rectus aponeuroplasty (LIRA technique): a step forward in minimally invasive abdominal wall reconstruction for ventral hernia repair (LVHR). <i>Surg Endosc.</i> 2018;32(8):3502-8. Available from: https://doi.org/10.1007/s00464-018-6070-y .	Population
Gursen C, Inanoglu D, Kaya S, Akbayrak T, Baltaci G. Effects of exercise and Kinesio taping on abdominal recovery in women with cesarean section: a pilot randomized controlled trial. <i>Arch Gynecol Obstet.</i>	Population

Referens	Exklusionsorsak
2016;293(3):557-65. Available from: https://doi.org/10.1007/s00404-015-3862-3 .	
Guzman-Valdivia Gomez G. Mesh prosthesis versus triangular flap from the anterior sheath of the abdominal rectus muscle in primary inguinal hernia repair. <i>J Invest Surg.</i> 2003;16(1):45-50.	Population
Hanssen A, Palmisano EM, Hanssen DA, Hanssen RA, Daes JE. Vasomotor changes in abdominal skin after endoscopic subcutaneous/Preaponeurotic Repair of Diastasis Recti (REPA). <i>Int J Surg Case Rep.</i> 2020;75:182-4. Available from: https://doi.org/10.1016/j.ijscr.2020.08.056 .	Studiedesign
Harlaar JJ, Deerenberg EB, Dwarkasing RS, Kamperman AM, Kleinrensink GJ, Jeekel J, et al. Development of incisional herniation after midline laparotomy. <i>BJS Open.</i> 2017;1(1):18-23. Available from: https://doi.org/10.1002/bjs5.3 .	Population
Harlaar JJ, Deerenberg EB, van Ramshorst GH, Lont HE, van der Borst EC, Schouten WR, et al. A multicenter randomized controlled trial evaluating the effect of small stitches on the incidence of incisional hernia in midline incisions. <i>BMC Surg.</i> 2011;11:20. Available from: https://doi.org/10.1186/1471-2482-11-20 .	Publikationstyp
Hicks CW, Krpata DM, Blatnik JA, Novitsky YW, Rosen MJ. Long-term effect on donor sites after components separation: a radiographic analysis. <i>Plast Reconstr Surg.</i> 2012;130(2):354-9. Available from: https://doi.org/10.1097/PRS.0b013e3182589d79 .	Studiedesign
Hodgman EI, Watson MJ. Revisiting the Anterior Rectus Sheath Repair for Incisional Hernia: A 10-Year Experience. <i>World J Surg.</i> 2017;41(3):713-21. Available from: https://doi.org/10.1007/s00268-016-3774-9 .	Population
Holm C. Perfusion zones of the DIEP flap revisited: a clinical study. <i>Plast Reconstr Surg.</i> 2006;118(4):1077. Available from: https://doi.org/10.1097/01.prs.0000233604.86936.e8 .	Publikationstyp
Horzic M, Kopljar M, Cupurdija K, Skupnjak M, Korica J, Lackovic Z, et al. Quality of life changes after inguinal hernia repair using anterior rectus sheath--a preliminary study. <i>Coll Antropol.</i> 2006;30(2):349-53.	Population
Huang GJ, Bajaj AK, Gupta S, Petersen F, Miles DAG. Increased intraabdominal pressure in abdominoplasty: delineation of risk factors. <i>Plast Reconstr Surg.</i> 2007;119(4):1319-25. Available from: https://doi.org/10.1097/01.prs.0000254529.51696.43 .	Studiedesign
Hughes KC, Weider L, Fischer J, Hopkins J, Antonetti A, Manders EK, et al. Ventral hernia repair with simultaneous panniculectomy. <i>Am Surg.</i> 1996;62(8):678-81.	Studiedesign
Iglesias M, Bravo L, Chavez-Munoz C, Barajas-Olivas A. Endoscopic abdominoplasty: an alternative approach. <i>Ann Plast Surg.</i> 2006;57(5):489-94. Available from: https://doi.org/10.1097/01.sap.0000229004.25059.87 .	Studiedesign
Ito F, Tsuchiya T, Otani S, Saito T, Matsuida H, Miura J. Scientific Session of the 16th World Congress of Endoscopic Surgery, Jointly Hosted by Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) & Canadian Association of General Surgeons (CAGS), Seattle, Washington,	Publikationstyp

Referens	Exklusionsorsak
USA, 11-14 April 2018: Poster Abstracts. <i>Surg Endosc.</i> 2018;32(Suppl 1):130-359. Available from: https://doi.org/10.1007/s00464-018-6121-4 .	
Jackson TL, Jackson RF, Freeman L. Minimally invasive abdominoplasty: surgical technique development and report of three cases. <i>Surg Laparosc Endosc.</i> 1995;5(4):301-5.	Studiedesign
Jacob CI, Rank B. Abdominal Remodeling in Postpartum Women by Using a High-intensity Focused Electromagnetic (HIFEM) Procedure: An Investigational Magnetic Resonance Imaging (MRI) Pilot Study. <i>The Journal of clinical and aesthetic dermatology.</i> 2020;13(9 Suppl 1):S16-S20.	Studiedesign
Jansen DA, Gailliot RV, Jr., Galli RA, Escobar JR, Kind G, Parry SW. An evaluation of fascial staples (a new technique) in wide fascial plication during reconstructive abdominoplasty. <i>Ann Plast Surg.</i> 1996;36(2):171-5. Available from: https://doi.org/10.1097/00000637-199602000-00012 .	Population
Jeffcoat H. Exercises for low back pain in pregnancy. <i>International Journal of Childbirth Education.</i> 2008;23(3):9-12.	Publikationstyp
Jones GG, Saour S, Botha AJ, Healy CM. Surgical tip--a novel method of applying fibrin sealant during repair of divarication of the recti via an abdominoplasty incision. <i>J Plast Reconstr Aesthet Surg.</i> 2009;62(11):e457-8. Available from: https://doi.org/10.1016/j.bjps.2008.08.035 .	Studiedesign
Joshi MA, Singh MB, Gadhire MA. Study of the outcome of modified shoelace repair for midline incisional hernia. <i>Hernia.</i> 2015;19(3):503-8. Available from: https://doi.org/10.1007/s10029-014-1234-8 .	Population
Juárez Muas D, Palmisano E, Santoja G, Mustone Paz O. Adjuvant botulinum toxin for endoscopic management (preaponeurotic endoscopic repair) of severe diastasis recti. <i>International Journal of Abdominal Wall and Hernia Surgery.</i> 2021;4(2):45-50. Available from: https://doi.org/10.4103/ijawhs.ijawhs_49_20 .	Studiedesign
Juarez Muas DM. Preaponeurotic endoscopic repair (REPA) of diastasis recti associated or not to midline hernias. <i>Surg Endosc.</i> 2019;33(6):1777-82. Available from: https://doi.org/10.1007/s00464-018-6450-3 .	Population
Kaddoura IL, Nasser A. Abdominoplasty: the use of stapled prolene mesh in severe musculoaponeurotic laxity. <i>Aesthet Surg J.</i> 1998;18(5):335-41. Available from: https://doi.org/10.1016/s1090-820x(98)70088-3 .	Studiedesign
Kale AA, Bellare B. Awareness about Exclusive Role of Postnatal Physical Therapy: A Preliminary Survey Conducted on Obstetricians at Aurangabad, Maharashtra. <i>Indian Journal of Physiotherapy and Occupational Therapy - An International Journal.</i> 2019;13(3):27-30. Available from: https://doi.org/10.5958/0973-5674.2019.00086.8 .	Studiedesign
Kalinovskiy SV, Vlasov VV, Pidmurnyak OO, Kharyshyn OM, Vlasova KV. Oral Communications. <i>Hernia.</i> 2016;20 Suppl 2:139-258. Available from: https://doi.org/10.1007/s10029-016-1482-x .	Publikationstyp
Kanjoor JR, Singh AK. Lipoabdominoplasty: An exponential advantage for a consistently safe and aesthetic outcome. <i>Indian J Plast Surg.</i> 2012;45(1):77-88. Available from: https://doi.org/10.4103/0970-0358.96592 .	Studiedesign

Referens	Exklusionsorsak
Karthikesalingam A, Kitcat M, Malata CM. Abdominoplasty in patients with and without pre-existing scars: a retrospective comparison. <i>J Plast Reconstr Aesthet Surg.</i> 2011;64(3):369-74. Available from: https://doi.org/10.1016/j.bjps.2010.05.025 .	Studiedesign
Kazmi K, Hashmi GM, Bilal M, Iqbal N, Wajeeha, Ilyas A, et al. Effect of Abdominal Physiotherapy in Reduction of Inter Recti Distance in Postpartum Females. <i>Pakistan Journal of Medical and Health Sciences.</i> 2021;15(6):1446-8. Available from: https://doi.org/10.53350/pjmhs211561446 .	Studiedesign
Keeler J, Albrecht M, Eberhardt L, Horn L, Donnelly C, Lowe D. Diastasis Recti Abdominis. <i>Journal of Women's Health Physical Therapy.</i> 2012;36(3):131-42. Available from: https://doi.org/10.1097/JWH.0b013e318276f35f .	Studiedesign
Kent DE, Jacob CI. Simultaneous Changes in Abdominal Adipose and Muscle Tissues Following Treatments by High-Intensity Focused Electromagnetic (HIFEM) Technology-Based Device: Computed Tomography Evaluation. <i>J Drugs Dermatol.</i> 2019;18(11):1098-102.	Population
Keshwani N, Mathur S, McLean L. Relationship Between Interrectus Distance and Symptom Severity in Women With Diastasis Recti Abdominis in the Early Postpartum Period. <i>Phys Ther.</i> 2018;98(3):182-90. Available from: https://doi.org/10.1093/ptj/pzx117 .	Studiedesign
Khandale SR, Hande D. Effects of Abdominal Exercises on Reduction of Diastasis Recti in Postnatal Women. <i>Int J Health Sci Res.</i> 2016;6(6):182-91.	Studiedesign
Khansa I, Janis J. 17th Annual Hernia Repair : March 30-April 2, 2016 Washington DC, USA. <i>Hernia.</i> 2016;20 Suppl 1(1):1-138. Available from: https://doi.org/10.1007/s10029-016-1468-8 .	Publikationstyp
Khetan M, Dey A, Bindal V, Suviraj J, Mittal T, Kalhan S, et al. e-TEP repair for midline primary and incisional hernia: technical considerations and initial experience. <i>Hernia.</i> 2021. Available from: https://doi.org/10.1007/s10029-021-02397-6 .	Population
Kim JK, Jang JY, Hong YG, Sim HB, Sun SH. Deep-Plane Lipoabdominoplasty in East Asians. <i>Arch Plast Surg.</i> 2016;43(4):352-9. Available from: https://doi.org/10.5999/aps.2016.43.4.352 .	Studiedesign
Kinder J, Davenport T, Chong W, Lee A. Telerehabilitation for Treating Pelvic Floor Dysfunction: A Case Series of 3 Patients' Experiences. <i>Journal of Women's Health Physical Therapy.</i> 2019;43(1):44-50. Available from: https://doi.org/10.1097/jwh.0000000000000120 .	Studiedesign
Kinney BM, Kent DE. American Society for Laser Medicine and Surgery Abstracts. <i>Lasers Surg Med.</i> 2019;51(S30):S4-S59. Available from: https://doi.org/10.1002/lsm.23059 .	Publikationstyp
Kinney BM, Kent DE. MRI and CT Assessment of Abdominal Tissue Composition in Patients After High-Intensity Focused Electromagnetic Therapy Treatments: One-Year Follow-Up. <i>Aesthet Surg J.</i> 2020;40(12):NP686-NP93. Available from: https://doi.org/10.1093/asj/sjaa052 .	Studiedesign

Referens	Exklusionsorsak
Kinney BM, Lozanova P. High intensity focused electromagnetic therapy evaluated by magnetic resonance imaging: Safety and efficacy study of a dual tissue effect based non-invasive abdominal body shaping. <i>Lasers Surg Med.</i> 2019;51(1):40-6. Available from: https://doi.org/10.1002/lsm.23024 .	Publikationstyp
Klein F, Ospina C, Rudolph B, Wustefeld J, Denecke T, Neuhaus P, et al. Formation of a chronic pain syndrome due to mesh shrinkage after laparoscopic intraperitoneal onlay mesh (IPOM). <i>Surg Laparosc Endosc Percutan Tech.</i> 2012;22(5):e288-90. Available from: https://doi.org/10.1097/SLE.0b013e31825efc3c .	Population
Kler A, Wilson P. Total endoscopic-assisted linea alba reconstruction (TESLAR) for treatment of umbilical/paraumbilical hernia and rectus abdominus diastasis is associated with unacceptable persistent seroma formation: a single centre experience. <i>Hernia.</i> 2020;24(6):1379-85. Available from: https://doi.org/10.1007/s10029-020-02266-8 .	Population
Ko JH, Wang EC, Salvay DM, Paul BC, Dumanian GA. Abdominal wall reconstruction: lessons learned from 200 "components separation" procedures. <i>Arch Surg.</i> 2009;144(11):1047-55. Available from: https://doi.org/10.1001/archsurg.2009.192 .	Studiedesign
Kockerling F, Botsinis MD, Rohde C, Reinbold W. Endoscopic-Assisted Linea Alba Reconstruction plus Mesh Augmentation for Treatment of Umbilical and/or Epigastric Hernias and Rectus Abdominis Diastasis - Early Results. <i>Front Surg.</i> 2016;3:27. Available from: https://doi.org/10.3389/fsurg.2016.00027 .	Population
Kockerling F, Botsinis MD, Rohde C, Reinbold W, Schug-Pass C. Endoscopic-assisted linea alba reconstruction: New technique for treatment of symptomatic umbilical, trocar, and/or epigastric hernias with concomitant rectus abdominis diastasis. <i>Eur Surg.</i> 2017;49(2):71-5. Available from: https://doi.org/10.1007/s10353-017-0473-1 .	Population
Kohler G, Fischer I, Kaltenbock R, Schrittwieser R. Minimal Invasive Linea Alba Reconstruction for the Treatment of Umbilical and Epigastric Hernias with Coexisting Rectus Abdominis Diastasis. <i>J Laparoendosc Adv Surg Tech A.</i> 2018;28(10):1223-8. Available from: https://doi.org/10.1089/lap.2018.0018 .	Population
Kohler G, Kaltenbock R, Pfandner R, Dauser B, Lechner M. Precostal top-down extended totally extraperitoneal ventral hernia plasty (eTEP): simplification of a complex technical approach. <i>Hernia.</i> 2020;24(3):527-35. Available from: https://doi.org/10.1007/s10029-019-02076-7 .	Population
Kohler G, Kaltenbock R, Pfandner R, Dauser B, Lechner M. Precostal top-down extended totally extraperitoneal ventral hernia plasty (eTEP): simplification of a complex technical approach. <i>Hernia.</i> 2020;24(3):527-35. Available from: https://doi.org/10.1007/s10029-019-02076-7 .	Population
Kohler G, Luketina RR, Emmanuel K. Sutured repair of primary small umbilical and epigastric hernias: concomitant rectus diastasis is a significant risk factor for recurrence. <i>World J Surg.</i> 2015;39(1):121-6; discussion 7. Available from: https://doi.org/10.1007/s00268-014-2765-y .	Population

Referens	Exklusionsorsak
Koshima I, Nanba Y, Tutsui T, Takahashi Y, Itoh S, Kobayashi R. Dynamic reconstruction of large abdominal defects using a free rectus femoris musculocutaneous flap with normal motor function. <i>Ann Plast Surg.</i> 2003;50(4):420-4. Available from: https://doi.org/10.1097/01.SAP.0000032304.45784.F6 .	Studiedesign
Kroese LF, van Eeghem LHA, Verhelst J, Jeekel J, Kleinrensink GJ, Lange JF. Long term results of open complex abdominal wall hernia repair with self-gripping mesh: A retrospective cohort study. <i>Int J Surg.</i> 2017;44:255-9. Available from: https://doi.org/10.1016/j.ijsu.2017.07.029 .	Population
Kudsi OY, Orthopoulos G, Bhurtel P, Shah J. 2017 Scientific Session of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) Houston, Texas, USA, 22-25 March 2017 : Poster Presentations. <i>Surg Endosc.</i> 2017;31(Suppl 1):137-334. Available from: https://doi.org/10.1007/s00464-017-5445-9 .	Publikationstyp
Kulacoglu H. Umbilical Hernia Repair and Pregnancy: Before, during, after. <i>Front Surg.</i> 2018;5:1. Available from: https://doi.org/10.3389/fsurg.2018.00001 .	Publikationstyp
Kulaylat MN, Karakousis CP. Large gaps of midline abdominal incisions and their management. <i>Am Surg.</i> 2008;74(11):1094-9.	Population
Kulhanek J, Mestak O. Treatment of umbilical hernia and recti muscles diastasis without a periumbilical incision. <i>Hernia.</i> 2013;17(4):527-30. Available from: https://doi.org/10.1007/s10029-013-1047-1 .	Studiedesign
Kurz J, Borello-France D. Movement System Impairment-Guided Approach to the Physical Therapist Treatment of a Patient With Postpartum Pelvic Organ Prolapse and Mixed Urinary Incontinence: Case Report. <i>Phys Ther.</i> 2017;97(4):464-77. Available from: https://doi.org/10.2522/ptj.20160035 .	Studiedesign
Lari A, Curinga P, Person H, Demian H, Braye F, Mabrut J, et al. Abdominoplasty with simultaneous laparoscopic umbilical hernia repair: A practical approach to preserve the umbilical vascularization. <i>Ann Chir Plast Esthet.</i> 2019;64(3):237-44. Available from: https://doi.org/10.1016/j.anplas.2018.06.015 .	Population
Lasheen AE. Mesh expansion and fixation at the recto-rectus plane through multiple stabs by using two tip hole needles in midline hernias repair. A prospective study. <i>Int J Surg.</i> 2008;6(5):367-70. Available from: https://doi.org/10.1016/j.ijsu.2008.06.002 .	Population
Lee D, Hodges PW. Behavior of the Linea Alba During a Curl-up Task in Diastasis Rectus Abdominis: An Observational Study. <i>J Orthop Sports Phys Ther.</i> 2016;46(7):580-9. Available from: https://doi.org/10.2519/jospt.2016.6536 .	Studiedesign
Li B, Qin C, Bittner R. Totally endoscopic sublay (TES) repair for midline ventral hernia: surgical technique and preliminary results. <i>Surg Endosc.</i> 2020;34(4):1543-50. Available from: https://doi.org/10.1007/s00464-018-6568-3 .	Population
Li B, Qin C, Liu D, Miao J, Yu J, Bittner R. Subxiphoid top-down endoscopic totally preperitoneal approach (eTPA) for midline ventral hernia repair.	Population

Referens	Exklusionsorsak
Langenbecks Arch Surg. 2021;406(6):2125-32. Available from: https://doi.org/10.1007/s00423-021-02259-w .	
Liaw LJ, Hsu MJ, Liao CF, Liu MF, Hsu AT. The relationships between interrecti distance measured by ultrasound imaging and abdominal muscle function in postpartum women: a 6-month follow-up study. J Orthop Sports Phys Ther. 2011;41(6):435-43. Available from: https://doi.org/10.2519/jospt.2011.3507 .	Studiedesign
Lincenberg SM. The Retro-Rectus Prosthesis for Core Myofascial Restoration in Cosmetic Abdominoplasty. Aesthet Surg J. 2017;37(8):930-8. Available from: https://doi.org/10.1093/asj/sjx027 .	Studiedesign
Lisiecki J, Kozlow JH, Agarwal S, Ranganathan K, Terjimanian MN, Rinkinen J, et al. Abdominal wall dynamics after component separation hernia repair. J Surg Res. 2015;193(1):497-503. Available from: https://doi.org/10.1016/j.jss.2014.08.008 .	Population
Litos K. Progressive Therapeutic Exercise Program for Successful Treatment of a Postpartum Woman With a Severe Diastasis Recti Abdominis. Journal of Women's Health Physical Therapy. 2014;38(2):58-73. Available from: https://doi.org/10.1097/jwh.0000000000000013 .	Studiedesign
Lo T, Candido G, Janssen P. Diastasis of the Recti abdominis in pregnancy: risk factors and treatment. Physiotherapy Canada. 1999;51(1):32-44.	Studiedesign
Losanoff JE, Basson MD, Gruber SA. Spontaneous hernia through the posterior rectus abdominis sheath: case report and review of the published literature 1937-2008. Hernia. 2009;13(5):555-8. Available from: https://doi.org/10.1007/s10029-009-0481-6 .	Studiedesign
Lu R, Addo A, Ewart Z, Broda A, Parlacoski S, Zahiri HR, et al. Comparative review of outcomes: laparoscopic and robotic enhanced-view totally extraperitoneal (eTEP) access retrorectus repairs. Surg Endosc. 2020;34(8):3597-605. Available from: https://doi.org/10.1007/s00464-019-07132-y .	Population
Lucon M, Ianhez LE, Lucon AM, Chambo JL, Sabbaga E, Srougi M. Bilateral nephrectomy of huge polycystic kidneys associated with a rectus abdominis diastasis and umbilical hernia. Clinics (Sao Paulo). 2006;61(6):529-34. Available from: https://doi.org/10.1590/s1807-59322006000600007 .	Population
Malik A, Macdonald AD, de Beaux AC, Tulloh BR. The peritoneal flap hernioplasty for repair of large ventral and incisional hernias. Hernia. 2014;18(1):39-45. Available from: https://doi.org/10.1007/s10029-013-1086-7 .	Population
Malik K, Bowers SP, Smith CD, Asbun H, Preissler S. A case series of laparoscopic components separation and rectus medialization with laparoscopic ventral hernia repair. J Laparoendosc Adv Surg Tech A. 2009;19(5):607-10. Available from: https://doi.org/10.1089/lap.2009.0155 .	Population
Maloney SR, Schlosser KA, Prasad T, Kasten KR, Gersin KS, Colavita PD, et al. Twelve years of component separation technique in abdominal wall reconstruction. Surgery. 2019;166(4):435-44. Available from: https://doi.org/10.1016/j.surg.2019.05.043 .	Population

Referens	Exklusionsorsak
Manetti G, Lolli MG, Belloni E, Nigri G. A new minimally invasive technique for the repair of diastasis recti: a pilot study. <i>Surg Endosc.</i> 2021;35(7):4028-34. Available from: https://doi.org/10.1007/s00464-021-08393-2 .	Studiedesign
Masurkar AA. Laparoscopic Trans-Abdominal Retromuscular (TARM) Repair for Ventral Hernia: A Novel, Low-Cost Technique for Sublay and Posterior Component Separation. <i>World J Surg.</i> 2020;44(4):1081-5. Available from: https://doi.org/10.1007/s00268-019-05298-z .	Population
Matarasso A, Swift RW, Rankin M. Abdominoplasty and abdominal contour surgery: a national plastic surgery survey. <i>Plast Reconstr Surg.</i> 2006;117(6):1797-808. Available from: https://doi.org/10.1097/01.prs.0000209918.55752.f3 .	Studiedesign
Matei OA, Runkel N. A novel technique of midline mesh repair for umbilical hernia associated with diastasis recti. <i>Surg Technol Int.</i> 2014;24:183-7.	Population
Mayer HF, Loustau HD. The Suprapubic Dermoadipose Flap for Aesthetic Reshaping of the Postpregnancy Abdomen. <i>Aesthet Surg J.</i> 2018;38(6):635-43. Available from: https://doi.org/10.1093/asj/sjx175 .	Studiedesign
Mazzocchi M, Dessy LA, Di Ronza S, Iodice P, Saggini R, Scuderi N. A study of postural changes after abdominal rectus plication abdominoplasty. <i>Hernia.</i> 2014;18(4):473-80. Available from: https://doi.org/10.1007/s10029-012-1015-1 .	Studiedesign
McPhail I. Abdominal aortic aneurysm and diastasis recti. <i>Angiology.</i> 2008;59(6):736-9. Available from: https://doi.org/10.1177/0003319708319940 .	Population
Medina JP, Busnelli G, Cerutti RR, Pirchi DE. Subcutaneous endoscopic approach for treatment of diastasis recti, associated with other midline abdominal wall defects. <i>Rev Argent Cir.</i> 2019;111(1):E20-E6.	Studiedesign
Mendes Dde A, Nahas FX, Veiga DF, Mendes FV, Figueiras RG, Gomes HC, et al. Ultrasonography for measuring rectus abdominis muscles diastasis. <i>Acta Cir Bras.</i> 2007;22(3):182-6. Available from: https://doi.org/10.1590/s0102-86502007000300005 .	Studiedesign
Mesquita LA, Machado AV, Andrade AV. Fisioterapia para redução da diástase dos músculos retos abdominais no pós-parto. <i>Revista Brasileira de Ginecologia e Obstetrícia.</i> 1999;21(5):267-72. Available from: https://doi.org/10.1590/s0100-72031999000500004 .	Språk
Mestak O, Kullac R, Mestak J, Nosek A, Krajcova A, Sukop A. Evaluation of the long-term stability of sheath plication using absorbable sutures in 51 patients with diastasis of the recti muscles: an ultrasonographic study. <i>Plast Reconstr Surg.</i> 2012;130(5):714e-9e. Available from: https://doi.org/10.1097/PRS.0b013e318267d806 .	Studiedesign
Misra MC, Bansal VK, Kulkarni MP, Pawar DK. Comparison of laparoscopic and open repair of incisional and primary ventral hernia: results of a prospective randomized study. <i>Surg Endosc.</i> 2006;20(12):1839-45. Available from: https://doi.org/10.1007/s00464-006-0118-0 .	Population
Mitura K, Skolimowska-Rzewuska M, Rzewuska A, Wyrzykowska D. Is mesh always necessary in every small umbilical hernia repair?	Population

Referens	Exklusionsorsak
Comparison of standardized primary sutured versus patch repair: retrospective cohort study. <i>Hernia</i> . 2021;25(3):571-7. Available from: https://doi.org/10.1007/s10029-020-02170-1 .	
Moga D, Buia F, Oprea V. Laparo-Endoscopic Repair of Ventral Hernia and Rectus Diastasis. <i>JSLS</i> . 2021;25(2). Available from: https://doi.org/10.4293/JSLS.2020.00103 .	Population
Montori G, Cocolini F, Manfredi R, Ceresoli M, Campanati L, Magnone S, et al. One year experience of swine dermal non-crosslinked collagen prostheses for abdominal wall repairs in elective and emergency surgery. <i>World J Emerg Surg</i> . 2015;10:28. Available from: https://doi.org/10.1186/s13017-015-0023-8 .	Population
Mossaad BM, Frame JD. Medial advancement of infraumbilical Scarpa's fascia improves waistline definition in "Brazilian" abdominoplasty. <i>Aesthetic Plast Surg</i> . 2013;37(1):3-10. Available from: https://doi.org/10.1007/s00266-012-0004-1 .	Intervention
Mota P, Pascoal AG, Carita AI, Bo K. The Immediate Effects on Inter-rectus Distance of Abdominal Crunch and Drawing-in Exercises During Pregnancy and the Postpartum Period. <i>J Orthop Sports Phys Ther</i> . 2015;45(10):781-8. Available from: https://doi.org/10.2519/jospt.2015.5459 .	Studiedesign
Munhoz AM, Sturtz G, Montag E, Arruda EG, Aldrichi C, Gemperli R, et al. Clinical outcome of abdominal wall after DIEP flap harvesting and immediate application of abdominoplasty techniques. <i>Plast Reconstr Surg</i> . 2005;116(7):1881-93. Available from: https://doi.org/10.1097/01.prs.0000191186.20698.0d .	Studiedesign
Nahabedian MY. Diastasis recti repair with onlay mesh. <i>Hernia</i> . 2021;25(4):855-62. Available from: https://doi.org/10.1007/s10029-021-02464-y .	Studiedesign
Nahabedian MY, Nahabedian AG. Closing the gap for patients with rectus abdominis diastasis. <i>Nursing</i> . 2018;48(1):49-52. Available from: https://doi.org/10.1097/01.NURSE.0000527601.09592.14 .	Publikationstyp
Nahas FX. An aesthetic classification of the abdomen based on the myoaponeurotic layer. <i>Plast Reconstr Surg</i> . 2001;108(6):1787-95; discussion 96-7. Available from: https://doi.org/10.1097/00006534-200111000-00057 .	Studiedesign
Nahas FX, Augusto SM, Ghelfond C. Should diastasis recti be corrected? <i>Aesthetic Plast Surg</i> . 1997;21(4):285-9. Available from: https://doi.org/10.1007/s002669900127 .	Studiedesign
Nahas FX, Ferreira LM, Augusto SM, Ghelfond C. Long-term follow-up of correction of rectus diastasis. <i>Plast Reconstr Surg</i> . 2005;115(6):1736-41; discussion 42-3. Available from: https://doi.org/10.1097/01.prs.0000161675.55337.f1 .	Studiedesign
Nahas FX, Ferreira LM, Ely PB, Ghelfond C. Rectus diastasis corrected with absorbable suture: a long-term evaluation. <i>Aesthetic Plast Surg</i> . 2011;35(1):43-8. Available from: https://doi.org/10.1007/s00266-010-9554-2 .	Studiedesign

Referens	Exklusionsorsak
Nahas FX, Ferreira LM, Mendes Jde A. An efficient way to correct recurrent rectus diastasis. <i>Aesthetic Plast Surg.</i> 2004;28(4):189-96. Available from: https://doi.org/10.1007/s00266-003-0097-7 .	Studiedesign
Nardi WS, Busnelli GL, Tchercansky A, Pirchi DE, Medina PJ. Diastasis recti associated with midline hernias: Totally subcutaneous video-endoscopic repair. <i>J Minim Access Surg.</i> 2018;14(2):161-3. Available from: https://doi.org/10.4103/jmas.JMAS_103_17 .	Studiedesign
Neely D. Recovery for the Postpartum Singer: Yoga and Yoga-Inspired Exercises for Regaining Abdominal Strength. <i>Journal of Singing.</i> 2020;77(2):191-200.	Publikationstyp
Neinstein RM, Matarasso A, Abramson DL. Concomitant abdominoplasty and umbilical hernia repair using the Ventralex hernia patch. <i>Plast Reconstr Surg.</i> 2015;135(4):1021-5. Available from: https://doi.org/10.1097/PRS.0000000000001135 .	Intervention
Nishihara Y, Asami M, Shimada T, Kawaguchi Y, Omoto K. Comorbid rectus abdominis diastasis is a risk factor for recurrence of umbilical hernia in Japanese patients. <i>Asian J Endosc Surg.</i> 2021;14(3):368-72. Available from: https://doi.org/10.1111/ases.12868 .	Population
Novitsky YW, Elliott HL, Orenstein SB, Rosen MJ. Transversus abdominis muscle release: a novel approach to posterior component separation during complex abdominal wall reconstruction. <i>Am J Surg.</i> 2012;204(5):709-16. Available from: https://doi.org/10.1016/j.amjsurg.2012.02.008 .	Population
O'Sullivan PB, Twomey L, Allison GT. Altered abdominal muscle recruitment in patients with chronic back pain following a specific exercise intervention. <i>J Orthop Sports Phys Ther.</i> 1998;27(2):114-24. Available from: https://doi.org/10.2519/jospt.1998.27.2.114 .	Population
Olsson A, Kiwanuka O, Wilhelmsson S, Sandblom G, Stackelberg O. Cohort study of the effect of surgical repair of symptomatic diastasis recti abdominis on abdominal trunk function and quality of life. <i>BJS Open.</i> 2019;3(6):750-8. Available from: https://doi.org/10.1002/bjs5.50213 .	Studiedesign
Palanivelu C, Rangarajan M, Jategaonkar PA, Amar V, Gokul KS, Srikanth B. Laparoscopic repair of diastasis recti using the 'Venetian blinds' technique of plication with prosthetic reinforcement: a retrospective study. <i>Hernia.</i> 2009;13(3):287-92. Available from: https://doi.org/10.1007/s10029-008-0464-z .	Studiedesign
Panait L, Bell RL, Roberts KE, Duffy AJ. Closing the gap: medialization of fascia with laparoscopic incisional hernia repair. <i>Hernia.</i> 2013;17(5):597-601. Available from: https://doi.org/10.1007/s10029-013-1133-4 .	Population
Parker MA, Millar AL, Dugan SA. Diastasis rectus abdominis and lumbo-pelvic pain and dysfunction -- are they related? <i>Journal of Women's Health Physical Therapy.</i> 2009;33(2):15-22.	Studiedesign
Pascoal AG, Dionisio S, Cordeiro F, Mota P. Inter-rectus distance in postpartum women can be reduced by isometric contraction of the abdominal muscles: a preliminary case-control study. <i>Physiotherapy.</i> 2014;100(4):344-8. Available from: https://doi.org/10.1016/j.physio.2013.11.006 .	Population

Referens	Exklusionsorsak
Pawar PA, Yeole UL, Navale M, Patil K. Effect of kinesiotaping on diastasis recti in post-partum women. Indian Journal of Public Health Research and Development. 2020;11(6):689-94.	Population
Pearson DG, Madura JA, 2nd. Modified retrorectus ventral hernia repair. Am J Surg. 2016;211(3):615-8. Available from: https://doi.org/10.1016/j.amjsurg.2015.11.012 .	Population
Pechter EA. Instant identification of redundant tissue in abdominoplasty with a marking grid. Aesthet Surg J. 2010;30(4):571-8. Available from: https://doi.org/10.1177/1090820X10377148 .	Studiedesign
Pereira N, Sciaraffia C, Danilla S, Parada F, Asfora C, Moral C. Effects of Abdominoplasty on Intra-Abdominal Pressure and Pulmonary Function. Aesthet Surg J. 2016;36(6):697-702. Available from: https://doi.org/10.1093/asj/sjv273 .	Studiedesign
Perry DJ, Driscoll DN. Corset Autodermis External Obliqueplasty Reinforcement of Rectus Diastasis and Umbilical Hernia Repairs during Abdominoplasty. Plast Reconstr Surg. 2021;147(4):860-3. Available from: https://doi.org/10.1097/PRS.0000000000007768 .	Publikationstyp
Persichetti P, Simone P, Scuderi N. Anchor-line abdominoplasty: a comprehensive approach to abdominal wall reconstruction and body contouring. Plast Reconstr Surg. 2005;116(1):289-94. Available from: https://doi.org/10.1097/01.prs.0000170049.94288.d2 .	Studiedesign
Pollock H, Pollock T. Progressive tension sutures: a technique to reduce local complications in abdominoplasty. Plast Reconstr Surg. 2000;105(7):2583-6; discussion 7-8. Available from: https://doi.org/10.1097/00006534-200006000-00047 .	Studiedesign
Povedano A, D'Alessandro GS, dos Santos LKIL, Munhoz AM, Gemperli R, Sampaio Góes JC. Subdermal neo-umbilicoplasty in abdominoplasty. European Journal of Plastic Surgery. 2017;41(3):335-44. Available from: https://doi.org/10.1007/s00238-017-1374-z .	Studiedesign
Prado A, Andrades P, Benitez S. Abdominoplasty: The Use of Polypropylene Mesh to Correct Myoaponeurotic-Layer Deformity. Aesthetic Plastic Surgery. 2004;28(3):144-7. Available from: https://doi.org/10.1007/s00266-004-3124-4 .	Studiedesign
Privett BJ, Ghusn M. Proposed technique for open repair of a small umbilical hernia and rectus divarication with self-gripping mesh. Hernia. 2016;20(4):527-30. Available from: https://doi.org/10.1007/s10029-016-1470-1 .	Population
Quilichini J, Leyder P. Simultaneous Prosthetic Mesh Abdominal Wall Reconstruction with Abdominoplasty for Ventral Hernia and Severe Rectus Diastasis Repairs. Plast Reconstr Surg. 2016;137(1):252e-4e. Available from: https://doi.org/10.1097/PRS.0000000000001932 .	Publikationstyp
Ramirez OM. Abdominoplasty and abdominal wall rehabilitation: a comprehensive approach. Plast Reconstr Surg. 2000;105(1):425-35. Available from: https://doi.org/10.1097/00006534-200001000-00069 .	Studiedesign
Ranney B. Diastasis recti and umbilical hernia causes, recognition and repair. S D J Med. 1990;43(10):5-8.	Studiedesign

Referens	Exklusionsorsak
Reinpold W, Schroder M, Berger C, Stoltenberg W, Kockerling F. MILOS and EMILOS repair of primary umbilical and epigastric hernias. Hernia. 2019;23(5):935-44. Available from: https://doi.org/10.1007/s10029-019-02056-x .	Population
Ribeiro L, Accorsi AJ, Buss A. Midabdominoplasty: indications and technique. Aesthetic Plast Surg. 1998;22(5):313-7. Available from: https://doi.org/10.1007/s002669900209 .	Publikationstyp
Rodrigues MA, Nahas FX, Gomes HC, Ferreira LM. Ventilatory function and intra-abdominal pressure in patients who underwent abdominoplasty with plication of the external oblique aponeurosis. Aesthetic Plast Surg. 2013;37(5):993-9. Available from: https://doi.org/10.1007/s00266-013-0158-5 .	Studiedesign
Rodrigues MA, Nahas FX, Reis RP, Ferreira LM. Does Diastasis Width Influence the Variation of the Intra-Abdominal Pressure After Correction of Rectus Diastasis? Aesthet Surg J. 2015;35(5):583-8. Available from: https://doi.org/10.1093/asj/sju091 .	Studiedesign
Romero-Morales C, Martin-Llantino PJ, Calvo-Lobo C, Beltran-Alacreu H, Lopez-Lopez D, Sanchez-Gomez R, et al. Effectiveness of Eccentric Exercise and a Vibration or Cryotherapy Program in Enhancing Rectus Abdominis Muscle Thickness and Inter-Rectus Distance in Patients with Chronic Mid-Portion Achilles Tendinopathy: A Randomized Clinical Trial. Int J Med Sci. 2018;15(14):1764-70. Available from: https://doi.org/10.7150/ijms.28656 .	Population
Rosen A, Hartman T. Repair of the midline fascial defect in abdominoplasty with long-acting barbed and smooth absorbable sutures. Aesthet Surg J. 2011;31(6):668-73. Available from: https://doi.org/10.1177/1090820X11415242 .	Studiedesign
Rozen WM, Kapila S, Enajat M, Tan MJ, Whitaker IS, Acosta R. Autologous dermal grafts for rectus sheath reconstruction and application in closure of ventral myofascial defects. Updates Surg. 2012;64(3):203-10. Available from: https://doi.org/10.1007/s13304-012-0167-y .	Population
Rutkowska-Kucharska A, Szpala A. The use of electromyography and magnetic resonance imaging to evaluate a core strengthening exercise programme. J Back Musculoskelet Rehabil. 2018;31(2):355-62. Available from: https://doi.org/10.3233/BMR-169780 .	Population
Saltz R. Two position comprehensive approach to abdominoplasty. Clin Plast Surg. 2014;41(4):681-704. Available from: https://doi.org/10.1016/j.cps.2014.07.008 .	Publikationstyp
Sancho MF, Pascoal AG, Mota P, Bo K. Abdominal exercises affect inter-rectus distance in postpartum women: a two-dimensional ultrasound study. Physiotherapy. 2015;101(3):286-91. Available from: https://doi.org/10.1016/j.physio.2015.04.004 .	Studiedesign
Sarlat Ribas MÁ. Diastasis of the recti. FMC Formacion Medica Continuada en Atencion Primaria. 2014;21(7):301-7.	Språk
Schwarz J, Reinpold W, Bittner R. Endoscopic mini/less open sublay technique (EMILOS)-a new technique for ventral hernia repair.	Population

Referens	Exklusionsorsak
Langenbecks Arch Surg. 2017;402(1):173-80. Available from: https://doi.org/10.1007/s00423-016-1522-0 .	
Senghaas A, Kremer T, Schmidt VJ, Harhaus L, Hirche C, Kneser U, et al. Sliding free transverse rectus abdominis myocutaneous flap for closure of a massive abdominal wall defect: A case report. Microsurgery. 2019;39(2):174-7. Available from: https://doi.org/10.1002/micr.30309 .	Studiedesign
Serra-Renom JM, Martinez-Teixido L, Serra-Mestre JM. Abdominoplasty with Customized Transverse Musculoaponeurotic Plications. Plast Reconstr Surg. 2015;136(6):741e-9e. Available from: https://doi.org/10.1097/PRS.0000000000001767 .	Population
Shah RH, Sharma A, Khullar R, Soni V, Baijal M, Chowbey PK. Laparoscopic repair of incarcerated ventral abdominal wall hernias. Hernia. 2008;12(5):457-63. Available from: https://doi.org/10.1007/s10029-008-0374-0 .	Population
Sharma A, Dey A, Khullar R, Soni V, Baijal M, Chowbey PK. Laparoscopic repair of suprapubic hernias: transabdominal partial extraperitoneal (TAPE) technique. Surg Endosc. 2011;25(7):2147-52. Available from: https://doi.org/10.1007/s00464-010-1513-0 .	Population
Shaw RB, Jr., Curet MJ, Kahn DM. Laparoscopic repair for recurrent abdominal wall hernia after TRAM flap breast reconstruction: case report of 2 patients. Ann Plast Surg. 2006;56(4):447-50. Available from: https://doi.org/10.1097/01.sap.0000200281.24169.1f .	Studiedesign
Sheppard S. Case study. The role of transversus abdominus in post partum correction of gross divarication recti. Manual Therapy. 1996;1(4):214-6.	Studiedesign
Sheppard S. Management of postpartum gross divarication recti... reprinted with permission. Copyright Journal of Association of Chartered Physiotherapists in Women's Health, 1996. MIDIRS Midwifery Digest. 1997;7(1):84-6.	Studiedesign
Shipkov HD, Mojallal A, Braye F. Simultaneous Abdominoplasty and Umbilical Hernia Repair via Laparoscopy: a Preliminary Report. Folia Med (Plovdiv). 2017;59(2):222-7. Available from: https://doi.org/10.1515/folmed-2017-0026 .	Population
Shirah BH, Shirah HA. The Effectiveness of Polypropylene Mesh in the Open and LaparoscopicRepair of Divarication of the Recti. J Med Imp Surg. 2016;1(1):105. Available from: https://doi.org/10.4172/jims.1000105 .	Studiedesign
Shubinets V, Fox JP, Sarik JR, Kovach SJ, Fischer JP. Surgically Treated Hernia following Abdominally Based Autologous Breast Reconstruction: Prevalence, Outcomes, and Expenditures. Plast Reconstr Surg. 2016;137(3):749-57. Available from: https://doi.org/10.1097/01.pr.s.0000479931.96538.c5 .	Population
Sickle KR, Baghai M, Mattar SG, Bowers SP, Ramaswamy A, Swafford V, et al. What happens to the rectus abdominus fascia after laparoscopic ventral hernia repair? Hernia. 2005;9(4):358-62. Available from: https://doi.org/10.1007/s10029-005-0018-6 .	Studiedesign

Referens	Exklusionsorsak
Siddiky AH, Kapadia CR. Laparoscopic plication of the linea alba as a repair for diastasis recti - a mesh free approach. <i>J Surg Case Rep.</i> 2010;2010(5):3. Available from: https://doi.org/10.1093/jscr/2010.5.3 .	Studiedesign
Sood R, Janes LE, Shah N, Sasson DC, Ellis MF, Dumanian GA. Mesh Repair of Rectus Diastasis for Abdominoplasty is Safer than Suture Plication. <i>Plast Reconstr Surg Glob Open.</i> 2021;9(8):e3721. Available from: https://doi.org/10.1097/GOX.00000000000003721 .	Studiedesign
Soybel DI. Abdominal rectus diastasis: Questions of wellness and biology. <i>Surgery.</i> 2016;160(5):1376-8. Available from: https://doi.org/10.1016/j.surg.2016.06.021 .	Publikationstyp
Staalesen T, Olsen MF, Elander A. The Effect of Abdominoplasty and Outcome of Rectus Fascia Plication on Health-Related Quality of Life in Post-Bariatric Surgery Patients. <i>Plast Reconstr Surg.</i> 2015;136(6):750e-61e. Available from: https://doi.org/10.1097/PRS.0000000000001772 .	Population
Stark B, Emanuelsson P, Gunnarsson U, Strigard K. Validation of Biomed system 4 for measuring the strength of muscles in patients with rectus diastasis. <i>J Plast Surg Hand Surg.</i> 2012;46(2):102-5. Available from: https://doi.org/10.3109/2000656X.2011.644707 .	Studiedesign
Stedge HL, Medina McKeon JM. A Supervised Pelvic Floor Training Program Has Minimal Effect on the Prevalence of Postpartum Diastasis Recti Abdominis: A Critically Appraised Paper. <i>International Journal of Athletic Therapy and Training.</i> 2020;25(6):299-302. Available from: https://doi.org/10.1123/ijatt.2019-0112 .	Publikationstyp
Strigard K, Clay L, Stark B, Gunnarsson U. Predictive Factors in the Outcome of Surgical Repair of Abdominal Rectus Diastasis. <i>Plast Reconstr Surg Glob Open.</i> 2016;4(5):e702. Available from: https://doi.org/10.1097/GOX.0000000000000688 .	Studiedesign
Swiatkiewicz J, Tomaszewski D, Kabala P. 25th International Congress of the European Association for Endoscopic Surgery (EAES) Frankfurt, Germany, 14-17 June 2017 Video Presentations. <i>Surg Endosc.</i> 2017;31(Suppl 2):190-259. Available from: https://doi.org/10.1007/s00464-017-5542-9 .	Publikationstyp
Tadiparthi S, Shokrollahi K, Doyle GS, Fahmy FS. Rectus sheath plication in abdominoplasty: assessment of its longevity and a review of the literature. <i>J Plast Reconstr Aesthet Surg.</i> 2012;65(3):328-32. Available from: https://doi.org/10.1016/j.bjps.2011.09.024 .	Studiedesign
Takaya K, Aramaki-Hattori N, Yabuki H, Wada N, Sakai S, Okabe K, et al. Correction of Diastasis Rectus Abdominis with Tacking the Rectus Sheath and Resection of Excess Skin for Cosmesis. <i>Case Rep Med.</i> 2020;2020:7635801. Available from: https://doi.org/10.1155/2020/7635801 .	Studiedesign
Takeuchi Y, Kurashima Y, Nakanishi Y, Asano T, Noji T, Ebihara Y, et al. Mesh trimming and suture reconstruction for wound dehiscence after huge abdominal intercostal hernia repair: A case report. <i>Int J Surg Case Rep.</i> 2018;53:381-5. Available from: https://doi.org/10.1016/j.ijscr.2018.11.028 .	Studiedesign

Referens	Exklusionsorsak
Talsman R, Kaplan B, Haik J, Aronov S, Shraga A, Orenstein A. Measuring alterations in intra-abdominal pressure during abdominoplasty as a predictive value for possible postoperative complications. <i>Aesthetic Plast Surg.</i> 2002;26(3):189-92. Available from: https://doi.org/10.1007/s00266-001-1469-5 .	Studiedesign
Tan W, McAllister J, Feaman S, Frisella M, Blatnik J, Kahan L. 18th Annual Hernia Repair. <i>Hernia.</i> 2017;21(Suppl 1):1-137. Available from: https://doi.org/10.1007/s10029-017-1571-5 .	Publikationstyp
Taylor DA, Merten SL, Sandercoe GD, Gahankari D, Ingram SB, Moncrieff NJ, et al. Abdominoplasty Improves Low Back Pain and Urinary Incontinence. <i>Plast Reconstr Surg.</i> 2018;141(3):637-45. Available from: https://doi.org/10.1097/PRS.0000000000004100 .	Studiedesign
Temel M, Turkmen A, Berberoglu O. Improvements in Vertebral-Column Angles and Psychological Metrics After Abdominoplasty With Rectus Plication. <i>Aesthet Surg J.</i> 2016;36(5):577-87. Available from: https://doi.org/10.1093/asj/sjv257 .	Studiedesign
Tenzel PL, Johnson RG, Bilezikian JA, Powers WF, Hope WW. A Preliminary Assessment of Abdominal Wall Tension in Patients Undergoing Retromuscular Hernia Repair. <i>Surg Technol Int.</i> 2019;34:251-4.	Population
Thornton SL, Thornton SJ. Management of Gross Divarication of the Recti Abdominis in Pregnancy and Labour. <i>Physiotherapy.</i> 1993;79(7):457-8. Available from: https://doi.org/10.1016/s0031-9406(10)60221-0 .	Studiedesign
Tiffert-Tłok J, Skrzypulec-Plinta V, Szuścik-Niewiadomy K, Drosdzol-Cop A, Czajkowska M. Pro-health behaviors of women in given trimesters of pregnancy - A preliminary study. <i>Ginekologia i Poloznictwo.</i> 2019;15(3):22-7.	Studiedesign
Toronto IR. The relief of low back pain with the WARP abdominoplasty: a preliminary report. <i>Plast Reconstr Surg.</i> 1990;85(4):545-55. Available from: https://doi.org/10.1097/00006534-199004000-00009 .	Population
Torregrosa Gallud A, Garcia Pastor P, Jimenez Rosellon R, Muniesa Gallardo C, Bueno Lledó J, Bonafe Diana S, et al. Oral and video presentations. <i>Hernia.</i> 2017;21(Suppl 2):139-273. Available from: https://doi.org/10.1007/s10029-017-1595-x .	Publikationstyp
Tuominen R, Vironen J, Jahkola T. Case series of a novel open plication supported by mesh (PSUM) - technique for symptomatic abdominal rectus diastasis repair with or without concomitant midline hernia: Early results and a review of the literature. <i>International Journal of Abdominal Wall and Hernia Surgery.</i> 2019;2(4):142-8. Available from: https://doi.org/10.4103/ijawhs.ijawhs_25_19 .	Studiedesign
Tuveri M, Tuveri A, Nicolo E. Repair of large abdominal incisional hernia by reconstructing the midline and use of an onlay of biological material. <i>Am J Surg.</i> 2011;202(1):e7-11. Available from: https://doi.org/10.1016/j.amjsurg.2010.06.005 .	Studiedesign
Uzair S, Babar Z, Sutton PA. Tissue regeneration mesh reinforcement during abdominoplasty for severe myoaponeurotic laxity after	Studiedesign

Referens	Exklusionsorsak
pregnancy. BMJ Case Rep. 2013;2013:05. Available from: https://doi.org/10.1136/bcr-2013-010306 .	
Vaishnavi G, Mohan Kumar G, Jayson CJ, Kirupa K, Tharani G, Kamatchi K, et al. Effectiveness of exercise in treating rectus abdominis diastasis. Biomedicine (India). 2018;38(4):583-7.	Population
Van Cleven S, Muysoms F. 25th International Congress of the European Association for Endoscopic Surgery (EAES) Frankfurt, Germany, 14-17 June 2017 Video Presentations. Surg Endosc. 2017;31(Suppl 2):190-259. Available from: https://doi.org/10.1007/s00464-017-5542-9 .	Publikationstyp
van Schalkwyk CP, Dusseldorf JR, Liang DG, Keshava A, Gilmore AJ, Merten S. Concomitant Abdominoplasty and Laparoscopic Umbilical Hernia Repair. Aesthet Surg J. 2018;38(12):NP196-NP204. Available from: https://doi.org/10.1093/asj/sjy100 .	Studiedesign
van Uchelen JH, Kon M, Werker PM. The long-term durability of plication of the anterior rectus sheath assessed by ultrasonography. Plast Reconstr Surg. 2001;107(6):1578-84. Available from: https://doi.org/10.1097/00006534-200105000-00046 .	Studiedesign
Vera Cucchiaro J, Lostia H, Velazquez P, Liska E. Liposubabdominoplasty with Progressive Traction Sutures. Plast Reconstr Surg Glob Open. 2017;5(6):e1338. Available from: https://doi.org/10.1097/GOX.0000000000001338 .	Studiedesign
Verissimo P, Nahas FX, Barbosa MV, de Carvalho Gomes HF, Ferreira LM. Is it possible to repair diastasis recti and shorten the aponeurosis at the same time? Aesthetic Plast Surg. 2014;38(2):379-86. Available from: https://doi.org/10.1007/s00266-014-0272-z .	Utfall
Walton LM, Costa A, LaVanture D, McIlrath S, Stebbins B. The effects of a 6 week dynamic core stability plank exercise program compared to a traditional supine core stability strengthening program on diastasis recti abdominis closure, pain, Oswestry disability index (ODI) and pelvic floor disability index scores (PFDI). Physical Therapy and Rehabilitation. 2016;3(1):3. Available from: https://doi.org/10.7243/2055-2386-3-3 .	Population
Wang Q, Yu X, Chen G, Sun X, Wang J. Does diastasis recti abdominis weaken pelvic floor function? A cross-sectional study. Int Urogynecol J. 2020;31(2):277-83. Available from: https://doi.org/10.1007/s00192-019-04005-9 .	Studiedesign
Wang TY, Elliott R, Low DW. Damage control abdomen: single-stage reconstruction using a vicryl mesh buttress. Ann Plast Surg. 2013;70(3):324-30. Available from: https://doi.org/10.1097/SAP.0b013e3182361af9 .	Population
Whitson BA, Ose KJ. Spontaneous posterior rectus sheath hernia: a new clinical entity? Hernia. 2007;11(5):445-7. Available from: https://doi.org/10.1007/s10029-007-0204-9 .	Studiedesign
Whittaker JL, Emery CA. Impact of the FIFA 11+ on the structure of select muscles in adolescent female soccer players. Phys Ther Sport. 2015;16(3):228-35. Available from: https://doi.org/10.1016/j.ptsp.2014.10.007 .	Population

Referens	Exklusionsorsak
Wiessner R, Vorwerk T, Tolla-Jensen C, Gehring A. Continuous Laparoscopic Closure of the Linea Alba with Barbed Sutures Combined with Laparoscopic Mesh Implantation (IPOM Plus Repair) As a New Technique for Treatment of Abdominal Hernias. <i>Front Surg.</i> 2017;4:62. Available from: https://doi.org/10.3389/fsurg.2017.00062 .	Population
Wilhelmsson S, Fagevik Olsen M, Staalesen T, Elander A, Nygren-Bonnier M. Abdominal plasty with and without plication-effects on trunk muscles, lung function, and self-rated physical function. <i>J Plast Surg Hand Surg.</i> 2017;51(3):199-204. Available from: https://doi.org/10.1080/2000656X.2016.1222293 .	Population
Williams TC, Hardaway M, Altuna B. Ambulatory abdominoplasty tailored to patients with an appropriate body mass index. <i>Aesthet Surg J.</i> 2005;25(2):132-7. Available from: https://doi.org/10.1016/j.asj.2005.01.011 .	Studiedesign
Winder J, Pauli E, Shope A. 18th Annual Hernia Repair. <i>Hernia.</i> 2017;21(Suppl 1):1-137. Available from: https://doi.org/10.1007/s10029-017-1571-5 .	Publikationstyp
Wong CH, Song C. The "intramuscular" rectus abdominis pedicle. <i>Plast Reconstr Surg.</i> 2007;119(6):1964-5. Available from: https://doi.org/10.1097/01.prs.0000259896.22578.29 .	Studiedesign
Wu L, Li J, Miao R. Intraoperative Observation during Total Extraperitoneal Repair (TEP). <i>Am Surg.</i> 2020;86(2):146-51.	Population
Yousif NJ, Lifchez SD, Nguyen HH. Transverse rectus sheath plication in abdominoplasty. <i>Plast Reconstr Surg.</i> 2004;114(3):778-84. Available from: https://doi.org/10.1097/01.prs.0000131023.09405.a8 .	Studiedesign
Zappile-Lucis M. Quality of life measurements and physical therapy management of a female diagnosed with Diastasis Recti Abdominis. <i>Journal of Women's Health Physical Therapy.</i> 2009;33(1):22-.	Publikationstyp
Zukowski ML, Ash K, Spencer D, Malanoski M, Moore G. Endoscopic intracorporeal abdominoplasty: a review of 85 cases. <i>Plast Reconstr Surg.</i> 1998;102(2):516-27. Available from: https://doi.org/10.1097/00006534-199808000-00037 .	Studiedesign