

Bilaga till rapport

1 (3)

Kontinuitet i vården / Continuity of care,
rapport 329 (2021)

Bilaga 8 Sammanställning av Risk of Bias-bedömning för psykisk sjukdom / Appendix 8 Templates for assessment of Risk of Bias regarding severe mental illness

Studie	1. Confounding	2. Exponering	3. Bortfall	4. Mätning och analys	5. Rapporterings	6. Jäv	Sammanvägd RoB
Adair et al.* [1]	Måttlig	Låg	Måttlig	Måttlig	Måttlig	Låg	Hög
Adanes et al. [2]	Måttlig	Låg	Måttlig	Låg	Låg	Låg	Måttlig
Bindman et al. [3]	Måttlig	Låg	Måttlig	Måttlig	Låg	Låg	Hög
Catty et al. [4]	Hög	Måttlig	Hög	Måttlig	Måttlig	Låg	Hög
Chien et al. [5]	Måttlig	Låg	Låg	Måttlig	Låg	Låg	Måttlig
Conti et al. [6]	Måttlig	Låg	Låg	Låg	Låg	Låg	Måttlig
Desai et al. [7]	Måttlig	Måttlig	Låg	Låg	Låg	Låg	Måttlig
Farley et al. [8]	Måttlig	Låg	Måttlig	Låg	Låg	Låg	Måttlig
Fleury et al. [9]	Hög	Hög	Måttlig	Hög	Måttlig	Låg	Oacceptabel
Giacco et al. [10]	Måttlig	Måttlig	Låg	Låg	Låg	Låg	Måttlig
Green et al. [11]	Hög	Måttlig	Måttlig	Hög	Måttlig	Låg	Oacceptabel
Hoertel et al. [12]	Måttlig	Låg	Låg	Låg	Låg	Låg	Måttlig
Kaltsidis et al. [13]	Måttlig	Måttlig	Låg	Låg	Låg	Låg	Måttlig
Mitton et al.* [14]	Måttlig	Låg	Låg	Låg	Låg	Låg	Måttlig
Puntis et al. [15]	Måttlig	Låg	Låg	Låg	Låg	Låg	Måttlig
Ride et al. [16]	Måttlig	Låg	Måttlig	Låg	Låg	Låg	Måttlig
Van der Lee et al. [17]	Hög	Låg	Måttlig	Låg	Låg	Låg	Hög
Watkins et al. [18]	Måttlig	Låg	Låg	Låg	Låg	Låg	Måttlig

*Artiklarna utgår från samma studie och rapporterar olika utfallsmått.

Referenslista:

1. Adair CE, McDougall GM, Mitton CR, Joyce AS, Wild TC, Gordon A, et al. Continuity of care and health outcomes among persons with severe mental illness. *Psychiatric Services*. 2005;56(9):1061-9. Available from: <https://doi.org/10.1176/appi.ps.56.9.1061>.
2. Adnanes M, Kalseth J, Ose SO, Ruud T, Rugkasa J, Puntis S. Quality of life and service satisfaction in outpatients with severe or non-severe mental illness diagnoses. *Quality of Life Research*. 2019;28(3):713-24. Available from: <https://doi.org/https://dx.doi.org/10.1007/s11136-018-2039-5>.
3. Bindman J, Johnson S, Szmukler G, Wright S, Kuipers E, Thornicroft G, et al. Continuity of care clinical outcome: A prospective cohort study. *Social Psychiatry and Psychiatric Epidemiology*. 2000;35(6):242-7. Available from: <https://doi.org/10.1007/s001270050234>.
4. Catty J, White S, Clement S, Cowan N, Geyer C, Harvey K, et al. Continuity of care for people with psychotic illness: its relationship to clinical and social functioning. *International Journal of Social Psychiatry*. 2013;59(1):5-17. Available from: <https://doi.org/https://dx.doi.org/10.1177/0020764011421440>.
5. Chien CF, Steinwachs DM, Lehman A, Fahey M, Skinner EA. Provider continuity and outcomes of care for persons with schizophrenia. *Mental Health Services Research*. 2000;2(4):201-11. Available from: <https://doi.org/10.1023/A:1010160419561>.
6. Conti V, Lora A, Cipriani A, Fortino I, Merlini L, Barbui C. Persistence with pharmacological treatment in the specialist mental healthcare of patients with severe mental disorders. *European Journal of Clinical Pharmacology*. 2012;68(12):1647-55. Available from: <https://doi.org/10.1007/s00228-012-1298-2>.
7. Desai RA, Dausey DJ, Rosenheck RA. Mental health service delivery and suicide risk: the role of individual patient and facility factors. *American Journal of Psychiatry*. 2005;162(2):311-8. Available from: <https://doi.org/10.1176/appi.ajp.162.2.311>.
8. Farley JF, Wang CC, Hansen RA, Voils CI, Maciejewski ML. Continuity of antipsychotic medication management for Medicaid patients with schizophrenia. *Psychiatric Services*. 2011;62(7):747-52. Available from: https://doi.org/https://dx.doi.org/10.1176/ps.62.7.pss6207_0747.
9. Fleury MJ, Grenier G, Bamvita JM, Caron J. Professional service utilisation among patients with severe mental disorders. *BMC health services research*. 2010;10:141. Available from: <https://doi.org/https://dx.doi.org/10.1186/1472-6963-10-141>.
10. Giacco D, Bird VJ, Ahmad T, Bauer M, Lasalvia A, Lorant V, et al. The same or different psychiatrists for in- and out-patient treatment? A multi-country natural experiment. *Epidemiology & Psychiatric Science*. 2018;29:e10. Available from: <https://doi.org/https://dx.doi.org/10.1017/S2045796018000732>.
11. Green CA, Polen MR, Janoff SL, Castleton DK, Wisdom JP, Vuckovic N, et al. Understanding how clinician-patient relationships and relational continuity of care affect recovery from serious mental illness: STARS study results. *Psychiatric Rehabilitation Journal*. 2008;32(1):9-22. Available from: <https://doi.org/https://dx.doi.org/10.2975/32.1.2008.9.22>.
12. Hoertel N, Limosin F, Leleu H. Poor longitudinal continuity of care is associated with an increased mortality rate among patients with mental disorders: results from the French

- National Health Insurance Reimbursement Database. European Psychiatry: the Journal of the Association of European Psychiatrists. 2014;29(6):358-64. Available from: <https://doi.org/https://dx.doi.org/10.1016/j.eurpsy.2013.12.001>.
- 13. Kaltsidis G, Bamvita JM, Grenier G, Fleury MJ. Predictors of Frequent Emergency Department Utilization for Mental Health Reasons. *Journal of Behavioral Health Services & Research*. 2020;17:17. Available from: <https://doi.org/https://dx.doi.org/10.1007/s11414-020-09695-4>.
 - 14. Mitton CR, Adair CE, McDougall GM, Marcoux G. Continuity of care and health care costs among persons with severe mental illness. *Psychiatric Services*. 2005;56(9):1070-6. Available from: <https://doi.org/10.1176/appi.ps.56.9.1070>.
 - 15. Puntis SR, Rugkasa J, Burns T. The association between continuity of care and readmission to hospital in patients with severe psychosis. *Social Psychiatry & Psychiatric Epidemiology*. 2016;51(12):1633-43. Available from: <https://doi.org/10.1007/s00127-016-1287-3>.
 - 16. Ride J, Kasteridis P, Gutacker N, Doran T, Rice N, Gravelle H, et al. Impact of family practice continuity of care on unplanned hospital use for people with serious mental illness. *Health Services Research*. 2019;54(6):1316-25. Available from: <https://doi.org/https://dx.doi.org/10.1111/1475-6773.13211>.
 - 17. van der Lee A, de Haan L, Beekman A. Schizophrenia in the Netherlands: Continuity of Care with Better Quality of Care for Less Medical Costs. *PLoS ONE [Electronic Resource]*. 2016;11(6):e0157150. Available from: <https://doi.org/https://dx.doi.org/10.1371/journal.pone.0157150>.
 - 18. Watkins KE, Paddock SM, Hudson TJ, Ounpraseuth S, Schrader AM, Hepner KA, et al. Association Between Quality Measures and Mortality in Individuals With Co-Occurring Mental Health and Substance Use Disorders. *Journal of Substance Abuse Treatment*. 2016;69:1-8. Available from: <https://doi.org/https://dx.doi.org/10.1016/j.jsat.2016.06.001>.