

## Bilaga 2 Exkluderade studier

För att en studie skulle inkluderas krävdes att följande kriterier uppfylldes.

- **Population.** Patienterna i studien måste vara 65 år eller äldre samt inkommit akut till sjukhus.
- **Intervention.** Studien måste utvärdera *strukturerat omhändertagande av äldre* på sjukhus där multiprofessionella team är delaktiga.
- **Jämförelsealternativ.** Jämförelsealternativet måste vara vanlig vård.
- **Effektmått.** Minst ett, av följande typer av effektmått, måste redovisas:  
(1) personlig funktionsförmåga (P-ADL), (2) instrumentell funktionsförmåga (I-ADL), (3) kognitiv funktion, (4) depression, (5) livskvalitet, (6) närlstående påverkan, (7) återinskrivning, (8) boende i eget hem efter sjukhusvistelsen eller (9) andel döda.
- **Studiedesign.** Studien måste vara en randomiserad kontrollerad studie (RCT). Observationsstudier (NRS) kan också accepteras under förutsättning att skillnader mellan interventions- och kontrollgrupp vid studiestarten hanterats på ett tillförlitligt sätt. Detta innebär att man måste ha använt multivariata metoder. Exempel på en till svensk metod är ”propensity scores”. Metoderna måste ha redovisats på ett sådant sätt i artiklarna att tillförlitligheten går att bedöma. Man måste även ha tillräckligt många variabler för att det ska vara möjligt att kontrollera för kända faktorer som kan snedvrida resultaten. Det måste dessutom finnas tillräckligt många individer för att problem med frihetsgrader inte ska uppstå då antalet variabler är många.

Ibland har artiklar exkluderats för att ett villkor inte varit uppfyllt och ibland handlar det om flera villkor. Oftast redovisas ett skäl även om fler är möjliga. Det förekommer även att artikeln inte redovisade forskningsresultat utan handlade om något annat, t ex en kommentar av en annan artikel eller att det rörde sig om en programbeskrivning. I dessa fall anges ”övrigt” som exkluderingsskäl. Efter genomläsning i fulltext exkluderades 420 studier.

Referenser	Exkluderings-skäl
Abatecola AM, Spazzafumo L, Corsonello A, Sirolla C, Bustacchini S, Guffanti E. Development and Validation of the HOPE Prognostic Index on 24-Month Posthospital Mortality and Rehospitalization: Italian National Research Center on Aging (INRCA). <i>Rejuvenation Res.</i> 2011.	Irrelevant studiedesign
Abizanda P, Leon M, Dominguez-Martin L, Lozano-Berrio V, Romero L, Luengo C, et al. Effects of a short-term occupational therapy intervention in an acute geriatric unit. A randomized clinical trial. <i>Maturitas.</i> 2011;69(3):273-8.	Irrelevant intervention
Abizanda P, Navarro JL, Romero L, Leon M, Sanchez-Jurado PM, Dominguez L. Upper extremity function, an independent predictor of adverse events in hospitalized elderly. <i>Gerontology.</i> 2007;53(5):267-73.	Irrelevant jämförelse-alternativ
ACE unit seeks to reduce elderly functional decline. <i>Hosp Case Manag.</i> 2002;10(5):71-3.	Ej levererad beställning
Adam SH, Eid HO, Barss P, Lunsjo K, Grivna M, Torab FC, et al. Epidemiology of geriatric trauma in United Arab Emirates. <i>Arch Gerontol Geriatr.</i> 2008;47(3):377-82.	Irrelevant intervention
Adamis D, Treloar A, Gregson N, Macdonald AJ, Martin FC. Delirium and the functional recovery of older medical inpatients after acute illness: the significance of biological factors. <i>Arch Gerontol Geriatr.</i> 2011;52(3):276-80.	Irrelevant intervention
Adams, A. L., Schiff, M. A., Koepsell, T. D., Rivara, F. P., Leroux, B. G., Becker, T. M., & Hedges, J. R. (2010). Physician consultation, multidisciplinary care, and 1-year mortality in Medicare recipients hospitalized with hip and lower extremity injuries. <i>J Am Geriatr Soc,</i> 58(10), 1835-1842. doi: 10.1111/j.1532-5415.2010.03087.x.	Irrelevant population
Adshead F, Cody DD, Pitt B. BASDEC: a novel screening instrument for depression in elderly medical inpatients. <i>BMJ.</i> 1992;305(6850):397.	Irrelevant intervention
Adunsky A, Arad M, Levi R, Blankstein A, Zeilig G, Mizrahi E. Five-year experience with the 'Sheba' model of comprehensive orthogeriatric care for elderly hip fracture patients. <i>Disability and Rehabilitation.</i> 2005;27(18-19):1123-7.	Irrelevant studiedesign
Adunsky A, Fleissig Y, Levenkrohn S, Arad M, Noy S. A comparative study of Mini-Mental Test, Clock Drawing task and Cognitive-FIM in evaluating functional outcome of elderly hip fracture patients. <i>Clin Rehabil.</i> 2002;16(4):414-9.	Irrelevant intervention
Adunsky A, Levi R, Cecic A, Arad M, Noy S, Barell V. The "Sheba" model of comprehensive orthogeriatric care for elderly hip fracture patients: A preliminary report. <i>Israel Medical Association Journal.</i> 2002;4(4):259-61.	Irrelevant studiedesign
Ahmed Nasiya, N., & Pearce Shannon, E. (2010). Acute care for the elderly: A literature review. <i>Population Health Management,</i> 13(4), 219-225.	Övrigt
Ahmed, N., Taylor, K., McDaniel, Y., & Dyer, C. B. (2012). The role of an acute care for the elderly unit in achieving hospital quality indicators while caring for frail hospitalized elders. <i>Population Health Management,</i> 15, 236-240.	Irrelevant studiedesign
Aimonino, N., Tibaldi, V., Barale, S., Bardelli, B., Pilon, S., Marchetto, C., . . . Molaschi, M. (2007). Depressive symptoms and quality of life in elderly patients with exacerbation of chronic obstructive pulmonary disease or cardiac heart failure: preliminary data of a randomized controlled trial. <i>Arch Gerontol Geriatr,</i> 44 Suppl 1, 7-12.	Irrelevant population
Aizen E, Swartzman R, Clarfield AM. Hospitalization of nursing home residents in an acute-care geriatric department: Direct versus emergency room admission. <i>Israel Medical Association Journal.</i> 2001;3(10):734-8.	Ej levererad beställning

Alarcon T, Barcena A, Gonzalez-Montalvo JI, Penalosa C, Salgado A. Factors predictive of outcome on admission to an acute geriatric ward. <i>Age and Ageing</i> . 1999;28(5):429-32.	Irrelevant studiedesign
Albiol, T., Martinez, A., Sanchez, R., J, L., Garcia, N., & J, A. (2011). Functional trajectories of elderly patients admitted in a geriatric rehabilitation unit after discharge of acute hospitalization. <i>European Geriatric Medicine</i> , 2, S49.	Irrelevant studiedesign
Anderson C, Deepak BV, Amoateng-Adjepong Y, Zarich S. Benefits of comprehensive inpatient education and discharge planning combined with outpatient support in elderly patients with congestive heart failure. <i>Congest Heart Fail</i> . 2005;11(6):315-21.	Irrelevant studiedesign
Andersson P, Hallberg IR, Renvert S. Comparison of oral health status on admission and at discharge in a group of geriatric rehabilitation patients. <i>Oral Health Prev Dent</i> . 2003;1(3):221-8.	Irrelevant studiedesign
Andrews, A., Sharp, S., McCullough, D., & Wennberg John, E. (2011). Fewer Hospitalizations Result When Primary Care Is Highly Integrated Into A Continuing Care Retirement Community. <i>Health Affairs</i> , 30(5), 975-984.	Irrelevant studiedesign
Andrews, B. C., Kaye, J., Bowcutt, M., & Campbell, J. (2001). Redesigning geriatric healthcare: how cross-functional teams and process improvement provide a competitive advantage. <i>Health Mark Q</i> . 19(2), 33-48.	Irrelevant studiedesign
Antonelli I, Landi F, Cipriani L, Bruno E, Pagano F, Gemma A, et al. Nutritional assessment: a primary component of multidimensional geriatric assessment in the acute care setting. <i>J Am Geriatr Soc</i> . 1996;44(2):166-74.	Irrelevant studiedesign
Applegate WB, Graney MJ, Miller ST, Elam JT. Impact of a geriatric assessment unit on subsequent health care charges. <i>Am J Public Health</i> . 1991;81(10):1302-6.	Irrelevant studiedesign
Arellano, P., Sanchez, R., M, D., Hernandez, L., Pineda, P., Sabartes, F, . . . M, A. (2011). Geriatric assessment as long-term mortality predictor. <i>European Geriatric Medicine</i> , 2, S36.	Irrelevant studiedesign
Armstrong J, Glenny C, Stolee P, Berg K. A comparison of two assessment systems in predicting functional outcomes of older rehabilitation patients. <i>Age Ageing</i> . 2010;39(3):394-9.	Irrelevant population
Arora, V. M., Plein, C., Chen, S., Siddique, J., Sachs, G. A., & Meltzer, D. O. (2009). Relationship between quality of care and functional decline in hospitalized vulnerable elders. <i>Medical Care</i> , 47(8), 895-901.	Irrelevant studiedesign
Asmus-Szepesi, K. J., de, V., P, L., Nieboer, A. P., van, W., J, D., . . . Mackenbach, J. P. (2011). Evaluation design of a reactivation care program to prevent functional loss in hospitalised elderly: A cohort study including a randomised controlled trial. <i>BMC Geriatr</i> , 11, 36.	Övrigt
Bachman SS, Collard AF, Greenberg JN, Fountain E, Huebner TW, Kimball B, et al. An innovative approach to geriatric acute care delivery: the Choate-Symmes experience. <i>Hosp Health Serv Adm</i> . 1987;32(4):509-20.	Övrigt
Baldwin, R., Pratt, H., Goring, H., Marriott, A., & Roberts, C. (2004). Does a nurse-led mental health liaison service for older people reduce psychiatric morbidity in acute general medical wards? A randomised controlled trial. <i>Age Ageing</i> , 33(5), 472-478.	Irrelevant intervention
Barer, D. (2011). Inpatient comprehensive geriatric assessment improves the likelihood of living at home at 12 months. <i>ACP Journal Club</i> , 155, 1.	Övrigt
Barker WH, Williams TF, Zimmer JG. Geriatric consultation teams in acute hospitals: Impact on back-up of elderly patients. <i>Journal of the American Geriatrics Society</i> . 1985;33(6):422-8.	Irrelevant studiedesign
Barone, A., Giusti, A., Pizzonia, M., Razzano, M., Palummo, E., & Pioli, G. (2006). A comprehensive geriatric intervention reduces short- and long-term mortality in older people with hip fracture. <i>Journal of the American Geriatrics Society</i> , 54(7), 1145-1147.	Irrelevant population

Barton C, Miller B, Yaffe K. Improved evaluation and management of cognitive impairment: results of a comprehensive intervention in long-term care. <i>Journal of the American Medical Directors Association</i> . 2006;7(2):84-9.	Irrelevant studiedesign
Bauer JM, Vogl T, Wicklein S, Trögner J, Mühlberg W, Sieber CC. Comparison of the Mini Nutritional Assessment, Subjective Global Assessment, and Nutritional Risk Screening (NRS 2002) for nutritional screening and assessment in geriatric hospital patients. <i>Zeitschrift für Gerontologie und Geriatrie</i> . 2005;38(5):322-7.	Irrelevanta effektmått
Baztan JJ, Galvez CP, Socorro A. Recovery of functional impairment after acute illness and mortality: one-year follow-up study. <i>Gerontology</i> . 2009;55(3):269-74.	Irrelevant studiedesign
Belknap SM, Gillespie U. A ward-based pharmacist intervention reduced emergency department visits and drug-related readmissions in elderly patients. <i>ACP Journal Club</i> . 2009;151(3):11-.	Övrigt
Bellelli G, Morghen S, Tirelli V, Trabucchi M. Motor recovery after rehabilitation and long-term clinical outcomes... Boyd CM, Landefield CS, Counsell SR, et al. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. <i>J Am Geriatr Soc</i> . 2008;56:2171-9. <i>Journal of the American Geriatrics Society</i> . 2009;57(7):1300-1.	Irrelevant studiedesign
Beltz SK. Comprehensive, in-hospital geriatric assessment plus an interdisciplinary home intervention after discharge reduced length of subsequent readmissions and improved functioning... commentary on Nikolaus T, Specht-Leible N, Bach M et al. A randomized trial of comprehensive geriatric assessment and home intervention in the care of hospitalized patients. <i>AGE AGEING</i> 1999 Oct;28(6):543-50. <i>Evidence Based Nursing</i> . 2000;3(3):83-.	Övrigt
Benedict L, Hazelett S, Fleming E, Ludwick R, Anthony M, Fosnight S, et al. Prevention, detection and intervention with delirium in an acute care hospital: a feasibility study. <i>Int J Older People Nurs</i> . 2009;4(3):194-202.	Övrigt
Bentur N, Eldar R, Davies MA. Process and outcome of care: comparison of two inpatient geriatric rehabilitation settings. <i>Clinical Rehabilitation</i> . 1994;8(4):307-13.	Irrelevant intervention
Bentur N, Eldar R. Quality of rehabilitation care in two inpatient geriatric settings. <i>Qual Assur Health Care</i> . 1993;5(3):237-42.	Irrelevant intervention
Berggren M, Stenvall M, Olofsson B, Gustafson Y. Evaluation of a fall-prevention program in older people after femoral neck fracture: a one-year follow-up. <i>Osteoporos Int</i> . 2008;19(6):801-9.	Irrelevant population
Bernardini B, Meinecke C, Pagani M, Grillo A, Fabbrini S, Zaccarini C, et al. Comorbidity and adverse clinical events in the rehabilitation of older adults after hip fracture. <i>Journal of the American Geriatrics Society</i> . 1995;43(8):894-8.	Irrelevant jämförelse-alternativ
Birks Y. A multifactorial intervention after a fall did not prevent falls in elderly patients with cognitive impairment and dementia. <i>Evidence Based Nursing</i> . 2003;6(4):114-5.	Övrigt
Blakemore, S. (2012). Specialist Care for Frail Older People. <i>Emergency Nurse</i> , 19, 12-16.	Irrelevant studiedesign
Blanc-Bisson C, Dechamps A, Gouspillou G, Dehail P, Bourdel-Marchasson I. A randomized controlled trial on early physiotherapy intervention versus usual care in acute care unit for elderly: potential benefits in light of dietary intakes. <i>J Nutr Health Aging</i> . 2008;12(6):395-9.	Irrelevant intervention
Bleijenberg, N., Drubbel, I., Ten, D., V, H., Numans, M. E., Schuurmans, M. J., . . . N, J. (2012). Proactive and integrated primary care for frail older people: design and methodological challenges of the Utrecht primary care PROactive frailty intervention trial (U-PROFIT). <i>BMC Geriatr</i> , 12, 16. doi: 10.1186/1471-2318-12-16.	Övrigt
Bo M, Martini B, Ruatta C, Massaia M, Ricauda NA, Varetti A, et al. Geriatric ward hospitalization reduced incidence delirium among older medical inpatients. <i>Am J Geriatr Psychiatry</i> . 2009;17(9):760-8.	Irrelevant studiedesign

Boltz M, Capezuti E, Bowar-Ferres S, Norman R, Secic M, Kim H, et al. Changes in the Geriatric Care Environment Associated with NICHE (Nurses Improving Care for HealthSystem Elders). <i>Geriatr Nurs.</i> 2008;29(3):176-85.	Irrelevant population
Borok GM, Reuben DB, Zendle LJ, Ershoff DH, Wolde-Tsadik G, Rubenstein LZ, et al. Rationale and design of a multi-center randomized trial of comprehensive geriatric assessment consultation for hospitalized patients in an HMO. <i>J Am Geriatr Soc.</i> 1994;42(5):536-44.	Övrigt
Boyd M, Koziol-McLain J, Yates K, Kerse N, McLean C, Pilcher C, et al. Emergency department case-finding for high-risk older adults: The Brief Risk Identification for Geriatric Health Tool (BRIGHT). <i>Academic Emergency Medicine.</i> 2008;15(7):598-606.	Irrelevant studiedesign
Boyd, CM, Landefeld, CS, Counsell, SR, Palmer, RM, Fortinsky, RH, Kresevic, D, Burant, C, Covinsky, K, (2008. Recovery in Activities of Daily Living Among Older Adults Following, Hospitalization for Acute Medical Illness. <i>J Am Geriatr Soc.</i> 56(12) 2171-2179.	Irrelevant studiedesign
Boyd M, Koziol-McLain J, Yates K, Kerse N, McLean C, Pilcher C. Brief screen for high risk older adults in an emergency department... 2007 Society for Academic Emergency Medicine Annual Meeting. <i>Academic Emergency Medicine.</i> 2007;14(5):S151-2.	Irrelevant studiedesign
Boyer N, Chuang JL, Gipner D. An acute care geriatric unit. <i>Nurs Manage.</i> 1986;17(5):22-5.	Irrelevant intervention
Brand, C. A., Martin-Khan, M., Wright, O., Jones, R. N., Morris, J. N., Travers, C. M., . . . Gray, L. C. (2011). Development of quality indicators for monitoring outcomes of frail elderly hospitalised in acute care health settings: study protocol. <i>BMC Health Serv Res,</i> 11, 281. doi: 10.1186/1472-6963-11-281.	Övrigt
Briggs D. A semirecumbent body position led to a lower rate of nosocomial pneumonia than a supine position in mechanically ventilated adults... commentary on Drakulovic MB, Torres A, Bauer TT et al. Supine body position as a risk factor for nosocomial pneumonia in mechanically ventilated patients: a randomised trial. <i>LANCET</i> 1999 Nov 27;354(9193):1851-8, 1835. <i>Evidence Based Nursing.</i> 2000;3(3):84-.	Irrelevant studiedesign
Broslawski GE, Elkins M, Algus M. Functional abilities of elderly survivors of intensive care. <i>J Am Osteopath Assoc.</i> 1995;95(12):712-7.	Irrelevant jämförelse-alternativ
Brymer C, Cavanagh P, Denomy E, Wells K, Cook C. The effect of a geriatric education program on emergency nurses. <i>J Emerg Nurs.</i> 2001;27(1):27-32.	Irrelevant intervention
Bugdayci D, Paker N, Dere D, Ozdemir E, Ince N. Frequency, features, and factors for falls in a group of subacute stroke patients hospitalized for rehabilitation in Istanbul. <i>Arch Gerontol Geriatr.</i> 2011;52(3):e215-9.	Irrelevant studiedesign
Burns R, Nichols LO, Graney MJ, Cloar FT. Impact of continued geriatric outpatient management on health outcomes of older veterans. <i>Arch Intern Med.</i> 1995;155(12):1313-8.	Irrelevant population
Buurman BM, Parlevliet JL, van D, B A, de H, R J, et al. A randomised clinical trial on a comprehensive geriatric assessment and intensive home follow-up after hospital discharge: the Transitional Care Bridge. <i>BMC Health Serv Res.</i> 2010;10:296.	Irrelevant studiedesign
Buurman BM, van M, B C, Korevaar JC, Abu-Hanna A, Levi M, et al. Prognostication in acutely admitted older patients by nurses and physicians. <i>J Gen Intern Med.</i> 2008;23(11):1883-9.	Irrelevant studiedesign
Buurman, B. M., Hoogerduijn, J. G., de, H., R, J., Abu-Hanna, A., Lagaay, A. M., . . . S, E. (2011). Geriatric conditions in acutely hospitalized older patients: prevalence and one-year survival and functional decline. <i>PLoS One,</i> 6, e26951. doi: 10.1371/journal.pone.0026951. Epub 2011 Nov 14.	Irrelevant jämförelse-alternativ

Buurman, B. M., Hoogerduijn, J. G., van, G., E, A., de, H., R, J., . . . S, E. (2012). Clinical characteristics and outcomes of hospitalized older patients with distinct risk profiles for functional decline: a prospective cohort study. <i>PLoS One</i> , 7, e29621. doi: 10.1371/journal.pone.0029621. Epub 2012 Jan 4.	Irrelevant jämförelse-alternativ
Camicoli RM, Kaye JA, Brummel-Smith K. Recognition of neurologic diseases in geriatric inpatients. <i>Acta Neurol Scand</i> . 1998;97(4):265-70.	Irrelevant intervention
Caplan GA, Harper EL. Recruitment of volunteers to improve vitality in the elderly: The REVIVE study. <i>Internal Medicine Journal</i> . 2007;37(2):95-100.	Irrelevant studiedesign
Caplan Gideon A, Williams Anthony J, Daly B, Abraham K. A Randomized, Controlled Trial of Comprehensive Geriatric Assessment and Multidisciplinary Intervention After Discharge of Elderly from the Emergency Department--The DEED II Study. <i>Journal of the American Geriatrics Society</i> . 2004;52(9):1417-23.	Irrelevant studiedesign
Cartwright DL, Madill HM, Dennis S. Cognitive impairment and functional performance of patients admitted to a geriatric assessment and rehabilitation centre. <i>Physical &amp; Occupational Therapy in Geriatrics</i> . 1996;14(3):1-21.	Irrelevant population
Cesari M, Colloca G, Cerullo F, Ferrini A, Testa AC, Foti E, et al. Onco-geriatric approach for the management of older patients with cancer. <i>Journal of the American Medical Directors Association</i> . 2011;12(2):153-9.	Irrelevant population
Cesari M, Kritchevsky SB, Penninx BW, Nicklas BJ, Simonsick EM, Newman AB, et al. Prognostic value of usual gait speed in well-functioning older people -- results from the health, aging and body composition study. <i>Journal of the American Geriatrics Society</i> . 2005;53(10):1675-80.	Irrelevant population
Chambers N. Into the unknown: establishing a new service in an acute hospital setting. <i>Nurs Older People</i> . 2004;16(8):18-20.	Irrelevant studiedesign
Chang H-H, Tsai S-L, Chen C-Y, Liu W-J. Outcomes of hospitalized elderly patients with geriatric syndrome: Report of a community hospital reform plan in Taiwan. <i>Archives of Gerontology and Geriatrics</i> . 2010;50(Suppl 1):S30-S3.	Irrelevant intervention
Chen CC, Huang G, Hung Y, Liu Y. Cognitive, nutritional, and functional decline associated with hospitalization in older Taiwanese patients admitted for med-surgical services: Sigma Theta Tau International; 2005.	Irrelevant intervention
Chen LK, Chen YM, Hwang SJ, Peng LN, Lin MH, Lee WJ, et al. Effectiveness of community hospital-based post-acute care on functional recovery and 12-month mortality in older patients: a prospective cohort study. <i>Ann Med</i> . 2010;42(8):630-6.	Irrelevant studiedesign
Chen Y-M, Chuang Y-W, Liao S-C, Lin C-S, Yang S-H, Tang Y-J, et al. Predictors of functional recovery (FR) for elderly hospitalized patients in a geriatric evaluation and management unit (GEMU) in Taiwan. <i>Archives of Gerontology and Geriatrics</i> . 2010;50(Suppl 1):S1-S5.	Irrelevant studiedesign
Chin JJ, Sahadevan S, Tan CY, Ho SC, Choo PW. Critical role of functional decline in delayed discharge from an acute geriatric unit. <i>Ann Acad Med Singapore</i> . 2001;30(6):593-9.	Irrelevant studiedesign
Chiavenda P, Vincentelli GM, Alegiani F. Cognitive impairment in elderly ED patients: need for multidimensional assessment for better management after discharge. <i>Am J Emerg Med</i> . 2002;20(4):332-5.	Irrelevant population
Chong MS, Chan MP, Kang J, Han HC, Ding YY, Tan TL. A new model of delirium care in the acute geriatric setting: geriatric monitoring unit. <i>BMC Geriatr</i> . 2011;11:41.	Övrigt
Chong, M. S., Empensando, E. F., Ding, Y. Y., & Tan, T. L. (2012). A subacute model of geriatric care for frail older persons: The tan tock seng hospital experience. <i>Annals of the Academy of Medicine Singapore</i> , 41, 354-361.	Irrelevant studiedesign

Chou, M. Y., Liang, C. K., Lin, Y. T., Chen, L. K., Peng, L. N., Liu, L. K., . . . Tu, M. S. (2012). Screening postacute care needs of hospital inpatients in Taiwan: A hospital-based study. <i>Journal of Clinical Gerontology and Geriatrics</i> , 3, 132-135.	Irrelevant studiedesign
Clarfield AM. A long-term geriatric teaching ward in an acute-care hospital: a three-year experience. <i>J Am Geriatr Soc</i> . 1982;30(7):457-65.	Irrelevant population
Close J, Ellis M, Hooper R, Glucksman E, Jackson S, Swift C. Prevention of falls in the elderly trial (PROFET): a randomised controlled trial. <i>Lancet</i> . 1999;353(9147):93-7.	Irrelevant intervention
Cockram A, Gibb R, Kalra L. The role of a specialist team in implementing continuing health care guidelines in hospitalized patients. <i>Age &amp; Ageing</i> . 1997;26(3):211-6.	Irrelevant population
Cogan, L., Martin, A. J., Kelly, L. A., Duggan, J., Hynes, D., & Power, D. (2010). An audit of hip fracture services in the Mater Hospital Dublin 2001 compared with 2006. <i>Ir J Med Sci</i> , 179(1), 51-55.	Irrelevant population
Cohen Harvey J, Saltz Constance C, Samsa G, McVey L, et al. Predictors of two-year post-hospitalization mortality among elderly veterans in a study evaluating a geriatric consultation team. <i>Journal of the American Geriatrics Society</i> . 1992;40(12):1231-5.	Irrelevant studiedesign
Coker E, Oliver D. Evaluation of the STRATIFY falls prediction tool on a geriatric unit. <i>Outcomes Manag</i> . 2003;7(1):8-14; quiz 5.	Irrelevant intervention
Cole MG, Fenton FR, Engelmann F, Mansouri I. Effectiveness of geriatric psychiatry consultation in an acute care hospital: a randomized clinical trial. <i>J Am Geriatr Soc</i> . 1991;39(12):1183-8.	Irrelevant intervention
Cole, M. G., McCusker, J., Elie, M., Dendukuri, N., Latimer, E., & Belzile, E. (2006). Systematic detection and multidisciplinary care of depression in older medical inpatients: a randomized trial. <i>CMAJ</i> , 174(1), 38-44.	Irrelevant intervention
Cole, M. G., Primeau, F. J., Bailey, R. F., Bonnycastle, M. J., Masciarelli, F., Engelmann, F., . . . Ducic, D. (1994). Systematic intervention for elderly inpatients with delirium: a randomized trial. <i>CMAJ</i> , 151(7), 965-970.	Irrelevant intervention
Coleman, S. A., Cunningham, C. J., Walsh, J. B., Coakley, D., Harbison, J., Casey, M., . . . Horgan, N. F. (2012). Outcomes among older people in a post-acute inpatient rehabilitation unit. <i>Disabil Rehabil</i> , 34, 1333-1338. doi: 10.3109/09638288.2011.636136. Epub 2011 Dec 8.	Irrelevant studiedesign
Collard AF. Predicting readmissions of elderly patients to the acute care hospital 1988.	Irrelevant population
Colprim, D., Martin, R., Parer, M., Prieto, J., Espinosa, L., & Inzitari, M. (2013). Direct Admission to Intermediate Care for Older Adults With Reactivated Chronic Diseases as an Alternative to Conventional Hospitalization. <i>J Am Med Dir Assoc</i> . doi: 10.1016/j.jamda.2012.12.003.	Irrelevant jämförelse-alternativ
Conroy Simon, P., Stevens, T., & Parker Stuart, G. (2011). A systematic review of comprehensive geriatric assessment to improve outcomes for frail older people being rapidly discharged from acute hospital: 'interface geriatrics'. <i>Age &amp; Ageing</i> , 40(4), 436-443.	Övrigt
Courtney, M., Edwards, H., Chang, A., Parker, A., Finlayson, K., & Hamilton, K. (2009). Fewer emergency readmissions and better quality of life for older adults at risk of hospital readmission: A randomized controlled trial to determine the effectiveness of a 24-week exercise and telephone follow-up program. <i>Journal of the American Geriatrics Society</i> , 57(3), 395-402.	Irrelevant intervention
Covinsky KE, King JT, Jr, Quinn LM, Siddique R, Palmer R, et al. Do acute care for elders units increase hospital costs? A cost analysis using the hospital perspective... presented, in part, at the annual session of the Society of General Internal Medicine in San Diego, California on May 5, 1995. <i>Journal of the American Geriatrics Society</i> . 1997;45(6):729-34.	Irrelevanta effektmått

Dawson JA, Reason JF, Chick JR. Measuring dependency in the elderly: The feasibility of incorporating formal assessments into occupational therapy workload. <i>British Journal of Occupational Therapy</i> . 1984;47(3):83-5.	Irrelevant jämförelse-alternativ
de M, N A, Lane K. Validity and reliability of the de Morton Mobility Index in the subacute hospital setting in a geriatric evaluation and management population. <i>J Rehabil Med</i> . 2010;42(10):956-61.	Irrelevant intervention
de O, P C, Dantas RO, de T, L E, Moriguti JC, et al. Clinical and scintigraphic assessment of swallowing of older patients admitted to a tertiary care geriatric ward. <i>Dysphagia</i> (0179051X). 2008;23(1):1-6.	Irrelevant intervention
de S-H, Jamart J, Boland B, Swine C, Cornette P. Comparison of three tools predicting functional decline after hospitalization of older patients. <i>J Am Geriatr Soc</i> . 2010;58(5):1003-5.	Irrelevant intervention
de S-H, Jamart J, Morrhaye G, Martens HJ, Geenen V, Duy V, et al. Serum IL-6 and IGF-1 improve clinical prediction of functional decline after hospitalization in older patients. <i>Aging Clin Exp Res</i> . 2011;23(2):106-11.	Irrelevant studiedesign
De V, O J, Peeters GMEE, Elders PJM, Muller M, Knol DL, et al. Multifactorial intervention to reduce falls in older people at high risk of recurrent falls: A randomized controlled trial. <i>Archives of Internal Medicine</i> . 2010;170(13):1110-7.	Irrelevant population
Del G, Ferretti E, Omiciuolo C, Sceusa R, Zanata C, Manganaro D, et al. The hospital-based, post-acute geriatric evaluation and management unit: the experience of the acute geriatric unit in Trieste. <i>Arch Gerontol Geriatr</i> . 2009;49 Suppl 1:49-60.	Irrelevant jämförelse-alternativ
Del, S., Pulignano, G., Di, L., Tarantini, L., Cioffi, G., Tolone, S., . . . Minardi, G. (2012). Role of a multidisciplinary program in improving outcomes in cognitively impaired heart failure older patients. <i>Monaldi Archives for Chest Disease - Cardiac Series</i> , 78, 20-28.	Irrelevant population
Dellasega CA, Zerbe TM. A multimethod study of advanced practice nurse postdischarge care. <i>Clin Excell Nurse Pract</i> . 2000;4(5):286-93.	Irrelevant studiedesign
Dendukuri N, McCusker J, Belzile E. The identification of seniors at risk screening tool: further evidence of concurrent and predictive validity. <i>J Am Geriatr Soc</i> . 2004;52(2):290-6.	Irrelevant studiedesign
Denkinger MD, Igl W, Coll-Planas L, Bleicher J, Nikolaus T, Jamour M. Evaluation of the short form of the late-life function and disability instrument in geriatric inpatients-validity, responsiveness, and sensitivity to change. <i>J Am Geriatr Soc</i> . 2009;57(2):309-14.	Irrelevant population
Denkinger MD, Igl W, Coll-Planas L, Nikolaus T, Bailer S, Bader A, et al. Practicality, validity and sensitivity to change of fear of falling self-report in hospitalised elderly—a comparison of four instruments. <i>Age Ageing</i> . 2009;38(1):108-12.	Irrelevant population
Deschondt M, Braes T, Broos P, Sermon A, Boonen S, Flamaing J, et al. Effect of an Inpatient Geriatric Consultation Team on Functional Outcome, Mortality, Institutionalization, and Readmission Rate in Older Adults with Hip Fracture: A Controlled Trial. <i>Journal of the American Geriatrics Society</i> . 2011;59(7):1299-308.	Irrelevant population
Desy PM, Prohaska TR. The Geriatric Emergency Nursing Education (GENE) course: an evaluation. <i>J Emerg Nurs</i> . 2008;34(5):396-402.	Irrelevant intervention
Dickes-Sotty, H., Chevalet, P., Fix, M. H., Riaudel, T., Serre-Sahel, C., Ould-Aoudia, V., . . . De, D. (2012). Six-months outcomes after admission in acute geriatric care unit secondary to a fall. <i>Geriatr Psychol Neuropsychiatr Vieil</i> , 10, 391-401.	Irrelevant studiedesign
Ding, Y. Y., Abisheganaden, J., Chong, W. F., Heng, B. H., & Lim, T. K. (2013). Effectiveness of acute geriatric units in the real world: The case of short-term mortality among seniors hospitalized for pneumonia. <i>Geriatr Gerontol Int</i> , 13, 55-62. doi: 10.1111/j.1447-0594.2012.00858.x. Epub 2012 Apr 11.	Irrelevant studiedesign

Drame M, Fierobe F, Lang PO, Jolly D, Boyer F, Mahmoudi R, et al. Predictors of institution admission in the year following acute hospitalisation of elderly people. <i>J Nutr Health Aging</i> . 2011;15(5):399-403.	Irrelevant jämförelse-alternativ
Drame M, Jovenin N, Novella JL, Lang PO, Somme D, Laniece I, et al. Predicting early mortality among elderly patients hospitalised in medical wards via emergency department: the SAFES cohort study. <i>J Nutr Health Aging</i> . 2008;12(8):599-604.	Irrelevant intervention
Drame M, Novella JL, Lang PO, Somme D, Jovenin N, Laniece I, et al. Derivation and validation of a mortality-risk index from a cohort of frail elderly patients hospitalised in medical wards via emergencies: the SAFES study. <i>Eur J Epidemiol</i> . 2008;23(12):783-91.	Irrelevant intervention
Drame, M., Lang, P. O., Novella, J. L., Narbey, D., Mahmoudi, R., Laniece, I., . . . Jolly, D. (2012). Six-month outcome of elderly people hospitalized via the emergency department: the SAFES cohort. <i>Rev Epidemiol Sante Publique</i> , 60, 189-196. doi: 10.1016/j.respe.2011.11.004. Epub 2012 May 16.	Irrelevant jämförelse-alternativ
Dunn RB, Lewis PA, Vetter NJ, Guy PM, Hardman CS, Jones RW. Health visitor intervention to reduce days of unplanned hospital re-admission in patients recently discharged from geriatric wards: The results of a randomised controlled study. <i>Archives of Gerontology and Geriatrics</i> . 1994;18(1):15-23.	Irrelevant intervention
Edmans J, Conroy S, Harwood R, Lewis S, Elliott RA, Logan P, et al. Acute medical unit comprehensive geriatric assessment intervention study (AMIGOS). <i>Trials</i> . 2011;12:200.	Övrigt
Edmans, J., Conroy, S., Harwood, R., Lewis, S., Elliott, R. A., Logan, P., . . . Gladman, J. (2011). Acute medical unit comprehensive geriatric assessment intervention study (AMIGOS): Study protocol for a randomised controlled trial. <i>Trials</i> , 12.	Övrigt
Ellis, G., Jamieson, C. A., Alcorn, M., & Devlin, V. (2012). An Acute Care for Elders (ACE) unit in the emergency department. <i>European Geriatric Medicine</i> , 3, 261-263.	Irrelevant studiedesign
Ellis, G., Whitehead Martin, A., Robinson, D., O'Neill, D., & Langhorne, P. (2011). Comprehensive geriatric assessment for older adults admitted to hospital: Meta-analysis of randomised controlled trials. <i>BMJ: British Medical Journal</i> , 343, 1-10.	Övrigt
Engelhardt JB, Toseland RW, O'Donnell JC, Richie JT, Jue D, Banks S. The effectiveness and efficiency of outpatient geriatric evaluation and management. <i>J Am Geriatr Soc</i> . 1996;44(7):847-56.	Irrelevant population
Ensberg MD, Paletta MJ, Galecki AT, Dacko CL, Fries BE. Identifying elderly patients for early discharge after hospitalization for hip fracture. <i>J Gerontol</i> . 1993;48(5):M187-95.	Irrelevant population
Fairhall N, Aggar C, Kurrle SE, Sherrington C, Lord S, Lockwood K, et al. Frailty Intervention Trial (FIT). <i>BMC Geriatr</i> . 2008;8:27.	Irrelevant population
Fisher S, Ottenbacher KJ, Goodwin JS, Graham JE, Ostir GV. Short Physical Performance Battery in hospitalized older adults. <i>Aging Clin Exp Res</i> . 2009;21(6):445-52.	Irrelevant intervention
Fisher Steven R. In hospital functional assessments in the elderly and their association to post discharge health outcomes. Dissertation Abstracts International: Section B: The Sciences and Engineering. 2009;70(4-B):2242.	Irrelevant intervention
Flamer HE, Christophidis N, Margetts C, Ugoni A, McLean AJ. Extended hospital stays with increasing age: the impact of an acute geriatric unit. <i>Med J Aust</i> . 1996;164(1):10-3.	Irrelevant population
Fleck, S. J., Bustamante-Ara, N., Ortiz, J., Vidan, M. T., Lucia, A., & Serra-Rexach, J. A. (2012). Activity in GEriatric acute CARe (AGECAR): rationale, design and methods. <i>BMC Geriatr</i> , 12, 28. doi: 10.1186/1471-2318-12-28.	Övrigt
Flood KL, Brown CJ, Carroll MB, Locher JL. Nutritional processes of care for older adults admitted to an oncology-acute care for elders unit. <i>Critical Reviews in Oncology/Hematology</i> . 2011;78(1):73-8.	Irrelevanta effektmått

Flood KL, Carroll MB, Le CV, Ball L, Esker DA, Carr DB. Geriatric syndromes in elderly patients admitted to an oncology-acute care for elders unit. <i>Journal of Clinical Oncology</i> . 2006;24(15):2298-303.	Irrelevant intervention
Foo Chik, L., Siu Vivian Wing, Y., Tan Thai, L., Ding Yew, Y., & Seow, E. (2012). Geriatric assessment and intervention in an emergency department observation unit reduced re-attendance and hospitalisation rates. <i>Australasian Journal on Ageing</i> , 31, 40-46.	Irrelevant studiedesign
Forasassi C, Golmard JL, Pautas E, Piette F, Myara I, Raynaud-Simon A. Inflammation and disability as risk factors for mortality in elderly acute care patients. <i>Archives of Gerontology and Geriatrics</i> . 2009;48(3):406-10.	Irrelevant intervention
Formiga F, Chivite D, Solé A, Manito N, Ramon Josep M, Pujol R. Functional outcomes of elderly patients after the first hospital admission for decompensated heart failure (HF): A prospective study. <i>Archives of Gerontology and Geriatrics</i> . 2006;43(2):175-85.	Irrelevant intervention
Fox, M. T., Persaud, M., Maimets, I., O'Brien, K., Brooks, D., Tregunno, D., & Schraa, E. (2012). Effectiveness of acute geriatric unit care using acute care for elders components: a systematic review and meta-analysis. <i>J Am Geriatr Soc</i> , 60, 2237-2245. doi: 10.1111/jgs.12028. Epub 2012 Nov 23.	Övrigt
Freeman M. Quick response programs: effective management of a population in crisis. <i>Leadersh Health Serv</i> . 1994;3(5):36-9.	Irrelevant population
Gayton D, Wood-Dauphinee S, de L, Tousignant P, Hanley J. Trial of a geriatric consultation team in an acute care hospital. <i>J Am Geriatr Soc</i> . 1987;35(8):726-36.	Stora brister studiekvalitet
Gazzotti C, Albert A, Pepinster A, Petermans J. Clinical usefulness of the mini nutritional assessment (MNA) scale in geriatric medicine. <i>J Nutr Health Aging</i> . 2000;4(3):176-81.	Irrelevant intervention
Germain M, Knoeffel F, Wieland D, Rubenstein LZ. A geriatric assessment and intervention team for hospital inpatients awaiting transfer to a geriatric unit: a randomized trial. <i>Aging (Milano)</i> . 1995;7(1):55-60.	Irrelevant population
Gharacholou, S. M., Sloane, R., Cohen, H. J., & Schmader, K. E. (2012). Geriatric inpatient units in the care of hospitalized frail adults with a history of heart failure. <i>International Journal of Gerontology</i> , 6, 112-116.	Subgrupp från Cohen et al 2002
Gilchrist WJ, Newman RJ, Hamblen DL, Williams BO. Prospective randomised study of an orthopaedic geriatric inpatient service. <i>BMJ</i> . 1988;297(6656):1116-8.	Irrelevant population
Gill Sudeep S, Gibson M, Al-Hammadi A. Economic implications of Discharge of Elderly from the Emergency Department trial results. <i>Journal of the American Geriatrics Society</i> . 2005;53(9):1642.	Övrigt
Gonzalez-Montalvo JI, Mauleon JL, Gil-Garay E, Gotor P, Martin-Vega A. The orthogeriatric unit for acute patients: A new model of care that improves efficiency in the management of patients with hip fracture. <i>HIP International</i> . 2010;20(2):229-35.	Irrelevant population
Gray LC, Bernabei R, Berg K, Finne-Soveri H, Fries BE, Hirdes JP, et al. Standardizing assessment of elderly people in acute care: the interRAI Acute Care instrument. <i>J Am Geriatr Soc</i> . 2008;56(3):536-41.	Irrelevant intervention
Greenberg NS, Rosin AJ. Factors influencing admission or nonadmission of the aged to the hospital. <i>J Am Geriatr Soc</i> . 1982;30(10):635-41.	Irrelevant intervention
Gregersen, M., Pedersen, A. B., & Damsgaard, E. M. (2012). Comprehensive geriatric assessment increases 30-day survival in the aged acute medical inpatients. <i>Dan Med J</i> , 59, A4442.	Irrelevant studiedesign
Gross PF. Long term care and alternative site therapy project. <i>Pride Inst J Long Term Home Health Care</i> . 1991;10(3):29-41.	Irrelevant population

Hamaker, M. E., Buurman, B. M., van, M., B, C., Kuper, I. M. J. A., Smorenburg, C. H., . . . S, E. (2011). The value of a comprehensive geriatric assessment for patient care in acutely hospitalized older patients with cancer. <i>Oncologist</i> , 16, 1403-1412.	Irrelevant population
Harari D, Martin FC, Buttery A, O'Neill S, Hopper A. The older persons' assessment and liaison team 'OPAL': evaluation of comprehensive geriatric assessment in acute medical inpatients. <i>Age Ageing</i> . 2007;36(6):670-5.	Irrelevant studiedesign
Harris RD, Henschke PJ, Popplewell PY, Radford AJ, Bond MJ, Turnbull RJ, et al. A randomised study of outcomes in a defined group of acutely ill elderly patients managed in a geriatric assessment unit or a general medical unit. <i>Aust N Z J Med</i> . 1991;21(2):230-4.	Övrigt
Hegney D, Buikstra E, Chamberlain C, March J, McKay M, Cope G, et al. Nurse discharge planning in the emergency department: a Toowoomba, Australia, study. <i>J Clin Nurs</i> . 2006;15(8):1033-44.	Irrelevant intervention
Hempsall VJ, Robertson DR, Campbell MJ, Briggs RS. Orthopaedic geriatric care--is it effective? A prospective population-based comparison of outcome in fractured neck of femur. <i>J R Coll Physicians Lond</i> . 1990;24(1):47-50.	Irrelevant population
Ho WWS, Kwan D, D L, Liu KW, Chow KM, Lau E, et al. To investigate the effect and cost-effectiveness of implementing an orthogeriatric intervention for elderly patients with acute hip fracture: the experience in Hong Kong. <i>Journal of the American Geriatrics Society</i> . 2009;57(11):2153-4.	Irrelevant population
Hogan DB, Fox RA. A prospective controlled trial of a geriatric consultation team in an acute-care hospital. <i>Age and Ageing</i> . 1990;19(2):107-13.	Irrelevant studiedesign
Holm, A., Michel, M., Stern, G. A., Hung, T., Klein, T., Flaherty, L., . . . Maletta, G. (1999). The outcomes of an inpatient treatment program for geriatric patients with dementia and dysfunctional behaviors. <i>Gerontologist</i> , 39(6), 668-676.	Irrelevant intervention
Holmboe, E. S., Meehan, T. P., Radford, M. J., Wang, Y., Marciak, T. A., & Krumholz, H. M. (1999). Use of critical pathways to improve the care of patients with acute myocardial infarction. <i>Am J Med</i> , 107(4), 324-331.	Irrelevant population
Hoogerduijn, J., Buurman, B., de, R., Sophia, Korevaar, J., Grobbee, D., & Schuurmans, M. (2010). ISAR-HP an instrument predicting functional decline in older hospitalized patients... Fourth European Nursing Congress. <i>Journal of Clinical Nursing</i> , 19, 37-37.	Irrelevant intervention
Hoolsma, E. M. (1996). Proving our worth: using treatment outcomes to demonstrate cost efficiency. <i>Topics in Geriatric Rehabilitation</i> , 11(4), 84-94.	Övrigt
Hopper A, Martin FC, Buttery A, O'Neill S, McGovern R, Shillo P, et al. The older persons assessment and liaison team 'OPAL': pilot evaluation of comprehensive geriatric assessment (CGA) in acutemedical inpatients... British Geriatrics Society: Abstracts of papers presented at the Spring Scientific Meeting, 6-7 April 2006. <i>Age &amp; Ageing</i> . 2006;35:i9-i.	Irrelevant intervention
Hoverman, C., Shugarman, L. R., Saliba, D., & Buntin, M. B. (2008). Use of postacute care by nursing home residents hospitalized for stroke or hip fracture: how prevalent and to what end? <i>J Am Geriatr Soc</i> , 56(8), 1490-1496.	Irrelevant population
Huang, T., & Liang, S. (2005). A randomized clinical trial of the effectiveness of a discharge planning intervention in hospitalized elders with hip fracture due to falling. <i>Journal of Clinical Nursing</i> , 14(10), 1193-1201.	Irrelevant population
Huber M, Kennard A. Functional and mental status outcomes of clients discharged from acute gerontological versus medical/surgical units. <i>J Gerontol Nurs</i> . 1991;17(7):20-4.	Irrelevant population
Hui E, Lum CM, Woo J, Or KH, Kay RL. Outcomes of elderly stroke patients. Day hospital versus conventional medical management. <i>Stroke</i> . 1995;26(9):1616-9.	Irrelevant intervention

Huusko Tiina M, Karppi P, Avikainen V, Kautiainen H, Sulkava R. Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: Subgroup analysis of patients with dementia. <i>BMJ: British Medical Journal</i> . 2000;321(7269):1107-11.	Irrelevant population
Huws, D. W., Cashmore, D., Newcombe, R. G., Roberts, C., Vincent, J., & Elwyn, G. (2008). Impact of case management by advanced practice nurses in primary care on unplanned hospital admissions: a controlled intervention study. <i>BMC Health Serv Res</i> , 8, 115.	Irrelevant intervention
Huynh, B. C., Rovner, A., & Rich, M. W. (2006). Long-term survival in elderly patients hospitalized for heart failure: 14-year follow-up from a prospective randomized trial. <i>Arch Intern Med</i> , 166(17), 1892-1898.	Irrelevant population
Inouye Sharon, K., Wagner, D. R., Acampora, D., Horwitz Ralph, I., & et al. (1993). A controlled trial of a nursing-centered intervention in hospitalized elderly medical patients: The Yale Geriatric Care Program. <i>Journal of the American Geriatrics Society</i> , 41(12), 1353-1360.	Övrigt
Inouye SK, Bogardus ST, Jr, Baker DI, Leo-Summers L, Cooney LM. The Hospital Elder Life Program: a model of care to prevent cognitive and functional decline in older hospitalized patients. <i>Hospital Elder Life Program. J Am Geriatr Soc</i> . 2000;48(12):1697-706.	Irrelevant intervention
Inouye SK, Wagner DR, Acampora D, Horwitz RI, Cooney LM, Jr, et al. A predictive index for functional decline in hospitalized elderly medical patients. <i>J Gen Intern Med</i> . 1993;8(12):645-52.	Irrelevant population
Inouye, S. K., Acampora, D., Miller, R. L., Fulmer, T., Hurst, L. D., Cooney, Jr., & L, M. (1993). The Yale Geriatric Care Program: A model of care to prevent functional decline in hospitalized elderly patients. <i>Journal of the American Geriatrics Society</i> , 41(12), 1345-1352.	Irrelevant studiedesign
Inouye, S. K., Bogardus, S. T., Jr, Baker, D. I., Leo-Summers, L., Cooney, L. M., & Jr. (2000). The Hospital Elder Life Program: a model of care to prevent cognitive and functional decline in older hospitalized patients. <i>Journal of the American Geriatrics Society</i> , 48(12), 1697-1706.	Irrelevant studiedesign
Inouye, S. K., Bogardus, S. T., Jr, Charpentier, P. A., Leo-Summers, L., Acampora, D., . . . Jr. (1999). A multicomponent intervention to prevent delirium in hospitalized older patients. <i>New England Journal of Medicine</i> , 340(9), 669-676.	Irrelevant population
Inouye, S. K., Peduzzi, P. N., Robison, J. T., Hughes, J. S., Horwitz, R. I., & Concato, J. (1998). Importance of functional measures in predicting mortality among older hospitalized patients. <i>JAMA: Journal of the American Medical Association</i> , 279(15), 1187-1193.	Övrigt
Ip SP, Leung YF, Ip CY, Mak WP. Outcomes of critically ill elderly patients: is high-dependency care for geriatric patients worthwhile? <i>Crit Care Med</i> . 1999;27(11):2351-7.	Irrelevant intervention
Jack, B., Hillier, V., Williams, A., & Oldham, J. (2003). Hospital based palliative care teams improve the symptoms of cancer patients. <i>Palliative Medicine</i> , 17(6), 498-502.	Irrelevant intervention
Jackson, A., & Warren, K. (2006). Does consultant geriatrician review on post take ward rounds (PTWR) effect elderly patients' length of stay?... British Geriatrics Society: Abstracts of papers presented at the Spring Scientific Meeting, 6-7 April 2006. <i>Age &amp; Ageing</i> , 35, i28-i28.	Irrelevant intervention
Jahangiri, M., Laborde, J. C., Roy, D., Williams, F., Abdulkareem, N., & Brecker, S. (2011). Outcome of patients with aortic stenosis referred to a multidisciplinary meeting for transcatheter valve. <i>Ann Thorac Surg</i> , 91(2), 411-415.	Irrelevant population
Jahnigen, D. W., Kramer, A. M., Robbins, L. J., Klingbeil, H., & DeVore, P. (1985). Academic affiliation with a nursing home. Impact on patient outcome. <i>J Am Geriatr Soc</i> , 33(7), 472-478.	Irrelevant population
Jain, R., Evenson, A., Biddison, E., Dalal, D., Kelly, K. M., Karmand, A. J., . . . Plantholt, S. (2010). Efficacy of multidisciplinary outpatient management (MOM) program in long term heart failure care. <i>Southern Medical Journal</i> , 103(2), 131-137.	Irrelevant population

Jayadevappa, R., Chhatre, S., Weiner, M., & Raziano, D. B. (2006). Health resource utilization and medical care cost of acute care elderly unit patients. <i>Value Health</i> , 9(3), 186-192.	Irrelevant studiedesign
Jitapunkul, S., Pillay, I., & Ebrahim, S. (1992). Delirium in newly admitted elderly patients: a prospective study. <i>Q J Med</i> , 83(300), 307-314.	Övrigt
Johanson, M. A., Cohen, B. A., Snyder, K. H., McKinley, A. J., & Scott, M. L. (2009). Outcomes for aging adults following total hip arthroplasty in an acute rehabilitation facility versus a subacute rehabilitation facility: a pilot study [corrected] [published erratum appears in J GERIATR PHYS THER 2009;32(3):110]. <i>Journal of Geriatric Physical Therapy</i> , 32(2), 29-34.	Irrelevant population
Johansson, B., Holmberg, L., Berglund, G., Brandberg, Y., Hellbom, M., Persson, C., . . . Sjoden, P. O. (2001). Reduced utilisation of specialist care among elderly cancer patients: a randomised study of a primary healthcare intervention. <i>Eur J Cancer</i> , 37(17), 2161-2168.	Irrelevant population
Jonsen, E., Ljunggren, G., Jonsson, P. V., & Gosta, B. (2007). Functional status in elderly people after acute care and quality of life at one-year follow-up. <i>Health Science Journal</i> , 1(1), 1-14.	Övrigt
Jonsson, P. V., Noro, A., Finne-Soveri, H., Jensdottir, A. B., Ljunggren, G., Bucht, G., . . . Schroll, M. (2008). Admission profile is predictive of outcome in acute hospital care. <i>Aging Clinical &amp; Experimental Research</i> , 20(6), 533-539.	Övrigt
Joseph, A., & Boult, C. (1998). Managed primary care of nursing home residents. <i>J Am Geriatr Soc</i> , 46(9), 1152-1156.	Irrelevant intervention
Kaczorowski, J., Chambers, L. W., Dolovich, L., Paterson, J. M., Karwalajtys, T., Gierman, T., . . . Sabaldt, R. J. (2011). Improving cardiovascular health at population level: 39 community cluster randomised trial of Cardiovascular Health Awareness Program (CHAP). <i>BMJ</i> , 342, d442. doi: 10.1136/bmj.d442.	Irrelevant studiedesign
Kafetz K, O'Farrell J, Parry A, Wijesuriya V, McElligott G, Rossiter B, et al. Age-related geriatric medicine: relevance of special skills of geriatric medicine to elderly people admitted to hospital as medical emergencies. <i>J R Soc Med</i> . 1995;88(11):629-33.	Irrelevant intervention
Kanda, A., Ebihara, S., Yasuda, H., Takashi, O., Sasaki, T., & Sasaki, H. (2004). A Combinatorial Therapy for Pneumonia in Elderly People. <i>Journal of the American Geriatrics Society</i> , 52(5), 846-847.	Irrelevant population
Karppi P, Tilvis R. Effectiveness of a Finnish geriatric inpatient assessment. Two-year follow up of a randomized clinical trial on community-dwelling patients. <i>Scand J Prim Health Care</i> . 1995;13(2):93-8.	Irrelevant population
Kay G, MacTavish M, Moffatt C, Lau G. Development and evaluation of a geriatric assessment unit in a community hospital. <i>Perspectives</i> . 1992;16(3):2-9.	Irrelevant population
Kehusmaa, S., Autti-Ramo, I., Valaste, M., Hinkka, K., & Rissanen, P. (2010). Economic evaluation of a geriatric rehabilitation programme: a randomized controlled trial. <i>J Rehabil Med</i> , 42(10), 949-955.	Irrelevant population
Kellogg FR, Brickner PW, Conley L, Conroy M. Controlling hospital readmission of elderly persons living at home: a risk factor analysis. <i>Home Health Care Services Quarterly</i> . 1991;12(2):5-16.	Irrelevant intervention
Kennie, D. C., Reid, J., Richardson, I. R., Kiamari, A. A., & Kelt, C. (1988). Effectiveness of geriatric rehabilitative care after fractures of the proximal femur in elderly women: a randomised clinical trial. <i>BMJ</i> , 297(6656), 1083-1086.	Irrelevant population
Kenny, R. A., O'Shea, D., & Walker, H. F. (2002). Impact of a dedicated syncope and falls facility for older adults on emergency beds. <i>Age &amp; Ageing</i> , 31(4), 272-275.	Irrelevant intervention
Kergoat, M.-J., Latour, J., Lebel, P., Leclerc, B.-S., Leduc, N., Béland, F., . . . Bolduc, A. (2012). Quality-of-Care Processes in Geriatric Assessment Units: Principles, Practice, and Outcomes. <i>Journal of the American Medical Directors Association</i> , 13, 459-463.	Irrelevant studiedesign

Khan, R., Fernandez, C., Kashif, F., Shedden, R., & Diggory, P. (2002). Combined orthogeriatric care in the management of hip fractures: A prospective study. <i>Annals of the Royal College of Surgeons of England</i> , 84(2), 122-124.	Irrelevant population
Khan, S. A., Millington, H., & Miskelly, F. G. (1997). Benefits of an accident and emergency short stay ward in the staged hospital care of elderly patients. <i>Journal of Accident and Emergency Medicine</i> , 14(3), 151-152.	Övrigt
Kim, M. M., Barnato, A. E., Angus, D. C., Fleisher, L. F., & Kahn, J. M. (2010). The effect of multidisciplinary care teams on intensive care unit mortality. <i>Archives of Internal Medicine</i> , 170(4), 369-376.	Irrelevant studiedesign
Kirchberger, I., Meisinger, C., Seidl, H., Wende, R., Kuch, B., & Holle, R. (2010). Nurse-based case management for aged patients with myocardial infarction: study protocol of a randomized controlled trial. <i>BMC Geriatr</i> , 10, 29.	Irrelevant population
Kleinpell, R. M., & Ferrans, C. E. (1998). Factors influencing intensive care unit survival for critically ill elderly patients. <i>Heart Lung</i> , 27(5), 337-343.	Irrelevant studiedesign
Klopotowska, J. E., Wierenga, P. C., de, R., S, E., Stuijt, C. C., Arisz, L., . . . Smorenburg, S. M. (2011). The effect of an active on-ward participation of hospital pharmacists in Internal Medicine teams on preventable Adverse Drug Events in elderly inpatients: protocol of the WINGS study (Ward-oriented pharmacy in newly admitted geriatric seniors). <i>BMC Health Serv Res</i> , 11, 124.	Irrelevant intervention
Koehler, B. E., Richter, K. M., Youngblood, L., Cohen, B. A., Prengler, I. D., Cheng, D., & Masica, A. L. (2009). Reduction of 30-day postdischarge hospital readmission or emergency department (ED) visit rates in high-risk elderly medical patients through delivery of a targeted care bundle. <i>J Hosp Med</i> , 4(4), 211-218.	Irrelevant intervention
Kohn, D., Sinoff, G., Strulov, A., Ciechanover, M., & Wei, J. Y. (1991). Long-term follow-up of patients aged 75 years and older admitted to an acute care hospital in Israel. <i>Aging - Clinical and Experimental Research</i> , 3(3), 279-285.	Irrelevant studiedesign
Komajda, M., Hanon, O., Hochadel, M., Lopez-Sendon, J. L., Follath, F., Ponikowski, P., . . . Nieminen, M. (2009). Contemporary management of octogenarians hospitalized for heart failure in Europe: Euro Heart Failure Survey II. <i>Eur Heart J</i> , 30(4), 478-486.	Irrelevant population
Kravitz Richard, L., Reuben David, B., Davis James, W., Mitchell, A., & et al. (1994). Geriatric home assessment after hospital discharge. <i>Journal of the American Geriatrics Society</i> , 42(12), 1229-1234.	Irrelevant intervention
Kroemer, D. J., Bloor, G., & Fiebig, J. (2004). Acute transition alliance: rehabilitation at the acute/aged care interface. <i>Australian Health Review</i> , 28(3), 266-274.	Övrigt
LaMantia MA, Platts-Mills TF, Biese K, Khandelwal C, Forbach C, Cairns CB, et al. Predicting hospital admission and returns to the emergency department for elderly patients. <i>Academic Emergency Medicine</i> . 2010;17(3):252-9.	Irrelevant studiedesign
Lamb, S. E. (2009). Multidisciplinary assessment of elderly people with a history of multiple falls reduces the risk of further falls. <i>Aust J Physiother</i> , 55(2), 139.	Irrelevant population
Landi F, Gambassi G, Pola R, Tabaccanti S, Cavinato T, Carbonin P, et al. Impact of integrated home care services on hospital use. <i>Journal of the American Geriatrics Society</i> . 1999;47(12):1430-4.	Irrelevant population
Landi, F., Zuccala, G., Bernabei, R., Cocchi, A., Manigrasso, L., Tafani, A., . . . Carbonin, P. (1997). Physiotherapy and occupational therapy: a geriatric experience in the acute care hospital. <i>American Journal of Physical Medicine &amp; Rehabilitation</i> , 76(1), 38-42.	Irrelevant intervention

Lang P, Heitz D, Hedelin G, Drame M, Jovenin N, Ankri J, et al. Early markers of prolonged hospital stays in older people: a prospective, multicenter study of 908 inpatients in French acute hospitals [corrected] [published erratum appears in J AM GERIATR SOC 2006 Sep;54(9):1479-80]. <i>Journal of the American Geriatrics Society</i> . 2006;54(7):1031-9.	Irrelevant studiedesign
Lang PO, Meyer N, Heitz D, Drame M, Jovenin N, Ankri J, et al. Loss of independence in Katz's ADL ability in connection with an acute hospitalization: early clinical markers in French older people. <i>Eur J Epidemiol</i> . 2007;22(9):621-30.	Irrelevant studiedesign
Lang PO, Zekry D, Michel JP, Drame M, Novella JL, Jolly D, et al. Early markers of prolonged hospital stay in demented inpatients: a multicentre and prospective study. <i>J Nutr Health Aging</i> . 2010;14(2):141-7.	Irrelevant studiedesign
Lang, P. O., Dramee, M., & Michel, J. P. (2011). Interdisciplinary geriatric and psychiatric care reduces potentially inappropriate prescribing in-hospital: Interventional study in 150 acutely ill elderly patients with mental and somatic co-morbid conditions. <i>European Geriatric Medicine</i> , 2, S173.	Dubbelpublicering
Lang, P. O., Vogt-Ferrier, N., Hasso, Y., Le, S., Drame, M., Zekry, D., . . . Michel, J. P. (2011). Interdisciplinary Geriatric and Psychiatric Care Reduces Potentially Inappropriate Prescribing in the Hospital: Interventional Study in 150 Acutely Ill Elderly Patients with Mental and Somatic Comorbid Conditions. <i>J Am Med Dir Assoc</i> , 13, 406 e401-407. doi: 10.1016/j.jamda.2011.03.008. Epub 2011 May 18.	Irrelevant studiedesign
Lang, P. O., Vogt-Ferrier, N., Hasso, Y., Le, S., Drame, M., Zekry, D., . . . Michel, J. P. (2012). Interdisciplinary geriatric and psychiatric care reduces potentially inappropriate prescribing in the hospital: interventional study in 150 acutely ill elderly patients with mental and somatic comorbid conditions. <i>J Am Med Dir Assoc</i> , 13, 406 e401-407. doi: 10.1016/j.jamda.2011.03.008. Epub 2011 May 18.	Irrelevant studiedesign
Lapane, K. L., Hughes, C. M., Daiello, L. A., Cameron, K. A., & Feinberg, J. (2011). Effect of a pharmacist-led multicomponent intervention focusing on the medication monitoring phase to prevent potential adverse drug events in nursing homes. <i>J Am Geriatr Soc</i> , 59(7), 1238-1245. doi: 10.1111/j.1532-5415.2011.03418.x. Epub 2011 Jun 7.	Irrelevant intervention
Laramee, A. S., Levinsky, S. K., Sargent, J., Ross, R., & Callas, P. (2003). Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. <i>Arch Intern Med</i> , 163(7), 809-817.	Irrelevant population
Launay, C., de, D., Hureaux-Huynh, R., Annweiler, C., & Beauchet, O. (2012). Mobile geriatric team and length of hospital stay among older inpatients: a case-control pilot study. <i>J Am Geriatr Soc</i> , 60, 1593-1594. doi: 10.1111/j.1532-5415.2012.04088.x.	Irrelevant studiedesign
Laursen, S. O., Henriksen, I. O., Dons, U., Jacobsen, B., & Gundertofte, L. (1995). [Intensive rehabilitation after apoplexy--a controlled pilot study]. <i>Ugeskr Laeger</i> , 157(14), 1996-1999.	Irrelevant population
Ledesert B, Lombraïl P, Yeni P, Carbon C, Brodin M. The impact of a comprehensive multi-dimensional geriatric assessment programme on duration of stay in a French acute medical ward. <i>Age and Ageing</i> . 1994;23(3):223-7.	Irrelevant studiedesign
Lee W, Peng L, Cheng Y, Liu C, Chen L, Yu H. Effectiveness of short-term interdisciplinary intervention on postacute patients in taiwan. <i>Journal of the American Medical Directors Association</i> . 2011;12(1):29-32.	Irrelevant studiedesign
Lee V, Ross B, Tracy B. Functional assessment of older adults in an emergency department. <i>Can J Occup Ther</i> . 2001;68(2):121-9.	Irrelevant studiedesign
Lee, J. S., Hurley, M. J., Carew, D., Fisher, R., Kiss, A., & Drummond, N. (2007). A Randomized Clinical Trial to Assess the Impact on an Emergency Response System on Anxiety and Health Care Use among Older Emergency Patients after a Fall. <i>Academic Emergency Medicine</i> , 14(4), 301-308.	Irrelevant intervention

Lee, J. S., Schwindt, G., Langevin, M., Moghabghab, R., Alibhai, S. M. H., Kiss, A., & Naglie, G. (2008). Validation of the Triage Risk Stratification Tool to identify older persons at risk for hospital admission and returning to the emergency department. <i>Journal of the American Geriatrics Society</i> , 56(11), 2112-2117.	Övrigt
Lee, K. H., Hui, K. P., Lim, T. K., & Tan, W. C. (1993). Acute physiology and chronic health evaluation (APACHE II) scoring in the Medical Intensive Care Unit, National University Hospital, Singapore. <i>Singapore Med J</i> , 34(1), 41-44.	Övrigt
Lee, L. (2008). A multifactorial intervention did not prevent falls or fractures in elderly patients during short hospital stays. <i>Evidence Based Nursing</i> , 11(4), 120-120.	Irrelevant studiedesign
Lee, W. J., Cheng, Y. Y., Liu, C. Y., Peng, L. N., Liu, L. K., & Chen, L. K. (2012). Dose-dependent effect of rehabilitation in functional recovery of older patients in the post-acute care unit. <i>Archives of Gerontology and Geriatrics</i> , 54, e290-e293.	Irrelevant studiedesign
Lee, Y. S., Lin, C. S., Jseng, Y. H., Luo, T. W., Hung, P. J., Wu, M. C., & Tang, Y. J. (2012). Predictive factors for patients discharged after participating in a post-acute care program. <i>Journal of Clinical Gerontology and Geriatrics</i> , 3, 25-28.	Irrelevant studiedesign
Lees L. A framework to promote the holistic assessment of older people in emergency care. <i>Nurs Older People</i> . 2005;16(10):16-8, 20.	Irrelevant studiedesign
Leff, B., Burton, L., Mader, S. L., Naughton, B., Burl, J., Inouye, S. K. Burton, J. R. (2005). Improving patient care. Hospital at home: feasibility and outcomes of a program to provide hospital-level care at home for acutely ill older patients. <i>Annals of Internal Medicine</i> , 143(11), 798.	Irrelevant intervention
Leff, B., Reider, L., Frick, K. D., Scharfstein, D. O., Boyd, C. M., Frey, K., . . . Boult, C. (2009). Guided care and the cost of complex healthcare: a preliminary report. <i>American Journal of Managed Care</i> , 15(8), 555-559.	Irrelevant studiedesign
Lefton E, Bonstelle S, Frengley JD. Success with an inpatient geriatric unit: a controlled study of outcome and follow-up. <i>J Am Geriatr Soc</i> . 1983;31(3):149-55.	Irrelevant studiedesign
Legrain, S., Tubach, F., Bonnet-Zamponi, D., Lemaire, A., Aquino, J. P., Paillaud, E., . . . Lacaille, S. (2011). A new multimodal geriatric discharge-planning intervention to prevent emergency visits and rehospitalizations of older adults: the optimization of medication in AGEd multicenter randomized controlled trial. <i>J Am Geriatr Soc</i> , 59, 2017-2028. doi: 10.1111/j.1532-5415.2011.03628.x. Epub 2011 Sep 27.	Irrelevant intervention
Lenze Eric, J., Host Helen, H., Hildebrand Mary, W., Morrow-Howell, N., Carpenter, B., Freedland Kenneth, E., . . . Binder Ellen, F. (2012). Enhanced Medical Rehabilitation Increases Therapy Intensity and Engagement and Improves Functional Outcomes in Postacute Rehabilitation of Older Adults: A Randomized-Controlled Trial. <i>Journal of the American Medical Directors Association</i> , 13, 708-712.	Irrelevant studiedesign
Leslie, D. L., Zhang, Y., Bogardus, S. T., Holford, T. R., Leo-Summers, L. S., & Inouye, S. K. (2005). Consequences of preventing delirium in hospitalized older adults on nursing home costs. <i>J Am Geriatr Soc</i> , 53(3), 405-409.	Övrigt
Leung, A. C., Liu, C., Chow, N. W., & Chi, I. (2004). Cost-benefit analysis of a case management project for the community-dwelling frail elderly in Hong Kong. <i>Journal of Applied Gerontology</i> , 23(1), 70-85.	Irrelevant intervention
Leung, L. P., & Fan, K. L. (2008). Who should be admitted to hospital? Evaluation of a screening tool. <i>Hong Kong Med J</i> , 14(4), 273-277.	Övrigt

Leventhal, M. E., Denhaerynck, K., Brunner-La, R., H, P., Burnand, B., Conca-Zeller, A. De, G. (2011). Swiss Interdisciplinary Management Programme for Heart Failure (SWIM-HF): a randomised controlled trial study of an outpatient inter-professional management programme for heart failure patients in Switzerland. <i>Swiss Med Wkly</i> , 141, w13171. doi: 10.4414/smw.2011.13171.	Irrelevant population
Li, M., Porter, E., Lam, R., & Jassal, S. V. (2007). Quality Improvement Through the Introduction of Interdisciplinary Geriatric Hemodialysis Rehabilitation Care. <i>American Journal of Kidney Diseases</i> , 50(1), 90-97.	Irrelevant intervention
Lightbody E, Baldwin R. Inpatient geriatric evaluation and management did not reduce mortality but reduced functional decline. <i>Evidence Based Mental Health</i> . 2002;5(4):109-.	Övrigt
Lightbody E, Watkins C, Leathley M, Sharma A, Lye M. Evaluation of a nurse-led falls prevention programme versus usual care: a randomized controlled trial. <i>Age Ageing</i> . 2002;31(3):203-10.	Irrelevant intervention
Lim, W. K., Lambert, S. F., & Gray, L. C. (2003). Effectiveness of case management and post-acute services in older people after hospital discharge. <i>Med J Aust</i> , 178(6), 262-266.	Irrelevant intervention
Lin, A., Cavendish, J., Boren, D., Ofstad, T., & Seidensticker, D. (2008). A pilot study: Reports of benefits from a 6-month, multidisciplinary, shared medical appointment approach for heart failure patients. <i>Military Medicine</i> , 173(12), 1210-1213.	Irrelevant population
Lin, J. S., Whitlock, E. P., Eckstrom, E., Fu, R., Perdue, L. A., Beil, T. L., & Leipzig, R. M. (2012). <i>Challenges in Synthesizing and Interpreting the Evidence From a Systematic Review of Multifactorial Interventions to Prevent Functional Decline in Older Adults</i> . Rockville MD.	Övrigt
Lindberg, M. C., & Sullivan, G. M. (1996). Effects of an inpatient geriatrics rotation on internal medicine residents' knowledge and attitudes. <i>J Gen Intern Med</i> , 11(7), 397-400.	Irrelevant intervention
Lisby, M., Thomsen, A., Nielsen, L. P., Lyhne, N. M., Breum-Leer, C., Fredberg, U., . . . Brock, B. (2010). The effect of systematic medication review in elderly patients admitted to an acute ward of internal medicine. <i>Basic Clin Pharmacol Toxicol</i> , 106(5), 422-427.	Irrelevant intervention
Lo, S. H., Chan, C. Y., Chan, C. H., Sze, W. K., Yuen, K. K., Wong, C. S., . . . Tung, Y. (2009). The implementation of an end-of-life integrated care pathway in a Chinese population. <i>Int J Palliat Nurs</i> , 15(8), 384-388.	Övrigt
Long, M. J., & Marshall, B. S. (1999). Case management and the cost of care in the last month of life: evidence from one managed care setting. <i>Health Care Manage Rev</i> , 24(4), 45-53.	Irrelevant intervention
Lorensen, M., & Eriksen, R. (2003). The effect of systematic assessment in rehabilitation on self-care capability, length of hospital stay, patient satisfaction, and interdisciplinary collaboration. <i>Self-Care, Dependent-Care &amp; Nursing</i> , 11(3), 4-18.	Irrelevant studiedesign
Lu JH, Chan DK, Ong B, Shen Q, Reuten S, Ko A. Management and outcomes of delirium in a secured, co-located geriatric and psychogeriatric unit. <i>J Am Geriatr Soc</i> . 2009;57(9):1725-7.	Irrelevant studiedesign
Luk, J. K. H., & Chan, C. F. (2011). Rehabilitation outcomes of older patients at 6 months follow-up after discharged from a geriatric day hospital (GDH). <i>Archives of Gerontology and Geriatrics</i> , 52(3), 327-330.	Irrelevant intervention
Luk, J. K., Kwok, T., & Woo, J. (1999). Geriatric screening in acute care wards--a novel method of providing care to elderly patients. <i>Hong Kong Med J</i> , 5(1), 34-38.	Irrelevant studiedesign
Lundstrom, M., Edlund, A., Karlsson, S., Brannstrom, B., Bucht, G., & Gustafson, Y. (2005). A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. <i>Journal of the American Geriatrics Society</i> , 53(4), 622-628.	Irrelevant studiedesign

Lushbough, R. S., Priddy, J. M., Sewell, H. H., Lovett, S. B., & Jones, T. C. (1988). The effectiveness of an occupational therapy program in an inpatient geropsychiatric setting. <i>Physical &amp; Occupational Therapy in Geriatrics</i> , 6(2), 63-73.	Irrelevant intervention
Maaravi, Y., & Stessman, J. (2008). The geriatric emergency department... Hwang U, Morrison S. The geriatric emergency department. <i>J Am Geriatr Soc</i> 2007;55:1873-6. <i>Journal of the American Geriatrics Society</i> , 56(8), 1579-1579.	Övrigt
Macdonald, A. J., Mann, A. H., Jenkins, R., Richard, L., Godlove, C., & Rodwell, G. (1982). An attempt to determine the impact of four types of care upon the elderly in London by the study of matched groups. <i>Psychol Med</i> , 12(1), 193-200.	Irrelevant studiedesign
Marengoni A, Cossi S, De M, Ghisla MK, Calabrese PA, Zanolini G, et al. Adverse outcomes in older hospitalized patients: The role of multidimensional geriatric assessment. <i>Aging - Clinical and Experimental Research</i> . 2003;15(1):32-7.	Irrelevant studiedesign
Margitic SE, Inouye SK, Thomas JL, Cassel CK, Regenstreif DI, Kowal J. Hospital Outcomes Project for the Elderly (HOPE): Rationale and design for a prospective pooled analysis. <i>Journal of the American Geriatrics Society</i> . 1993;41(3):258-67.	Irrelevant studiedesign
Martinez-Selles, M., Datino, T., & Bueno, H. (2006). Coronary care unit admission of very old patients with acute myocardial infarction. <i>Heart</i> , 92(4), 549-550.	Irrelevant population
Mason, S., O'Keeffe, C., Knowles, E., Bradburn, M., Campbell, M., Coleman, P., . . . Patterson, M. (2012). A pragmatic quasi-experimental multi-site community intervention trial evaluating the impact of Emergency Care Practitioners in different UK health settings on patient pathways (NEECap Trial). <i>Emergency Medicine Journal</i> , 29, 47-53.	Irrelevant intervention
Mateev A, Gaspoz JM, Borst F, Waldvogel F, Weber D. Use of a short-form screening procedure to detect unrecognized functional disability in the hospitalized elderly. <i>J Clin Epidemiol</i> . 1998;51(4):309-14.	Irrelevant studiedesign
Mathew, L. J., Gutsch, H. M., Hackney, N. W., & Munsat, E. M. (1994). Promoting quality and cost-effective care to geropsychiatric patients. <i>Issues in Mental Health Nursing</i> , 15(2), 169-185.	Irrelevant intervention
Mazzola P, De F, Castoldi G, Galetti P, Zatti G, Annoni G. A comparison between two co-managed geriatric programmes for hip fractured elderly patients. <i>Aging Clin Exp Res</i> . 2010.	Övrigt
McCloskey, R. (2004). Functional and self-efficacy changes of patients admitted to a geriatric rehabilitation unit. <i>Journal of Advanced Nursing</i> , 46(2), 186-193.	Irrelevant studiedesign
McCormack B. An acute care unit for elderly medical patients did not increase hospital costs [commentary on Covinsky KE, King JT Jr, Quinn LM, et al. Do acute care for elders units increase hospital costs? A cost analysis using the hospital perspective. <i>J AM GERIATR SOC</i> 1997;45(6):729-34]. <i>Evidence Based Nursing</i> . 1998;1(1):26-.	Övrigt
McCusker J, Dendukuri N, Tousignant P, Verdon J, Belzile E. Rapid two-stage emergency department intervention for seniors: impact on continuity of care. <i>Academic Emergency Medicine</i> . 2003;10(3):233-43.	Irrelevant studiedesign
McCusker J, Jacobs P, Dendukuri N, Latimer E, Tousignant P, Verdon J. Cost-effectiveness of a brief two-stage emergency department intervention for high-risk elders: results of a quasi-randomized controlled trial. <i>Ann Emerg Med</i> . 2003;41(1):45-56.	Irrelevant intervention
McCusker J, Verdon J, Tousignant P, de C, L P, Dendukuri N, et al. Rapid emergency department intervention for older people reduces risk of functional decline: results of a multicenter randomized trial. <i>J Am Geriatr Soc</i> . 2001;49(10):1272-81.	Irrelevant intervention
McCusker, J., & Verdon, J. (2006). Do geriatric interventions reduce emergency department visits? A systematic review. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 61(1), 53-62.	Övrigt

McCusker, J., Dendukuri, N., Tousignant, P., Verdon, J., De, C., L, P., & Belzile, E. (2003). Rapid two-stage emergency department intervention for seniors: Impact on continuity of care. <i>Academic Emergency Medicine</i> , 10(3), 233-243.	Irrelevant intervention
McCusker, J., Verdon, J., Tousignant, P., Poulin de, C., Louise, Dendukuri, N., & Belzile, E. (2001). Rapid emergency department intervention for older people reduces risk of functional decline: Results of a multicenter randomized trial. <i>Journal of the American Geriatrics Society</i> , 49(10), 1272-1281.	Irrelevant intervention
Meissner P, Andolsek K, Mears PA, Fletcher B. Maximizing the functional status of geriatric patients in an acute community hospital setting. <i>Gerontologist</i> . 1989;29(4):524-8.	Irrelevant studiedesign
Meldon, S. W., Mion, L. C., Palmer, R. M., Drew, B. L., Connor, J. T., Lewicki, L. J., . . . Emerman, C. L. (2003). A brief risk-stratification tool to predict repeat emergency department visits and hospitalizations in older patients discharged from the emergency department. <i>Academic Emergency Medicine</i> , 10(3), 224-232.	Irrelevant intervention
Melin, A. L., Hakansson, S., & Bygren, L. O. (1993). The cost-effectiveness of rehabilitation in the home: a study of Swedish elderly. <i>Am J Public Health</i> , 83(3), 356-362.	Irrelevant intervention
Melis, R. J. F., Adang, E., Teerenstra, S., van, E., M, I. J., Wimo, A., . . . M, G. M. (2008). Cost-effectiveness of a multidisciplinary intervention model for community-dwelling frail older people. <i>Journals of Gerontology Series A: Biological Sciences &amp; Medical Sciences</i> , 63A(3), 275-282.	Irrelevant population
Merkert, J., Butz, S., Nieczaj, R., Steinhagen-Thiessen, E., & Eckardt, R. (2011). Combined whole body vibration and balance training using Vibrosphere(R): improvement of trunk stability, muscle tone, and postural control in stroke patients during early geriatric rehabilitation. <i>Z Gerontol Geriatr</i> , 44(4), 256-261.	Irrelevant population
Michael, R., Wheeler, B., Wichmann, H., Horner, B., & Downie, J. (2005). The healthy ageing unit: a comparative controlled intervention. <i>Journal of the Australasian Rehabilitation Nurses' Association (JARNA)</i> , 8(2), 9-15.	Irrelevant studiedesign
Michael, R., Wichmann, H., Wheeler, B., Horner, B., & Downie, J. (2004). A multidisciplinary model of transitional rehabilitation in acute aged care. <i>Journal of the Australasian Rehabilitation Nurses' Association (JARNA)</i> , 7(4), 10-16.	Övrigt
Michael, R., Wichmann, H., Wheeler, B., Horner, B., & Downie, J. (2005). The healthy ageing unit: beyond discharge. <i>Journal of the Australasian Rehabilitation Nurses' Association (JARNA)</i> , 8(4), 8-16.	Irrelevant studiedesign
Milisen K, Foreman MD, Abraham IL, De G, Godderis J, Vandermeulen E, et al. A nurse-led interdisciplinary intervention program for delirium in elderly hip-fracture patients. <i>Journal of the American Geriatrics Society</i> . 2001;49(5):523-32.	Irrelevant population
Miller DK, Lewis LM, Nork MJ, Morley JE. Controlled trial of a geriatric case-finding and liaison service in an emergency department. <i>J Am Geriatr Soc</i> . 1996;44(5):513-20.	Irrelevant studiedesign
Miller, P., Gladman, J. R. F., Cunliffe, A. L., Husbands, S. L., Dewey, M. E., & Harwood, R. H. (2005). Economic analysis of an early discharge rehabilitation service for older people. <i>Age &amp; Ageing</i> , 34(3), 274-280.	Irrelevant intervention
Mion LC, Palmer RM, Meldon SW, Bass DM, Singer ME, Payne SM, et al. Case finding and referral model for emergency department elders: a randomized clinical trial. <i>Ann Emerg Med</i> . 2003;41(1):57-68.	Irrelevant intervention
Miralles, R., Sabartes, O., Ferrer, M., Esperanza, A., Llorach, I., Garcia-Palleiro, P., & Cervera, A. M. (2003). Development and validation of an instrument to predict probability of home discharge from a geriatric convalescence unit in Spain. <i>J Am Geriatr Soc</i> , 51(2), 252-257.	Irrelevant studiedesign

Montuclard, L., Garrouste-Orgeas, M., Timsit, J., Misset, B., De, J., & Carlet, J. (2000). Outcome, functional autonomy, and quality of life of elderly patients with a long-term intensive care unit stay. <i>Critical Care Medicine</i> , 28(10), 3389-3395.	Irrelevant intervention
Mor, V., Intrator, O., Fries, B. E., Phillips, C., Teno, J., Hiris, J., . . . Morris, J. (1997). Changes in hospitalization associated with introducing the resident assessment instrument. <i>Journal of the American Geriatrics Society</i> , 45(8), 1002-1010.	Irrelevant population
Mudge, A. M., Giebel, A. J., & Cutler, A. J. (2008). Exercising body and mind: an integrated approach to functional independence in hospitalized older people. <i>J Am Geriatr Soc</i> , 56(4), 630-635.	Irrelevant intervention
Mudge, A., Laracy, S., Richter, K., & Denaro, C. (2006). Controlled trial of multidisciplinary care teams for acutely ill medical inpatients: Enhanced multidisciplinary care. <i>Internal Medicine Journal</i> , 36(9), 558-563.	Irrelevant population
Munin, M. C., Begley, A., Skidmore, E. R., & Lenze, E. J. (2006). Influence of rehabilitation site on hip fracture recovery in community-dwelling subjects at 6-month follow-up. <i>Archives of Physical Medicine &amp; Rehabilitation</i> , 87(7), 1004-1006.	Irrelevant population
Murphy TH, Labonte P, Klock M, Houser L. Falls prevention for elders in acute care: an evidence-based nursing practice initiative. <i>Crit Care Nurs Q</i> . 2008;31(1):33-9.	Irrelevant studiedesign
Myint, P. K., Kamath, A. V., Vowler, S. L., Maisey, D. N., & Harrison, B. D. W. (2005). The CURB (confusion, urea, respiratory rate and blood pressure) criteria in community-acquired pneumonia (CAP) in hospitalised elderly patients aged 65 years and over: a prospective observational cohort study. <i>Age &amp; Ageing</i> , 34(1), 75-77.	Irrelevant intervention
Naglie, G., Tansey, C., Kirkland, J. L., Ogilvie-Harris, D. J., Detsky, A. S., Etchells, E., . . . Goldlist, B. (2002). Interdisciplinary inpatient care for elderly people with hip fracture: A randomized controlled trial. <i>Canadian Medical Association Journal</i> , 167(1), 25-32.	Irrelevant population
Nankhonya JM. Screening for social needs in accident and emergency departments. <i>J Accid Emerg Med</i> . 1994;11(2):135-6.	Irrelevant studiedesign
Naughton BJ, Saltzman S, Ramadan F, Chadha N, Priore R, Mylotte JM. A multifactorial intervention to reduce prevalence of delirium and shorten hospital length of stay. <i>J Am Geriatr Soc</i> . 2005;53(1):18-23.	Irrelevant studiedesign
Naylor, M. D. (1990). Comprehensive discharge planning for hospitalized elderly: a pilot study. <i>Nurs Res</i> , 39(3), 156-161.	Irrelevant population
Naylor, M. D. (1990). Comprehensive discharge planning for the elderly. <i>Res Nurs Health</i> , 13(5), 327-347.	Irrelevanta effektmått
Naylor, M. D. (2000). A decade of transitional care research with vulnerable elders. <i>Journal of Cardiovascular Nursing</i> , 14(3), 1.	Övrigt
Naylor, M. D., Brooten, D. A., Campbell, R. L., Maislin, G., McCauley, K. M., & Schwartz, J. S. (2004). Transitional care of older adults hospitalized with heart failure: a randomized, controlled trial [corrected] [published erratum appears in J AM GERIATR SOC 2004 Jul;52(7):1228]. <i>Journal of the American Geriatrics Society</i> , 52(5), 675-684.	Irrelevant population
Naylor, M. D., Brooten, D., Campbell, R., Jacobsen, B. S., Mezey, M. D., Pauly, M. V., & Schwartz, J. S. (1999). Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. <i>JAMA</i> , 281(7), 613-620.	Irrelevant intervention
Nazareth, I., Burton, A., Shulman, S., Smith, P., Haines, A., & Timberal, H. (2001). A pharmacy discharge plan for hospitalized elderly patients--a randomized controlled trial. <i>Age Ageing</i> , 30(1), 33-40.	Irrelevant intervention
Newcomer, R., Kang, T., & Graham, C. (2006). Outcomes in a nursing home transition case-management program targeting new admissions. <i>Gerontologist</i> , 46(3), 385-390.	Irrelevant intervention

Ngian VJJ, Ong BS, O'Rourke F, Nguyen HV, Chan DKY. Review of a rapid geriatric medical assessment model based in emergency department. <i>Age &amp; Ageing</i> . 2008;37(6):696-9.	Irrelevant studiedesign
NICHE (Nurses Improving Care to the Hospitalized Elderly) project helps improve geriatric outcomes. <i>Patient Focus Care Satisf</i> . 1998;6(1):10-1.	Irrelevant studiedesign
Nikolaus T, Bach M, Oster P, Schlierf G. Prospective value of self-report and performance-based tests of functional status for 18-month outcomes in elderly patients. <i>Aging (Milano)</i> . 1996;8(4):271-6.	Irrelevant studiedesign
Nikolaus T, Bach M. Preventing falls in community-dwelling frail older people using a home intervention team (HIT): results from the Randomized Falls-HIT Trial. <i>Journal of the American Geriatrics Society</i> . 2003;51(3):300-5.	Irrelevant intervention
Nikolaus T, Specht-Leible N, Bach M, Oster P, Schuerf G. A randomised trial of Comprehensive geriatric assessment and home intervention in the care of hospitalised patients. <i>Age &amp; Ageing</i> 1999;28:543-50.	Irrelevant intervention
Nikolaus T, Specht-Leible N, Bach M, et al. (2000) Comprehensive inhospital geriatric assessment plus an interdisciplinary home intervention after discharge reduced length of subsequent readmissions and improved functioning. <i>Evidence Based Nursing</i> . 3(83).	Övrigt
Noro, A., Poss, J. W., Hirdes, J. P., Finne-Soveri, H., Ljunggren, G., Bjornsson, J., . . . Jonsson, P. V. (2011). Method for Assigning Priority Levels in Acute Care (MAPLe-AC) predicts outcomes of acute hospital care of older persons--a cross-national validation. <i>BMC Med Inform Decis Mak</i> , 11, 39.	Irrelevant intervention
O'Donnell JC, Toseland RW. Does geriatric evaluation and management improve the health behavior of older veterans in psychological distress? <i>J Aging Health</i> . 1997;9(4):473-97.	Irrelevant population
O'Grady, S., Fairbrother, G., & Farrington, C. (1996). Matching needs to services: the quick response. <i>Australian Health Review</i> , 19(4), 100-112.	Övrigt
Olofsson, B., Stenvall, M., Lundström, M., Svensson, O., & Gustafson, Y. (2007). Malnutrition in hip fracture patients: An intervention study. <i>Journal of Clinical Nursing</i> , 16(11), 2027-2038.	Irrelevant population
Olsson, L. E., Hansson, E., Ekman, I., & Karlsson, J. (2009). A cost-effectiveness study of a patient-centred integrated care pathway. <i>J Adv Nurs</i> , 65(8), 1626-1635.	Irrelevant studiedesign
Olsson, L., Hansson, E., Ekman, I., & Karlsson, J. (2009). A cost-effectiveness study of a patient-centred integrated care pathway. <i>Journal of Advanced Nursing</i> , 65(8), 1626-1635.	Irrelevant population
Olsson, L., Karlsson, J., & Ekman, I. (2007). Effects of nursing interventions within an integrated care pathway for patients with hip fracture. <i>Journal of Advanced Nursing</i> , 58(2), 116-125.	Irrelevant population
Onder, G., Liperoti, R., Soldato, M., Carpenter, I., Steel, K., Bernabei, R., & Landi, F. (2007). Case management and risk of nursing home admission for older adults in home care: results of the AgeD in HOme Care Study. <i>J Am Geriatr Soc</i> , 55(3), 439-444.	Irrelevant population
Opasich, C., Patrignani, A., Mazza, A., Gualco, A., Cobelli, F., & Pinna, G. D. (2010). An elderly-centered, personalized, physiotherapy program early after cardiac surgery. <i>Eur J Cardiovasc Prev Rehabil</i> , 17(5), 582-587.	Irrelevant population
O'Reilly, J., Lowson, K., Green, J., Young, J. B., & Forster, A. (2008). Post-acute care for older people in community hospitals--a cost-effectiveness analysis within a multi-centre randomised controlled trial. <i>Age &amp; Ageing</i> , 37(5), 513-520.	Irrelevant intervention
O'Reilly, J., Lowson, K., Young, J., Forster, A., Green, J., & Small, N. (2006). A cost effectiveness analysis within a randomised controlled trial of post-acute care of older people in a community hospital. <i>BMJ: British Medical Journal (International Edition)</i> , 333(7561), 228-231.	Irrelevant intervention

Ornstein, K., Smith Kristofer, L., Foer Dinah, H., Lopez-Cantor Maria, T., & Soriano, T. (2011). To the hospital and back home again: a nurse practitioner-based transitional care program for hospitalized homebound people. <i>Journal of the American Geriatrics Society</i> , 59(3), 544-551.	Irrelevant population
Ouslander, J. G., Griffiths, P. C., McConnell, E., Riolo, L., Kutner, M., & Schnelle, J. (2005). Functional incidental training: a randomized, controlled, crossover trial in Veterans Affairs nursing homes. <i>Journal of the American Geriatrics Society</i> , 53(7), 1091-1100.	Irrelevant population
Palleschi L, De A, Salani B, Fimognari FL, Marsili A, Pierantozzi A, et al. Functional recovery of elderly patients hospitalized in geriatric and general medicine units. the PROgetto DImissioni in GEriatria study. <i>Journal of the American Geriatrics Society</i> . 2011;59(2):193-9.	Irrelevant studiedesign
Palmer RM, Counsell S, Landefeld CS. Clinical intervention trials: the ACE unit. <i>Clinics in Geriatric Medicine</i> . 1998;14(4):831-49.	Övrigt
Palnum, K. D., Petersen, P., Sorensen, H. T., Ingeman, A., Mainz, J., Bartels, P., & Johnsen, S. P. (2008). Older patients with acute stroke in Denmark: quality of care and short-term mortality. A nationwide follow-up study. <i>Age &amp; Ageing</i> , 37(1), 90-95.	Irrelevant population
Panella, M., Marchisio, S., Barbieri, A., & Di, S. (2008). A cluster randomized trial to assess the impact of clinical pathways for patients with stroke: rationale and design of the Clinical Pathways for Effective and Appropriate Care Study [NCT00673491]. <i>BMC Health Serv Res</i> , 8, 223.	Irrelevant population
Paquet, N., Desrosiers, J., Demers, L., & Robichaud, L. (2009). Predictors of daily mobility skills 6 months post-discharge from acute care or rehabilitation in older adults with stroke living at home. <i>Disability and Rehabilitation: An International, Multidisciplinary Journal</i> , 31(15), 1267-1274.	Irrelevant population
Partridge RA, Shapiro MJ, Micalone M, Jenouri I, Woolard RH, Gifford DR. Rapid computerized cognitive assessment: performance of highly functional elder patients with minor traumatic injury in the emergency department. <i>Acad Emerg Med</i> . 1999;6(7):758-60.	Irrelevant studiedesign
Pascual, C. R., Galan, E. P., Guerrero, J. L., Colino, R. M., Soler, P. A., Calvo, M. H., . . . Rodriguez-Artalejo, F. (2011). Rationale and methods of the multicenter randomised trial of a heart failure management programme among geriatric patients (HF-Geriatrics). <i>BMC Public Health</i> , 11, 627.	Irrelevant population
Peleg, R., Press, Y., Asher, M., Pugachev, T., Glicenstein, H., Lederman, M., & Biderman, A. (2008). An intervention program to reduce the number of hospitalizations of elderly patients in a primary care clinic. <i>BMC Health Serv Res</i> , 8, 36.	Irrelevant population
Perez-Zepeda MU, Gutierrez-Robledo LM, Sanchez-Garcia S, Juarez-Cedillo T, Gonzalez JJ, Franco-Marina F, et al. Comparison of a geriatric unit with a general ward in Mexican elders. <i>Arch Gerontol Geriatr</i> . 2011.	Irrelevant studiedesign
Perez-Zepeda, M. U., Gutierrez-Robledo, L. M., Sanchez-Garcia, S., Juarez-Cedillo, T., Gonzalez, J. J., Franco-Marina, F., & Garcia-Pena, C. (2012). Comparison of a geriatric unit with a general ward in Mexican elders. <i>Arch Gerontol Geriatr</i> , 54, e370-375. doi: 10.1016/j.archger.2011.05.028. Epub 2011 Jul 22.	Irrelevant population
Phillips, E. M., Abrandt, B. L., Cesta, T., & Gallucci, M. A. (1999). Rehabilitation after hip fracture. <i>Topics in Geriatric Rehabilitation</i> , 15(1), 56-65.	Irrelevant population
Pitkälä Kaisu, H., Laurila Jouko, V., Strandberg Timo, E., & Tilvis Reijo, S. (2006). Multicomponent Geriatric Intervention for Elderly Inpatients With Delirium: A Randomized, Controlled Trial. <i>The Journals of Gerontology: Series A: Biological Sciences and Medical Sciences</i> , 61A(2), 176-181.	Irrelevant intervention

Popplewell PY, Henschke PJ. What is the value of a Geriatric Assessment Unit in a teaching hospital? A comparative study of the management of elderly inpatients. <i>Aust Health Rev</i> . 1983;6(2):23-5.	Irrelevant studiedesign
Pound, P., Sabin, C., & Ebrahim, S. (1999). Observing the process of care: a stroke unit, elderly care unit and general medical ward compared. <i>Age &amp; Ageing</i> , 28(5), 433-440.	Irrelevant studiedesign
Pugh, L. C., Tringali, R. A., Boehmer, J., Blaha, C., Kruger, N. R., Capauna, T. A., . . . Xie, S. (1999). Partners in care: a model of collaboration. <i>Holistic Nursing Practice</i> , 13(2), 61-65.	Irrelevant population
Pulignano, G., Del, S., Di, L., Tarantini, L., Cioffi, G., Gregori, D., . . . Minardi, G. (2010). Usefulness of frailty profile for targeting older heart failure patients in disease management programs: A cost-effectiveness, pilot study. <i>Journal of Cardiovascular Medicine</i> , 11(10), 739-747.	Irrelevant population
Randomised controlled trial of a nurse-led falls prevention programme [8]. (2003). <i>Age and Ageing</i> , 32(3), 361-362.	Övrigt
Resnick, B. (1998). Efficacy beliefs in geriatric rehabilitation. <i>J Gerontol Nurs</i> , 24(7), 34-44.	Irrelevant population
Resnick, B. M. (1996). <i>Self-efficacy in geriatric rehabilitation</i> . Retrieved from <a href="http://search.ebscohost.com/login.aspx?direct=true&amp;db=c8h&amp;AN=2000013812&amp;site=ehost-live">http://search.ebscohost.com/login.aspx?direct=true&amp;db=c8h&amp;AN=2000013812&amp;site=ehost-live</a>	Irrelevant studiedesign
Richards, S. H., Coast, J., Gunnell, D. J., Peters, T. J., Pounsford, J., & Darlow, M. A. (1998). Randomised controlled trial comparing effectiveness and acceptability of an early discharge, hospital at home scheme with acute hospital care. <i>BMJ</i> , 316(7147), 1796-1801.	Irrelevant population
Rideout, E. (2004). Review: comprehensive discharge planning plus post-discharge support reduced total readmissions in older patients with congestive heart failure. <i>Evidence Based Nursing</i> , 7(4), 115-115.	Övrigt
Riordan J, Dierking L, Clements J. Client outcomes of a pilot acute geriatric unit. <i>Kansas Nurse</i> . 1993;68(5):5-.	Irrelevant studiedesign
Rizzo JA, Bogardus ST, Jr, Leo-Summers L, Williams CS, Acampora D, et al. Multicomponent targeted intervention to prevent delirium in hospitalized older patients: what is the economic value? <i>Med Care</i> . 2001;39(7):740-52.	Irrelevant intervention
Robinson S, Rich C, Weitzel T, Vollmer C, Eden B. Delirium prevention for cognitive, sensory, and mobility impairments. <i>Res Theory Nurs Pract</i> . 2008;22(2):103-13.	Irrelevant population
Roder F, Schwab M, Aleker T, Morike K, Thon K, Klotz U. Proximal femur fracture in older patients -- rehabilitation and clinical outcome. <i>Age &amp; Ageing</i> . 2003;32(1):74-80.	Irrelevant jämförelse-alternativ
Rolland, Y., Pillard, F., Lauwers-Cances, V., Busquere, F., Vellas, B., & Lafont, C. (2004). Rehabilitation outcome of elderly patients with hip fracture and cognitive impairment. <i>Disability and Rehabilitation</i> , 26(7), 425-431.	Irrelevant population
Rollins, G. (2001). Coordinated care reduces functional declines in hospitalized elderly patients. <i>Rep Med Guidel Outcomes Res</i> , 12(2), 1-2, 5.	Övrigt
Rollins, G. (2004). Specialized nurse intervention improves post-hospitalization outcomes in elderly heart failure patients. <i>Rep Med Guidel Outcomes Res</i> , 15(12), 9-10, 12.	Irrelevant studiedesign
Ross, D. M. (2008). <i>Healthcare transitions and the aging population: a framework to measure the value of rapid rehabilitation</i> . Retrieved from <a href="http://search.ebscohost.com/login.aspx?direct=true&amp;db=c8h&amp;AN=2010154951&amp;site=ehost-live">http://search.ebscohost.com/login.aspx?direct=true&amp;db=c8h&amp;AN=2010154951&amp;site=ehost-live</a>	Irrelevant studiedesign
Rosswurm MA, Lanham DM. Discharge planning for elderly patients. <i>J Gerontol Nurs</i> . 1998;24(5):14-21.	Irrelevant population

Roughead, E. E., Barratt, J. D., Ramsay, E., Pratt, N., Ryan, P., Peck, R., . . . Gilbert, A. L. (2009). The effectiveness of collaborative medicine reviews in delaying time to next hospitalization for patients with heart failure in the practice setting: results of a cohort study. <i>Circ Heart Fail</i> , 2(5), 424-428.	Irrelevant population
Roughead, E. E., Barratt, J. D., Ramsay, E., Pratt, N., Ryan, P., Peck, R., . . . Gilbert, A. L. (2011). Collaborative home medicines review delays time to next hospitalization for warfarin associated bleeding in Australian war veterans. <i>Journal of Clinical Pharmacy &amp; Therapeutics</i> , 36(1), 27-32.	Irrelevant intervention
Rowe, J. (2002). Care Awaiting Placement Unit: the first 12 months. <i>Geraction</i> , 20(4), 9-14.	Irrelevant studiedesign
Rubenstein Laurence Z, et al. The Sepulveda VA Geriatric Evaluation Unit: Data on four-year outcomes and predictors of improved patient outcomes. <i>Journal of the American Geriatrics Society</i> . 1984;32(7):503-12.	Irrelevant intervention
Rubenstein LZ, Alessi CA, Josephson KR, Hoyl MT, Harker JO, Pietruszka FM. A randomized trial of a screening, case finding, and referral system for older veterans in primary care. <i>Journal of the American Geriatrics Society</i> . 2007;55(2):166-74.	Irrelevant population
Rubin FH, Neal K, Fenlon K, Hassan S, Inouye SK. Sustainability and scalability of the hospital elder life program at a community hospital. <i>J Am Geriatr Soc</i> . 2011;59(2):359-65.	Irrelevant studiedesign
Runciman P, Currie CT, Nicol M, Green L, McKay V. Discharge of elderly people from an accident and emergency department: evaluation of health visitor follow-up. <i>J Adv Nurs</i> . 1996;24(4):711-8.	Irrelevant population
Rypkema G, Adang E, Dicke H, Naber T, de S, Disselhorst L, et al. Cost-effectiveness of an interdisciplinary intervention in geriatric inpatients to prevent malnutrition. <i>J Nutr Health Aging</i> . 2004;8(2):122-7.	Irrelevant population
Sacanella E, Perez-Castejon JM, Nicolas JM, Masanes F, Navarro M, Castro P, et al. Functional status and quality of life 12 months after discharge from a medical ICU in healthy elderly patients: a prospective observational study. <i>Crit Care</i> . 2011;15(2):R105.	Irrelevant studiedesign
Sacanella E, Perez-Castejon JM, Nicolas JM, Masanes F, Navarro M, Castro P, et al. Mortality in healthy elderly patients after ICU admission. <i>Intensive Care Medicine</i> . 2009;35(3):550-5.	Irrelevant studiedesign
Sager, M. A., Rudberg, M. A., Jalaluddin, M., Franke, T., Inouye, S. K., Seth, L., . . . Winograd, C. H. (1996). Hospital Admission Risk Profile (HARP): Identifying older patients at risk for functional decline following acute medical illness and hospitalization. <i>Journal of the American Geriatrics Society</i> , 44(3), 251-257.	Irrelevant studiedesign
Salvi, F., Morichi, V., Grilli, A., Giorgi, R., Spazzafumo, L., Polonara, S., . . . Dessí-Fulgheri, P. (2008). A geriatric emergency service for acutely ill elderly patients: Pattern of use and comparison with a conventional emergency department in Italy. <i>Journal of the American Geriatrics Society</i> , 56(11), 2131-2138.	Irrelevant studiedesign
Sampson, E. L., Jones, L., Thune-Boyle, I. C., Kukkastenvehmaa, R., King, M., Leurent, B., . . . Blanchard, M. R. (2011). Palliative assessment and advance care planning in severe dementia: An exploratory randomized controlled trial of a complex intervention. <i>Palliative Medicine</i> , 25(3), 197-209.	Irrelevant studiedesign
Sankaran, S., Kenealy, T., Adair, A., Adair, V., Coster, H., Whitehead, N., . . . Rea, H. (2010). A complex intervention to support 'rest home' care: a pilot study. <i>N Z Med J</i> , 123(1308), 41-53.	Irrelevant population
Santamaria, N., Houghton, L., Kimmel, L., & Graham, A. (2003). Clinical pathways for fractured neck of femur: a cohort study of health related quality of life, patient satisfaction and clinical outcome. <i>Australian Journal of Advanced Nursing</i> , 20(3), 24-29.	Irrelevant population

Schmader KE, Hanlon JT, Pieper CF, Sloane R, Ruby CM, Twersky J, et al. Effects of geriatric evaluation and management on adverse drug reactions and suboptimal prescribing in the frail elderly. <i>Am J Med.</i> 2004;116(6):394-401.	Irrelevant population
Schoenenberger RA, Conzelmann M, Dubach UC, Schwander J. Quality of emergency room triage of medical inpatients to an acute care clinic or chronic health care facilities. <i>J Gen Intern Med.</i> 1992;7(3):321-7.	Irrelevant intervention
Seneff, M. G., Wagner, D., Thompson, D., Honeycutt, C., & Silver, M. R. (2000). The impact of long-term acute-care facilities on the outcome and cost of care for patients undergoing prolonged mechanical ventilation. <i>Crit Care Med,</i> 28(2), 342-350.	Irrelevant studiedesign
Shah, M. N., Clarkson, L., Lerner, E. B., Fairbanks, R. J., McCann, R., & Schneider, S. M. (2006). An emergency medical services program to promote the health of older adults. <i>J Am Geriatr Soc,</i> 54(6), 956-962.	Irrelevant studiedesign
Shamian J, Clarfield AM, Maclean J. A randomized trial of intra-hospital relocation of geriatric patients in a tertiary-care teaching hospital. <i>J Am Geriatr Soc.</i> 1984;32(11):794-800.	Irrelevant population
Shanley C, Sutherland S, Tumeth R, Stott K, Whitmore E. Caring for the older person in the emergency department: the ASET program and the role of the ASET clinical nurse consultant in South Western Sydney, Australia. <i>J Emerg Nurs.</i> 2009;35(2):129-33.	Irrelevant studiedesign
Shaw FE, Bond J, Richardson DA, Dawson P, Steen IN, McKeith IG, et al. Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: randomised controlled trial. <i>BMJ.</i> 2003;326(7380):73.	Irrelevant intervention
Shaw, F. E., Bond, J., Richardson, D. A., Dawson, P., Nicholas, S., McKeith, I. G., & Anne, K. (2003). Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: Randomised controlled trial. <i>British Medical Journal,</i> 326(7380), 73-75.	Irrelevant intervention
Shyu YL, Liang J, Wu C, Su J, Cheng H, Chou S, et al. Two-year effects of interdisciplinary intervention for hip fracture in older Taiwanese. <i>Journal of the American Geriatrics Society.</i> 2010;58(6):1081-9.	Irrelevant population
Silliman RA. Predictors of family caregivers' physical and psychological health following hospitalization of their elders. <i>J Am Geriatr Soc.</i> 1993;41(10):1039-46.	Irrelevant intervention
Silliman Rebecca A, McGarvey Stephen T, Raymond Patricia M, Fretwell Marsha D. 'The Senior Care Study: Does inpatient interdisciplinary geriatric assessment help the family caregivers of acutely ill older patients?': Erratum. <i>Journal of the American Geriatrics Society.</i> 1990;38(10):1163.	Övrigt
Siu AL, Kravitz RL, Keeler E, Hemmerling K, Kington R, Davis JW, et al. Postdischarge geriatric assessment of hospitalized frail elderly patients. <i>Archives of Internal Medicine.</i> 1996;156(1):76-81.	Irrelevant intervention
Sledge, W. H., Brown, K. E., Levine, J. M., Fiellin, D. A., Chawarski, M., White, W. D., & O'Connor, P. (2006). A randomized trial of primary intensive care to reduce hospital admissions in patients with high utilization of inpatient services. <i>Dis Manag,</i> 9(6), 328-338.	Ej levererad
Sletvold O, Helbostad JL, Thingstad P, Taraldsen K, Prestmo A, Lamb SE, et al. Effect of in-hospital comprehensive geriatric assessment (CGA) in older people with hip fracture. The protocol of the Trondheim Hip Fracture Trial. <i>BMC Geriatr.</i> 2011;11:18.	Övrigt
Soejono CH. The impact of 'comprehensive geriatric assessment (CGA)' implementation on the effectiveness and cost (CEA) of healthcare in an acute geriatric ward. <i>Acta Med Indones.</i> 2008;40(1):3-10.	Irrelevant population

Solanki, N. S., Greenwood, J. E., Wagstaff, M. J., & Franchi, B. F. (2012). Length of stay of elderly patients in an acute burns unit: has the pilot appointment of a geriatrician to the service made a difference? <i>J Burn Care Res</i> , 33, e178. doi: 10.1097/BCR.0b013e31823b262b.	Övrigt
Somme D, Andrieux N, Guéröt E, Lahjibi-Paulet H, Lazarovici C, Gisselbrecht M, et al. Loss of autonomy among elderly patients after a stay in a medical intensive care unit (ICU): A randomized study of the benefit of transfer to a geriatric ward. <i>Archives of Gerontology and Geriatrics</i> . 2010;50(3):e36-e40.	Övrigt
Somme D, Lazarovici C, Drame M, Blanc P, Lang PO, Gauvain JB, et al. The geriatric patient: Use of acute geriatrics units in the emergency care of elderly patients in France. <i>Archives of Gerontology and Geriatrics</i> . 2011;52(1):40-5.	Irrelevant studiedesign
Sorbero, M. E., Saul, M. I., Liu, H., & Resnick, N. M. (2012). Are geriatricians more efficient than other physicians at managing inpatient care for elderly patients? <i>J Am Geriatr Soc</i> , 60, 869-876. doi: 10.1111/j.1532-5415.2012.03934.x.	Irrelevant intervention
Spillane, L. L., Lumb, E. W., Cobaugh, D. J., Wilcox, S. R., Clark, J. S., & Schneider, S. M. (1997). Frequent users of the emergency department: Can we intervene? <i>Academic Emergency Medicine</i> , 4(6), 574-580.	Ej levererad
Spinewine A, Swine C, Dhillon S, Lambert P, Nachega Jean B, Wilmotte L, et al. Effect of a collaborative approach on the quality of prescribing for geriatric inpatients: A randomized, controlled trial. <i>Journal of the American Geriatrics Society</i> . 2007;55(5):658-65.	Irrelevanta effektmått
Srinivasan M, Roffe C. Nutritional status after acute stroke: Is a stroke unit better than a general geriatric ward? <i>Journal of Nutrition, Health and Aging</i> . 2008;12(5):323-5.	Irrelevant intervention
Steele Denise K. Older hospitalized patients who are cared for in a specialized unit called the delirium room. <i>Dissertation Abstracts International: Section B: The Sciences and Engineering</i> . 2009;70(4-B):2183.	Irrelevant studiedesign
Steeman, E., Moons, P., Milisen, K., De, B., De, G., De, F., . . . Abraham, I. (2006). Implementation of discharge management for geriatric patients at risk of readmission or institutionalization. <i>International Journal for Quality in Health Care</i> , 18(5), 352-358.	Irrelevant population
Stenvall M, Olofsson B, Lundstrom M, Englund U, Borssen B, Svensson O, et al. A multidisciplinary, multifactorial intervention program reduces postoperative falls and injuries after femoral neck fracture. <i>Osteoporos Int</i> . 2007;18(2):167-75.	Irrelevant population
Stenvall M, Olofsson B, Nyberg L, Lundstrom M, Gustafson Y. Improved performance in activities of daily living and mobility after a multidisciplinary postoperative rehabilitation in older people with femoral neck fracture: a randomized controlled trial with 1-year follow-up. <i>J Rehabil Med</i> . 2007;39(3):232-8.	Irrelevant population
Stewart DA, Burns JM, Beard K, Dall JL, Lennox IM, Roberts MA, et al. The roles of general and geriatric medicine in the provision of acute medical care for elderly patients. <i>Health Bull (Edinb)</i> . 1992;50(3):259-66.	Irrelevant studiedesign
Stewart M, Suchak N, Scheve A, Popat-Thakkar V, David E, Laquinte J, et al. The impact of a geriatrics evaluation and management unit compared to standard care in a community teaching hospital. <i>Md Med J</i> . 1999;48(2):62-7.	Irrelevant studiedesign
Stineman MG, Ross RN, Hamilton BB, Maislin G, Bates B, Granger CV, et al. Inpatient rehabilitation after stroke: a comparison of lengths of stay and outcomes in the Veterans Affairs and non-Veterans Affairs health care system. <i>Med Care</i> . 2001;39(2):123-37.	Irrelevant studiedesign
Stott DJ, Buttery AK, Bowman A, Agnew R, Burrow K, Mitchell SL, et al. Comprehensive geriatric assessment and home-based rehabilitation for elderly people with a history of recurrent non-elective hospital admissions. <i>Age and Ageing</i> . 2006;35(5):487-91.	Irrelevant intervention

Stromberg L, Ohlen G, Nordin C, Lindgren U, Svensson O. Postoperative mental impairment in hip fracture patients. A randomized study of reorientation measures in 223 patients. <i>Acta Orthop Scand.</i> 1999;70(3):250-5.	Irrelevant population
Sturdy, D. (2010). Best practice for older people in acute care settings. <i>Int J Nurs Stud,</i> 47(1), 1-2.	Övrigt
Styrborn K. Early discharge planning for elderly patients in acute hospitals - An intervention study. <i>Scandinavian Journal of Social Medicine.</i> 1995;23(4):273-85.	Irrelevant studiedesign
Swenson JR, Perez EL. The impact of a geriatric assessment unit on a psychiatric consultation service in a general hospital. <i>Psychiatric Hospital.</i> 1986;17(3):121-5.	Irrelevant intervention
Takun, K., Sahakian, G. D., Bloch, F., Kansao, J., & Dhainaut, J. F. (2006). Health network to improve elderly patients' flow in emergency department. <i>European Journal of Emergency Medicine,</i> 13(5), 260-263.	Irrelevant studiedesign
Teasdale TA, Shuman L, Snow E, Luchi RJ. A comparison of placement outcomes of geriatric cohorts receiving care in a geriatric assessment unit and on general medicine floors. <i>J Am Geriatr Soc.</i> 1983;31(9):529-34.	Irrelevant population
Tibaldi, V., Aimonino, N., Ponzetto, M., Stasi, M. F., Amati, D., Raspo, S., . . . Fabris, F. (2004). A randomized controlled trial of a home hospital intervention for frail elderly demented patients: behavioral disturbances and caregiver's stress. <i>Arch Gerontol Geriatr Suppl</i> (9), 431-436.	Övrigt
Tousignant, M., Boissy, P., Corriveau, H., & Moffet, H. (2006). In home telerehabilitation for older adults after discharge from an acute hospital or rehabilitation unit: A proof-of-concept study and costs estimation. <i>Disability and Rehabilitation: Assistive Technology,</i> 1(4), 209-216.	Irrelevant studiedesign
Traissac, T., Videau, M. N., Bourdil, M. J., Bourdel-Marchasson, I., & Salles, N. (2011). The short mean length of stay of post-emergency geriatric units is associated with the rate of early readmission in frail elderly. <i>Aging - Clinical and Experimental Research,</i> 23, 217-222.	Irrelevant studiedesign
Trella RS. A multidisciplinary approach to case management of frail, hospitalized older adults. <i>Journal of Nursing Administration.</i> 1993;23(2):20-6.	Irrelevant studiedesign
Wald HL, Glasheen JJ, Guerrasio J, Youngwerth JM, Cumbler EU. Evaluation of a hospitalist-run acute care for the elderly service. <i>J Hosp Med.</i> 2011;6(6):313-21.	Irrelevant studiedesign Irrelevant intervention
Walther, S. M., & Jonasson, U. (2004). Outcome of the elderly critically ill after intensive care in an era of cost containment. <i>Acta Anaesthesiologica Scandinavica,</i> 48(4), 417-422.	Irrelevant intervention
Weatherall M. Geriatric evaluation and management in a New Zealand hospital. <i>Australian and New Zealand Journal of Medicine.</i> 2000;30(1):101.	Irrelevant studiedesign
Webster, S., & Harrison, L. (2004). The multidisciplinary approach to mental health crisis management: an Australian example. <i>J Psychiatr Ment Health Nurs,</i> 11(1), 21-29.	Irrelevant population
Wendel, V. I., Durso, S. C., Cayea, D., Arbaje, A. I., & Tanner, E. (2010). Implementing staff nurse geriatric education in the acute hospital setting. <i>Medsurg Nurs,</i> 19(5), 274-280; quiz 281.	Irrelevant population
White SJ, Powers JS, Knight JR, Harrell D, Varnell L, Vaughn C, et al. Effectiveness of an inpatient geriatric service in a university hospital. <i>J Tenn Med Assoc.</i> 1994;87(10):425-8.	Stora brister studiequalitet
Vidán María T, Sánchez E, Alonso M, Montero B, Ortiz J, Serra José A. An intervention integrated into daily clinical practice reduces the incidence of delirium during hospitalization in elderly patients. <i>Journal of the American Geriatrics Society.</i> 2009;57(11):2029-36.	Irrelevant studiedesign

Wiechula, R., Kitson, A., Marcoionni, D., Page, T., Zeitz, K., & Silverston, H. (2009). Improving the fundamentals of care for older people in the acute hospital setting: facilitating practice improvement using a Knowledge Translation Toolkit. <i>Int J Evid Based Healthc</i> , 7(4), 283-295. doi: 10.1111/j.1744-1609.2009.00145.x.	Irrelevant studiedesign
Wilhelmsen K, Duner A, Eklund K, Gosman-Hedstrom G, Blomberg S, Hasson H, et al. Design of a randomized controlled study of a multi-professional and multidimensional intervention targeting frail elderly people. <i>BMC Geriatr</i> . 2011;11(1):24.	Övrigt
Williams ME, Williams TF, Zimmer JG, Hall WJ, Podgorski CA. How does the team approach to outpatient geriatric evaluation compare with traditional care: a report of a randomized controlled trial. <i>J Am Geriatr Soc</i> . 1987;35(12):1071-8.	Irrelevant intervention
Wilson NM, Dawson P. A comparison of primary nursing and team nursing in a geriatric long-term care setting. <i>International Journal of Nursing Studies</i> . 1989;26(1):1-13.	Irrelevant studiedesign
Vind, A. B., Andersen, H. E., Pedersen, K. D., Joergensen, T., & Schwarz, P. (2010). Effect of a program of multifactorial fall prevention on health-related quality of life, functional ability, fear of falling and psychological well-being. A randomized controlled trial. <i>Aging - Clinical and Experimental Research</i> , 22(3), 249-254.	Irrelevant intervention
von, R.-K., Wolfgang, & Krause, T. (2007). Incidence of in-hospital falls in geriatric patients before and after the introduction of an interdisciplinary team-based fall-prevention intervention. <i>Journal of the American Geriatrics Society</i> , 55(12), 2068-2074.	Irrelevant studiedesign
Young, J., Green, J., Forster, A., Small, N., Lawson, K., Bogle, S., . . . Rowe, J. (2007). Postacute care for older people in community hospitals: A multicenter randomized, controlled trial. <i>Journal of the American Geriatrics Society</i> , 55(12), 1995-2002.	Irrelevant intervention
Zelada MA, Salinas R, Baztan JJ. Reduction of functional deterioration during hospitalization in an acute geriatric unit. <i>Arch Gerontol Geriatr</i> . 2009;48(1):35-9.	Irrelevant studiedesign